

**CORR 2001-06**

**Two Theorems on Euclidean Distance Matrices and  
Gale Transform**

**Abdo Y. Alfakih & Henry Wolkowicz**

**Abstract** We present a characterization of those euclidean distance matrices  $D$  which can be expressed as  $D = \lambda(E - C)$  for some nonnegative scalar  $\lambda$  and some correlation matrix  $C$ , where  $E$  is the matrix of all ones. this shows that the cones

$$\text{cone}(E - \mathcal{E}_n) \not\subset \overline{\text{cone}(E - \mathcal{E}_n)} = \mathcal{D}_n,$$

where  $\mathcal{E}_n$  is the ellitope (set of correlation matrices) and  $\mathcal{D}_n$  is the (closed convex) cone of Euclidean distance matrices.

The characterization is given using the Gale transform of the points generating  $D$ . We also show that given points  $p^1, p^2, \dots, p^n \in \mathbb{R}^r$ , for any scalars  $\lambda_1, \lambda_2, \dots, \lambda_n$  such that

$$\sum_{j=1}^n \lambda_j p^j = 0, \quad \sum_{j=1}^n \lambda_j = 0,$$

we have

$$\sum_{j=1}^n \lambda_j \|p^i - p^j\|^2 = \alpha \text{ for all } i = 1, \dots, n,$$

for some scalar *alpha* independent of  $i$ .

**Keywords** Euclidean distance matrices, semidefinite matrices, correlation matrices, tangent cones, Gale transform.

**AMS Subject Classification** 51K05, 15A48, 52B35.