

TODA EQUATIONS AND PIECEWISE POLYNOMIALITY FOR MIXED DOUBLE HURWITZ NUMBERS

I. P. GOULDEN, MATHIEU GUAY-PAQUET, AND JONATHAN NOVAK

ABSTRACT. This article introduces *mixed double Hurwitz numbers*, which interpolate combinatorially between the classical double Hurwitz numbers studied by Okounkov and the monotone double Hurwitz numbers introduced recently by Goulden, Guay-Paquet and Novak. Generalizing a result of Okounkov, we prove that a certain generating series for the mixed double Hurwitz numbers solves the 2-Toda hierarchy of partial differential equations. We also prove that the mixed double Hurwitz numbers are piecewise polynomial, thereby generalizing a result of Goulden, Jackson and Vakil.

1. INTRODUCTION

Consider the right Cayley graph of the symmetric group $S(d)$, as generated by the full conjugacy class of transpositions. This is a $\binom{d}{2}$ -regular graded graph with levels L_0, L_1, \dots, L_{d-1} , where L_k is the set of permutations with $d-k$ cycles. Each level L_k decomposes as the disjoint union of those conjugacy classes in $S(d)$ labelled by Young diagrams with $d-k$ rows.

Let us introduce an edge labelling of the Cayley graph by marking each edge corresponding to the transposition $\tau = (s \ t)$ with t , the larger of the two elements interchanged by τ . This edge labelling was used by Stanley [20] and Biane [1] to study various connections between permutations, parking functions, and noncrossing partitions¹.

Given integers $k, l \geq 0$ and partitions $\alpha, \beta \vdash d$, let $W^{k,l}(\alpha, \beta)$ denote the number of $(k+l)$ -step walks

$$\sigma = \rho \underbrace{(s_1 \ t_1) \dots (s_k \ t_k)}_k \underbrace{(s_{k+1} \ t_{k+1}) \dots (s_{k+l} \ t_{k+l})}_l$$

on the Cayley graph beginning in the conjugacy class C_α and ending in the conjugacy class C_β which satisfy

$$t_1 \leq \dots \leq t_k.$$

In words, this monotonicity condition states that the labels of the edges traversed in the first k steps of the walk form a weakly increasing sequence.

While elementary to define, the numbers $W^{k,l}(\alpha, \beta)$ are related to some rather sophisticated mathematics. Let $z, t, u, a_1, a_2, \dots, b_1, b_2, \dots$ be commuting indeterminates, and form the generating function

¹Stanley and Biane use s as the edge label instead of t , but this is a minor difference.

$$\mathbf{W}(z, t, u, A, B) = 1 + \sum_{d=1}^{\infty} \frac{z^d}{d!} \sum_{k,l=0}^{\infty} t^k \frac{u^l}{l!} \sum_{\alpha, \beta \vdash d} W^{k,l}(\alpha, \beta) p_{\alpha}(A) p_{\beta}(B),$$

where $p_{\alpha}(A)$ and $p_{\beta}(B)$ denote the power-sum symmetric functions in the variables $A = \{a_1, a_2, \dots\}$ and $B = \{b_1, b_2, \dots\}$, respectively. The series

$$\mathbf{H}(z, t, u, A, B) = \log \mathbf{W}(z, t, u, A, B)$$

is a well-defined element of $\mathbb{Q}[[z, t, u, A, B]]$. Set

$$H^{k,l}(\alpha, \beta) = \left[z^d t^k \frac{u^l}{l!} p_{\alpha}(A) p_{\beta}(B) \right] \mathbf{H}(z, t, u, A, B),$$

where $[X]Y$ denotes the coefficient of the term X in a series Y .

The numbers $H^{0,l}(\alpha, \beta)$ were first studied by Okounkov [13], who called them the *double Hurwitz numbers*. By a classical construction due to Hurwitz — the monodromy construction — $H^{0,l}(\alpha, \beta)$ is a weighted count of degree d branched covers of the Riemann sphere by a compact, connected Riemann surface such that the covering map has profile α over 0, β over ∞ , and simple ramification over each of the l th roots of unity. The Riemann-Hurwitz formula determines the genus of the covering surface in terms of the ramification data of the covering map:

$$g = \frac{l + 2 - \ell(\alpha) - \ell(\beta)}{2}.$$

Verifying and extending a conjecture of Pandharipande [18] in Gromov-Witten theory, Okounkov proved that the generating function $\mathbf{H}(z, 0, u, A, B)$ is a solution of the 2-Toda hierarchy of Ueno and Takasaki. The 2-Toda hierarchy is a countable collection of partial differential equations, each of which yields a recurrence relation satisfied by the double Hurwitz numbers. A construction of the Toda hierarchy may be found in [15, §4]. Kazarian and Lando [12] subsequently showed that, when combined with the ELSV formula [4], Okounkov's result yields a streamlined proof of the Kontsevich-Witten theorem relating intersection theory in moduli spaces of curves to integrable hierarchies.

Goulden, Jackson and Vakil [10] gave an alternative interpretation of the double Hurwitz number $H^{0,l}(\alpha, \beta)$ as counting lattice points in a certain integral polytope. As a consequence of this interpretation and Ehrhart's theorem, it was shown in [10] that, after a simple rescaling, $H^{0,l}(\alpha, \beta)$ is a piecewise polynomial function of the parts of α and β , when $\ell(\alpha)$ and $\ell(\beta)$ are held fixed. Detailed structural properties of this piecewise polynomial behaviour were postulated in [10], and subsequently shown to hold by Johnson [11] using the combinatorics of the infinite wedge representation of $\mathfrak{gl}(\infty)$. Shadrin, Spitz and Zvonkine [19] have recently generalized double Hurwitz numbers to the setting of the completed cycle theory introduced by Okounkov and Pandharipande [14]. It is shown in [19] that piecewise polynomiality of double Hurwitz numbers with completed cycle insertions follows from a suitable modification of Johnson's arguments.

The *monotone double Hurwitz numbers* $H^{k,0}(\alpha, \beta)$ were introduced by the present authors in [6], where it was shown that they are the combinatorial objects underlying the asymptotic expansion of the Harish-Chandra-Itzykson-Zuber integral, an important special function in random matrix theory. Structural properties of the monotone single Hurwitz numbers $H^{k,0}(\alpha) = H^{k,0}(\alpha, 1^d)$ were studied in detail in

[7, 8], where it was shown that they enjoy a high degree of structural similarity with the classical single Hurwitz numbers $H^{0,l}(\alpha) = H^{0,l}(\alpha, 1^d)$.

In this article, we extend the theorems of Okounkov and Goulden-Jackson-Vakil to the more general setting of the *mixed double Hurwitz numbers* $H^{k,l}(\alpha, \beta)$, which interpolate between the classical double Hurwitz numbers ($k = 0$) and the monotone double Hurwitz numbers ($l = 0$).

Theorem 1. *The generating function \mathbf{H} is a solution of the 2-Toda hierarchy.*

Theorem 2. *The mixed double Hurwitz numbers are piecewise polynomial.*

2. TODA EQUATIONS

Let us group the transposition generators of $S(d)$ into a matrix,

$$T = \begin{bmatrix} (1\ 2) & (1\ 3) & \dots & (1\ d) \\ & (2\ 3) & \dots & (2\ d) \\ & & \ddots & \vdots \\ & & & (d-1\ d) \end{bmatrix}.$$

Denote by

$$\begin{aligned} J_2 &= (1\ 2) \\ J_3 &= (1\ 3) + (2\ 3) \\ &\vdots \\ J_d &= (1\ d) + (2\ d) + \dots + (d-1\ d), \end{aligned}$$

the column sums of this matrix, viewed as elements of the group algebra $\mathbb{Q}S(d)$. These elements commute. Set $J_1 := 0$, and introduce the multiset

$$\Xi_d = \{\{J_1, \dots, J_d, 0, 0, \dots\}\}.$$

Let Λ denote the \mathbb{Q} -algebra of symmetric functions. We will consider the evaluation of the complete symmetric function $h_\mu \in \Lambda$ indexed by the (k, l) -hook Young diagram $\mu = (k, 1^l)$ on the alphabet Ξ_d . From the definition of the complete symmetric functions, we have

$$\begin{aligned} h_{(k, 1^l)}(\Xi_d) &= h_k(\Xi_d) h_1(\Xi_d)^l \\ &= \left(\sum_{2 \leq t_1 \leq \dots \leq t_k \leq d} J_{t_1} \dots J_{t_k} \right) \left(\sum_{t=2}^d J_t \right)^l \\ &= \left(\sum_{2 \leq t_1 \leq \dots \leq t_k \leq d} \left(\sum_{s_1 < t_1} (s_1\ t_1) \right) \dots \left(\sum_{s_k < t_k} (s_k\ t_k) \right) \right) \left(\sum_{2 \leq t \leq d} \sum_{s < t} (s\ t) \right)^l \\ &= \sum_{\substack{t_1, \dots, t_{k+l}=2 \\ t_1 \leq \dots \leq t_k \\ s_i < t_i}}^d (s_1\ t_1) \dots (s_k\ t_k) (s_{k+1}\ t_{k+1}) \dots (s_{k+l}\ t_{k+l}). \end{aligned}$$

Thus

$$W^{k,l}(\alpha, \beta) = [C_{(1^d)}]C_\alpha h_{(k,1^l)}(\Xi_d)C_\beta,$$

where we have identified each conjugacy class in $S(d)$ with the formal sum of its elements in $\mathbb{Q}S(d)$. In other words, $W^{k,l}(\alpha, \beta)$ is the normalized character of $C_\alpha h_{(k,1^l)}(\Xi_d)C_\beta$ in the regular representation of $\mathbb{Q}S(d)$.

The columns sums of T are known as the *Jucys-Murphy elements* of $\mathbb{Q}S(d)$. It was observed by Jucys and Murphy (see [3] for a proof) that $e_r(\Xi_d)$, the r th elementary symmetric function evaluated on the alphabet of Jucys-Murphy elements, is precisely the sum of the permutations on level L_r of the Cayley graph:

$$e_r(\Xi_d) = \sum_{\substack{\mu \vdash d \\ \ell(\mu) = d-r}} C_\mu.$$

In particular, $e_r(\Xi_d)$ belongs to the center $Z(d)$ of $\mathbb{Q}S(d)$. Since $\Lambda = \mathbb{Q}[e_1, e_2, \dots]$, the substitution $f \mapsto f(\Xi_d)$ defines a specialization $\Lambda \rightarrow Z(d)$. In fact, since the levels of the Cayley graph generate $Z(d)$, this specialization is surjective [5].

Since $C_\alpha h_{(k,1^l)}(\Xi_d)C_\beta$ belongs to the centre of $\mathbb{Q}S(d)$, we can calculate its character in the regular representation using the Fourier transform. Let $(V^\lambda, \rho^\lambda)$, $\lambda \vdash d$, be pairwise non-isomorphic irreducible representations of $\mathbb{Q}S(d)$, so that the map

$$\sigma \mapsto (\rho^\lambda(\sigma) : \lambda \vdash d)$$

defines an algebra isomorphism

$$\mathbb{Q}S(d) \rightarrow \bigoplus_{\lambda \vdash d} \text{End } V^\lambda.$$

The normalized character of $C_\alpha h_{(k,1^l)}(\Xi_d)C_\beta$ in the regular representation of $\mathbb{Q}S(d)$ is thus

$$\begin{aligned} [C_{(1^d)}]C_\alpha h_{(k,1^l)}(\Xi_d)C_\beta &= \sum_{\lambda \vdash d} \text{Tr } \rho^\lambda(C_\alpha h_{(k,1^l)}(\Xi_d)C_\beta) \frac{\dim V^\lambda}{d!} \\ &= \sum_{\lambda \vdash d} \text{Tr } \rho^\lambda(C_\alpha) \rho^\lambda(h_{(k,1^l)}(\Xi_d)) \rho^\lambda(C_\beta) \frac{\dim V^\lambda}{d!} \\ &= \sum_{\lambda \vdash d} \text{Tr } \omega^\lambda(C_\alpha) \omega^\lambda(h_{(k,1^l)}(\Xi_d)) \omega^\lambda(C_\beta) \frac{(\dim V^\lambda)^2}{d!}, \end{aligned}$$

where, for any $C \in Z(d)$, we denote by $\omega^\lambda(C)$ the unique eigenvalue of the scalar operator $\rho^\lambda(C) \in \text{End } V^\lambda$, i.e. $\rho^\lambda(C) = \omega^\lambda(C)I_{V^\lambda}$. The eigenvalue $\omega^\lambda(C)$ is known as the *central character* of C in the representation $(V^\lambda, \rho^\lambda)$.

The central character of any conjugacy class C_μ is given, in terms of the usual character

$$\chi_\mu^\lambda = \text{Tr } \rho^\lambda(\sigma), \quad \sigma \in C_\mu,$$

by the formula

$$\omega^\lambda(C_\mu) = |C_\mu| \frac{\chi_\mu^\lambda}{\dim V^\lambda}.$$

The central character of any symmetric function f evaluated on Ξ_d is simply

$$\omega^\lambda(f(\Xi_d)) = f(\text{Cont}_\lambda),$$

the evaluation of f on the multiset of contents of the Young diagram λ . This remarkable result is due to Jucys and Murphy, see [3] for a proof.

Recalling that the Schur functions have the expansion

$$s_\lambda = \sum_{\mu \vdash d} \frac{|C_\mu|}{d!} \chi_\mu^\lambda p_\mu,$$

where $\lambda \vdash d$, the generating function \mathbf{W} may be rewritten as follows:

$$\begin{aligned} \mathbf{W}(z, t, u, A, B) &= 1 + \sum_{d=1}^{\infty} \frac{z^d}{d!} \sum_{k,l=0}^{\infty} t^k \frac{u^l}{l!} \sum_{\alpha, \beta, \lambda \vdash d} W^{k,l}(\alpha, \beta) p_\alpha(A) p_\beta(B) \\ &= 1 + \sum_{d=1}^{\infty} \frac{z^d}{d!} \sum_{k,l=0}^{\infty} t^k \frac{u^l}{l!} \sum_{\alpha, \beta, \lambda \vdash d} \left(\frac{1}{d!} \sum_{\lambda \vdash d} |C_\alpha| \chi_\alpha^\lambda h_{(k,1^l)}(\text{Cont}_\lambda) |C_\beta| \chi_\beta^\lambda \right) p_\alpha(A) p_\beta(B) \\ &= 1 + \sum_{d=1}^{\infty} z^d \sum_{k,l=0}^{\infty} t^k \frac{u^l}{l!} \sum_{\lambda \vdash d} h_{(k,1^l)}(\text{Cont}_\lambda) \left(\sum_{\alpha \vdash d} \frac{|C_\alpha|}{d!} \chi_\alpha^\lambda p_\alpha(A) \right) \left(\sum_{\beta \vdash d} \frac{|C_\beta|}{d!} \chi_\beta^\lambda p_\beta(B) \right) \\ &= 1 + \sum_{d=1}^{\infty} z^d \sum_{k,l=0}^{\infty} t^k \frac{u^l}{l!} \sum_{\lambda \vdash d} h_{(k,1^l)}(\text{Cont}_\lambda) s_\lambda(A) s_\lambda(B) \\ &= 1 + \sum_{d=1}^{\infty} z^d \sum_{\lambda \vdash d} s_\lambda(A) s_\lambda(B) \left(\sum_{k=0}^{\infty} t^k h_k(\text{Cont}_\lambda) \right) \left(\sum_{l=0}^{\infty} \frac{u^l}{l!} h_1(\text{Cont}_\lambda)^l \right) \\ &= \sum_{\lambda \in \mathcal{Y}} Y(\lambda) s_\lambda(A) s_\lambda(B). \end{aligned}$$

Here \mathcal{Y} is the set of all Young diagrams (including the empty diagram) and

$$Y(\lambda) = Y(\lambda; z, t, u) = \prod_{\square \in \lambda} \frac{z e^{c(\square)u}}{1 - c(\square)t},$$

where for any Young diagram λ and cell \square in λ , $c(\square)$ denotes the content of this cell, i.e. its column index less its row index. By convention, an empty product equals 1.

In order to complete the proof of Theorem 1, we appeal to the following result of Orlov and Scherbin [17], and Carrell [2].

Theorem 3. *Let $\{y_k : k \in \mathbb{Z}\}$ be a set of variables indexed by the integers, and set*

$$Y(\lambda) = \prod_{\square \in \lambda} y_{c(\square)}$$

for each $\lambda \in \mathcal{Y}$. The series

$$\log \left(\sum_{\lambda \in \mathcal{Y}} Y(\lambda) s_\lambda(A) s_\lambda(B) \right)$$

is a solution of the 2-Toda lattice hierarchy in the variables $p_1(A), p_2(A), \dots$ and $p_1(B), p_2(B), \dots$.

Solutions of the Toda equations of the form described in Theorem 3 are known as *diagonal content-product solutions*. Our computations above show that the generating function

$$\mathbf{H}(z, t, u, A, B) = \log \vec{\mathbf{W}}(z, t, u, A, B)$$

of the mixed double Hurwitz numbers is a diagonal content-product solution, with

$$y_k = y_k(z, t, u) = \frac{ze^{ku}}{1-kt}.$$

In particular, Okounkov's generating function $\mathbf{H}(z, 0, u, A, B)$ for the classical double Hurwitz numbers is the diagonal content product solution with

$$y_k = ze^{ku},$$

while the generating function $\mathbf{H}(z, t, 0, A, B)$ for the monotone double Hurwitz numbers is the diagonal content product solution with

$$y_k = \frac{z}{1-kt}.$$

3. PIECEWISE POLYNOMIALITY

To prove Theorem 2, let us begin by formulating precisely the relationship between the numbers $W^{k,l}(\alpha, \beta)$ and $H^{k,l}(\alpha, \beta)$. To this end, consider the generating functions \mathbf{W} and \mathbf{H} as elements of $\mathbb{Q}[[t, u, A, B]][[z]]$ by writing

$$\mathbf{W}(z) = 1 + \sum_{d=1}^{\infty} \frac{z^d}{d!} \mathbf{W}_d(t, u, A, B), \quad \mathbf{H}(z) = \sum_{d=1}^{\infty} \frac{z^d}{d!} \mathbf{H}_d(t, u, A, B),$$

where, for each $d \geq 1$,

$$\begin{aligned} \mathbf{W}_d(t, u, A, B) &= \sum_{k,l=0}^{\infty} t^k \frac{u^l}{l!} \sum_{\alpha, \beta \vdash d} W^{k,l}(\alpha, \beta) p_{\alpha}(A) p_{\beta}(B) \\ \mathbf{H}_d(t, u, A, B) &= d! \sum_{k,l=0}^{\infty} t^k \frac{u^l}{l!} \sum_{\alpha, \beta \vdash d} H^{k,l}(\alpha, \beta) p_{\alpha}(A) p_{\beta}(B). \end{aligned}$$

Then, by the exponential formula, we have

$$\mathbf{W}_d(t, u, A, B) = \sum_{r=1}^d \sum_{P_1 \sqcup \dots \sqcup P_r} \prod_{j=1}^r |P_j|! \mathbf{H}_{|P_j|}(t, u, A, B)$$

for each $d \geq 1$, as an identity in $\mathbb{Q}[[t, u, A, B]]$. In this identity, the inner sum is over partitions $P_1 \sqcup \dots \sqcup P_r$ of $\{1, \dots, d\}$ into r disjoint nonempty sets. Equating the coefficient of $t^k \frac{u^l}{l!}$ on either side of this identity, we obtain

$$\sum_{\alpha, \beta \vdash d} W^{k,l}(\alpha, \beta) p_{\alpha}(A) p_{\beta}(B) = \sum_{r=1}^d \sum_{P_1 \sqcup \dots \sqcup P_r} \prod_{j=1}^r |P_j|! \sum_{\zeta^j, \eta^j \vdash |P_j|} H^{k,l}(\zeta^j, \eta^j) p_{\eta^j}(A) p_{\zeta^j}(B),$$

for each $d \geq 1$ and $k, l \geq 0$, as an identity in $\mathbb{Q}[[A, B]]$. Note that the sum on the right hand side depends only on the overall block structure of the set partition

$P_1 \sqcup \cdots \sqcup P_r$, and not on the internal structure of the individual blocks. We can thus replace the right hand side by a sum over integer partitions,

$$\sum_{\alpha, \beta \vdash d} W^{k,l}(\alpha, \beta) p_\alpha(A) p_\beta(B) = \sum_{\theta \vdash d} c_\theta \sum_{\substack{(\zeta^1, \dots, \zeta^{\ell(\theta)}) \\ \zeta^j \vdash \theta_j}} \sum_{\substack{(\eta^1, \dots, \eta^{\ell(\theta)}) \\ \eta^j \vdash \theta_j}} \left(\prod_{j=1}^{\ell(\theta)} H^{k,l}(\zeta^j, \eta^j) \right) p_{\zeta^1 \cup \dots \cup \zeta^{\ell(\theta)}}(A) p_{\eta^1 \cup \dots \cup \eta^{\ell(\theta)}}(B),$$

where the coefficient c_θ is given by

$$c_\theta = |f^{-1}(\theta)| \prod_{i=1}^{\ell(\theta)} \theta_i!$$

and f denotes the surjection

$$P_1 \sqcup \cdots \sqcup P_r \mapsto \left(\max_{1 \leq j \leq r} |P_j|, \dots, \min_{1 \leq j \leq r} |P_j| \right)$$

from set partitions of $\{1, \dots, d\}$ onto integer partitions of d . The internal sums run over sequences of partitions whose j th element is a partition of the j th part of θ , and if (μ^1, \dots, μ^k) is any sequence of partitions then $\mu^1 \cup \cdots \cup \mu^k$ denotes the partition obtained by arranging the parts of μ^1, \dots, μ^k in weakly decreasing order.

Extracting the coefficient of $p_\alpha(A) p_\beta(B)$ on each side of the above identity in $\mathbb{Q}[[A, B]]$, we obtain the numerical identity

$$W^{k,l}(\alpha, \beta) = \sum_{\theta \vdash d} c_\theta \sum_{\substack{(\zeta^1, \dots, \zeta^{\ell(\theta)}) \\ \zeta^j \vdash \theta_j \\ \zeta^1 \cup \dots \cup \zeta^{\ell(\theta)} = \alpha}} \sum_{\substack{(\eta^1, \dots, \eta^{\ell(\theta)}) \\ \eta^j \vdash \theta_j \\ \eta^1 \cup \dots \cup \eta^{\ell(\theta)} = \beta}} \prod_{j=1}^{\ell(\theta)} H^{k,l}(\zeta^j, \eta^j).$$

The sum on the right hand side receives non-zero contributions from Young diagrams θ whose rows can be obtained by gluing together rows of α , and by gluing together rows of β . This is possible if and only if, for each row θ_k of θ , there exist subsets $I_k, J_k \subseteq \{1, \dots, d\}$ such that

$$(3.1) \quad \sum_{i \in I_k} \alpha_i = \sum_{j \in J_k} \beta_j = \theta_k.$$

In the case of the one-row Young diagram $\theta = (d)$, the required sets are simply $I_1 = J_1 = \{1, \dots, d\}$, so we have

$$W^{k,l}(\alpha, \beta) = d! H^{k,l}(\alpha, \beta) + \dots,$$

where the ellipsis stands for contributions from Young diagrams θ that have at least two rows. These contributions are zero unless the constraint 3.1 is met for each row of θ . We thus conclude that $W^{k,l}(\alpha, \beta)$ and $H^{k,l}(\alpha, \beta)$ typically agree, up to a factor of $d!$.

We can give a geometric interpretation of the above as follows. Fix two positive integers m and n and consider the convex region

$$\mathfrak{R}_{m,n} = \{(x_1, \dots, x_m, y_1, \dots, y_n) : x_1 \geq \cdots \geq x_m > 0, y_1 \geq \cdots \geq y_n > 0, \sum_{i=1}^m x_i = \sum_{j=1}^n y_j\}$$

in Euclidean space \mathbb{R}^{m+n} . Pairs of partitions (α, β) such that

$$|\alpha| = |\beta|, \quad \ell(\alpha) = m, \quad \ell(\beta) = n$$

may be identified with lattice points in $\mathfrak{A}_{m,n}$ via

$$(\alpha, \beta) \mapsto (\alpha_1, \dots, \alpha_m, \beta_1, \dots, \beta_n).$$

Given proper nonempty subsets $I \subset \{1, \dots, m\}$ and $J \subset \{1, \dots, n\}$, define a hyperplane \mathfrak{W}_{IJ} in \mathbb{R}^{m+n} by

$$\mathfrak{W}_{IJ} = \{(x_1, \dots, x_m, y_1, \dots, y_n) \in \mathbb{R}^{m+n} : \sum_{i \in I} x_i = \sum_{j \in J} y_j\}.$$

The hyperplanes \mathfrak{W}_{IJ} , as I ranges over proper nonempty subsets of $\{1, \dots, m\}$ and J ranges over proper nonempty subsets of $\{1, \dots, n\}$, constitute the *resonance arrangement* of [11]. A chamber \mathfrak{c} of the resonance arrangement is a connected component of the complement of a hyperplane \mathfrak{W}_{IJ} in $\mathfrak{A}_{m,n}$. On any chamber \mathfrak{c} of the resonance arrangement, we have that

$$W^{k,l}(\alpha, \beta) = d! H^{k,l}(\alpha, \beta)$$

for all lattice points $(\alpha, \beta) \in \mathfrak{c}$.

Theorem 2 claims that for each chamber \mathfrak{c} there exists a polynomial $p_{\mathfrak{c}}^{k,l}$ in $m+n$ variables such that

$$H^{k,l}(\alpha, \beta) = p_{\mathfrak{c}}^{k,l}(\alpha_1, \dots, \alpha_m, \beta_1, \dots, \beta_n)$$

for all lattice points $(\alpha, \beta) \in \mathfrak{c}$. From the above discussion and Section 2, we know that $H^{k,l}(\alpha, \beta)$ is given by the character formula

$$(3.2) \quad H^{k,l}(\alpha, \beta) = \frac{|C_{\alpha}|}{d!} \frac{|C_{\beta}|}{d!} \sum_{\lambda \vdash d} \chi_{\alpha}^{\lambda} h_{(k,1^l)}(\text{Cont}_{\lambda}) \chi_{\beta}^{\lambda}, \quad d = |\alpha| = |\beta|,$$

on any chamber of the resonance arrangement. In order to deduce Theorem 2 from this character formula, we appeal to a recent result of Shadrin, Spitz and Zvonkine [19] which asserts the piecewise polynomiality of a general class of sums of the above form.

Let $\mathbb{C}^{\mathcal{Y}}$ denote the algebra of all functions $\mathcal{Y} \rightarrow \mathbb{C}$. Following Olshanski [16, Proposition 2.4], we define the algebra \mathbb{A} of *regular functions* on Young diagrams to be the subalgebra of $\mathbb{C}^{\mathcal{Y}}$ generated by the functions

$$\lambda \mapsto f(\text{Cont}_{\lambda}), \quad f \in \mathbb{A},$$

together with the function $\lambda \mapsto |\lambda|$. Consider the transform

$$S : \mathbb{C}^{\mathcal{Y}} \rightarrow \mathbb{C}^{\mathcal{Y} \times \mathcal{Y}}$$

from functions on Young diagrams to functions on pairs of Young diagrams defined by

$$S^f(\alpha, \beta) = \frac{|C_{\alpha}|}{d!} \frac{|C_{\beta}|}{d!} \sum_{\lambda \vdash d} \chi_{\alpha}^{\lambda} f(\lambda) \chi_{\beta}^{\lambda},$$

where $d = |\alpha| = |\beta|$. This definition assumes that α, β have the same size; if $|\alpha| > |\beta|$ or vice versa, complete the smaller diagram by adding unicellular rows. The result of Shadrin, Spitz and Zvonkine that we need may be stated as follows: if $f \in \mathbb{A}$ is a regular function, then S^f is piecewise polynomial with respect to the resonance arrangement.

Theorem 4. *Given a regular function $f \in \mathbb{A}$, positive integers m and n , and a chamber \mathfrak{c} of the resonance arrangement in $\mathfrak{R}_{m,n}$, there exists a polynomial $p_{\mathfrak{c}}^f$ in $m+n$ variables such that*

$$S^f(\alpha, \beta) = p_{\mathfrak{c}}^f(\alpha_1, \dots, \alpha_m, \beta_1, \dots, \beta_n)$$

for all lattice points $(\alpha, \beta) \in \mathfrak{c}$.

From the character formula 3.2, we see that Theorem 2 follows by applying Theorem 4 with f the regular function

$$\lambda \mapsto h_{(k, 1^l)}(\text{Cont}_{\lambda}).$$

REFERENCES

1. P. Biane, *Parking functions of types A and B*, Electron. J. Combin. **9** (2002), #N7.
2. S. R. Carrell, *Diagonal solutions to the 2-Toda hierarchy*, Math. Res. Lett., to appear.
3. P. Diaconis, C. Greene, *Applications of Murphy's elements*, Stanford University Technical Report No. 335, Sept. 1989.
4. T. Ekedahl, S. Lando, M. Shapiro, A. Vainshtein, *Hurwitz numbers and intersections on moduli spaces of curves*, Invent. Math. **146** (2001), 297-327.
5. H. K. Farahat, G. Higman, *The centres of symmetric group rings*, Proc. R. Soc. Lond. Ser. A Math. Phys. Eng. Sci. **250**(1261) (1959), 212-221.
6. I. P. Goulden, M. Guay-Paquet, J. Novak, *Monotone Hurwitz numbers and the HCIZ integral*, arXiv:1107.1015.
7. I. P. Goulden, M. Guay-Paquet, J. Novak, *Monotone Hurwitz in genus zero*, Canad. J. Math, in press.
8. I. P. Goulden, M. Guay-Paquet, J. Novak, *Polynomiality of monotone Hurwitz in higher genera*, Adv. Math. **238** (2013), 1-23.
9. I. P. Goulden, D. M. Jackson, *The KP hierarchy, branched covers, and triangulations*, Adv. Math. **219** (2008), 932-951.
10. I. P. Goulden, D. M. Jackson, R. Vakil, *Towards the geometry of double Hurwitz numbers*, Adv. Math. **198** (2005), 43-92.
11. P. Johnson, *Double Hurwitz numbers via the infinite wedge*, arXiv:1008.3266.
12. M. Kazarian, S. Lando, *An algebro-geometric proof of Witten's conjecture*, J. Amer. Math. Soc. **20**(4) (2007), 1079-1089.
13. A. Okounkov, *Toda equations for Hurwitz numbers*, Math. Res. Lett. **7** (2000), 447-453.
14. A. Okounkov, R. Pandharipande, *Gromov-Witten theory, Hurwitz theory, and completed cycles*, Ann. Math. **163** (2006), 517-560.
15. A. Okounkov, R. Pandharipande, *The equivariant Gromov-Witten theory of \mathbf{P}^1* , Ann. Math. **163** (2006), 561-605.
16. G. Olshanski, *Plancherel averages: remarks on a paper by Stanley*, Electron. J. Combin. **17** (2010), #R43.
17. A. Yu. Orlov, D. M. Scherbin, *Hypergeometric solutions of soliton equations*, Theoretical and Mathematical Physics **128**(1) (2001), 906-926.
18. R. Pandharipande, *The Toda equations and the Gromov-Witten theory of the Riemann sphere*, Lett. Math. Phys. **53** (2000), 59-74.
19. S. Shadrin, L. Spitz, D. Zvonkine, *On double Hurwitz numbers with completed cycles*, J. London Math. Soc., to appear.
20. R. P. Stanley, *Parking functions and noncrossing partitions*, Electron. J. Combin. **4**(2) (1997), #R20.

DEPARTMENT OF COMBINATORICS & OPTIMIZATION, UNIVERSITY OF WATERLOO, CANADA
E-mail address: `ipgoulden@uwaterloo.ca`

LACIM, UNIVERSITÉ DU QUÉBEC À MONTRÉAL, CANADA
E-mail address: `mathieu.guaypaquet@lacim.ca`

DEPARTMENT OF MATHEMATICS, MASSACHUSETTS INSTITUTE OF TECHNOLOGY, USA
E-mail address: `jnovak@math.mit.edu`