

The Moduli Space of Curves, Double Hurwitz Numbers, and Faber’s Intersection Number Conjecture*

I. P. Goulden¹, D. M. Jackson¹, and R. Vakil²

¹Department of Combinatorics and Optimization, University of Waterloo, Waterloo, Ontario N2L 3G1, Canada

{ipgoulden, dmjackson}@math.uwaterloo.ca

²Department of Mathematics, Stanford University, 450 Serra Mall, Stanford, CA 94305, USA
vakil@math.stanford.edu

Received January 28, 2009

Mathematics Subject Classification: 14H10, 05E99, 14K30

Abstract. We define the dimension $2g - 1$ Faber-Hurwitz Chow/homology classes on the moduli space of curves, parametrizing curves expressible as branched covers of \mathbb{P}^1 with given ramification over ∞ and sufficiently many fixed ramification points elsewhere. Degeneration of the target and judicious localization expresses such classes in terms of localization trees weighted by “top intersections” of tautological classes and genus 0 double Hurwitz numbers. This identity of generating series can be inverted, yielding a “combinatorialization” of top intersections of ψ -classes. As genus 0 double Hurwitz numbers with at most 3 parts over ∞ are well understood, we obtain Faber’s Intersection Number Conjecture for up to 3 parts, and an approach to the Conjecture in general (bypassing the Virasoro Conjecture). We also recover other geometric results in a unified manner, including Looijenga’s theorem, the socle theorem for curves with rational tails, and the hyperelliptic locus in terms of κ_{g-2} .

Keywords: moduli space of curves, branched covers, Chow rings, enumerative geometry

Part I. Introduction and Summary of Results

Since we shall be using arguments from geometry and combinatorics, we have separated the material into three parts to assist the reader. Part I gives the background to the topic and a summary of our results. Part II contains the geometry that uses degeneration to obtain a recursion for the Faber-Hurwitz classes, and localization to express these as tree sums involving the Faber symbol. Part III contains an approach through algebraic combinatorics to transform and then solve the formal partial differential equations and functional equations that originate from degeneration and localization in Part II and thence to obtain the top intersection numbers. We have sought to make the transition from the geometry of Part II to the combinatorics of Part III pellucid.

* The first two authors are partially supported by NSERC grants. The third author is partially supported by NSF PECASE/CAREER grant DMS-0238532.

1. Summary of Results

The purpose of this paper is to give a geometrico-combinatorial approach that is direct and, we hope, enlightening, to the three known results that are listed below. We give a summary of results for those quite familiar with moduli spaces of curves, and Faber’s foundational conjectures on their cohomology or Chow rings. A more detailed introduction to this paper is given in Section 2, and most readers should turn immediately to this.

The three results are:

- (I) $R_{2g-1}(\mathcal{M}_{g,n}^{rt})$ is generated by a single element, which we denote $\mathbb{G}_{g,1}$ (Theorem 3.12). This argument was promised in [26, Sec. 5.7]. (Since $R_{2g-1}(\mathcal{M}_g)$ was shown earlier by Faber to be non-zero [7, Thm. 2], this single element is also non-zero by Remark 2.3(iii) below.)
- (II) A combinatorial description of $\psi_1^{\alpha_1} \cdots \psi_n^{\alpha_n} \in R_{2g-1}(\mathcal{M}_{g,n}^{rt})$ as a multiple of this generator, in terms of genus 0 double Hurwitz numbers (Theorem 3.11). (These intersections of ψ -classes determine all top intersections in the tautological ring, and are the subject of Faber’s Intersection Number Conjecture.)
- (III) Hence a proof of Faber’s Intersection Number Conjecture for up to three points[†] and arbitrary genus (Theorem 2.5).

Past proofs of some of these results are described in Section 1.2. In the above statements we have used the following notation. Let $\mathcal{M}_{g,n}^{rt}$ be the moduli space of n -pointed genus g stable curves with “rational tails”, and $R^*(\mathcal{M}_{g,n}^{rt})$ its tautological ring. Genus 0 double Hurwitz numbers enumerate branched covers of the sphere by another sphere, with branching over 0 and ∞ specified by partitions α and β respectively, and the simplest non-trivial branching over an appropriate number of other given points. They are well understood in the case where one of the partitions has at most three parts [21].

The “one-part analysis” yields as corollaries new proofs of a number of important facts, such as the class of the hyperelliptic locus in \mathcal{M}_g as a multiple of κ_{g-2} . These results are collected as Corollaries 6.3 through 6.6. A direct proof of these consequences by our methods would be much shorter; much of our effort will be to develop techniques to deal with more parts.

We note that Faber’s Intersection Number Conjecture for *bounded genus* involves a finite amount of information, thanks to the string and dilaton equation (Proposition 2.6); for any (reasonably small) genus, this finite amount of information can be directly computed ([9]). (We point out that although Faber’s verification of his conjecture up to genus 21, [10], using work of Pandharipande, involves a finite amount of information, it is a large amount, and is very difficult.) However, Faber’s Intersection Number Conjecture for a *bounded number of points*, the case considered here, involves an infinite amount of information.

The methods here can readily be adapted to deal with a larger number of points. For example, the case $n = 4$ follows from the formulae for genus 0 double Hurwitz

[†] Note added in revision: A proof for the general case has recently been given in [36]. See also math.AG/0611659.

numbers from [21], and a Maple computation. The case $n = 5$ should also be computationally tractable. However, we content ourselves with calculations that could be done by hand, as the route to a natural proof of Faber's Intersection Number Conjecture is clearly not through any hoped-for closed form description of the genus 0 double Hurwitz generating series in general — such series might be expected to get quite complicated. Instead, we hope for a proof using the structure of the double Hurwitz generating series as a whole, and there are indications that this may be tractable. We point out in particular the very recent preprint [42], giving a good description of genus 0 double Hurwitz numbers in general, and a particularly elegant description in many cases.

1.1. Motivation

The ELSV formula provides a remarkable link between Hurwitz numbers (counting branched covers of \mathbb{P}^1 or, combinatorially, transitive factorizations of elements of the symmetric group into transpositions) and the intersection theory on the moduli space of curves. See [5, 6], and [25] for a proof in the context of Gromov-Witten theory. The ELSV formula describes Hurwitz numbers in terms of top intersections of the moduli space of curves. This relation can be “inverted” to prove results on top intersections on the moduli space of curves. This has been used by a large number of authors, including Ekedahl, Kazarian, Lando, Okounkov, Pandharipande, Shadrin, Shapiro, Vainshtein, and Zvonkine, to great effect. Notable examples are the proofs of Witten's Conjecture [30, 39].

This paper relies on the observation that similar methods can be applied to the (*non-compact*) moduli space of *smooth* curves. In this case, localization turns question about top intersections into “genus 0 combinatorics” (involving trees rather than general graphs), which should in principle be simpler than the genus g combinatorics that arises in the case of compactified moduli space. Indeed, we immediately get some results (for example, results **(I)** and **(II)**) from the statement of the analogue of the ELSV formula. Explicitly inverting this ELSV-analogue requires more work, but then our knowledge of genus 0 double Hurwitz numbers with few parts translates directly into Faber's Intersection Number Conjecture with few parts (result **(III)**).

We point out that these techniques of algebraic combinatorics are useful in geometry in a wider context. A further example of their use appears in [22] where we give a direct proof of Getzler and Pandharipande's λ_g -Conjecture (without Gromov-Witten theory), and related techniques have been used by a number of authors mentioned above.

1.2. Background

The one-dimensionality of **(I)** was established in [11], and may also follow from Looijenga [38]. The argument given here can be seen as an extension of the tautological vanishing theorem of [26]. The entire argument is outlined in a few pages in [45, Sec. 7].

Getzler and Pandharipande showed that Faber's Intersection Number Conjecture is a formal consequence of the Virasoro Conjecture for the projective plane ([14]), in fact just the degree 0 part (the “large volume” limit). Givental proved the Virasoro

Conjecture for projective space (and more generally Fano toric manifolds) [15, 16]. Lee and Pandharipande are writing a book [31] giving details. Givental’s result is one of the most important results in Gromov-Witten theory, and is a marvelous feat. However, it seems circuitous to prove the Intersection Number Conjecture by means of the Virasoro Conjecture. The latter is a very heavy instrument which conceals the combinatorial structure that lies behind the intersection numbers. As noted by Liu and Xu [35], it is very desirable to have a shorter and direct explanation. (Liu and Xu show how the conjecture cleanly follows from another attractive conjectural identity.) For this reason, we give such an argument, paralleling our understanding of top intersection numbers on $\overline{\mathcal{M}}_{g,n}$ via Hurwitz numbers.

1.3. Outline of Paper

The strategy of the paper is as follows. We define dimension $2g - 1$ Faber-Hurwitz Chow/homology classes on the moduli space of curves, by considering (“virtually”) branched covers of \mathbb{P}^1 with given ramification over ∞ and sufficiently many fixed ramification points elsewhere. We consider only curves with rational tails, which simplifies the combinatorics dramatically. We use the two most effective techniques of Gromov-Witten theory, degeneration and localization.

- Degeneration of the target yields a recursion for such classes, which we can solve explicitly. The story is analogous to that of genus 0 Hurwitz numbers (which are combinatorially straightforward), *not* genus g Hurwitz numbers.
- Localization expresses such classes in terms of the desired top intersections and genus 0 double Hurwitz numbers. The rational tails constraint forces the resulting localization graphs to be trees.

More precisely, localization expresses Faber-Hurwitz classes as a sum over certain decorated trees (Section 3.10) of linear combinations of top intersections (including those that are the subject of Faber’s conjecture) and genus 0 double Hurwitz numbers. The relation between the top intersections and Faber-Hurwitz classes can be easily seen to be invertible, i.e., the top intersections are linear combinations of Faber-Hurwitz classes (which we already understood *via* degeneration). From this, Looijenga’s Theorem drops out quickly (from Theorem 3.12), for example. The central idea of this paper is that this inversion may be done explicitly: The localization sum gives an expression which readily can be unwound, and from which the top intersections can be extracted. This is done by a change of variables.

Using this approach, we quickly recover various geometric results in a unified manner (Section 6.2). Also, as genus 0 double Hurwitz numbers with at most 3 parts over ∞ are well understood, we obtain Faber’s Intersection Number Conjecture for up to 3 parts, and an approach to the conjecture in general.

In Section 2, we state Faber’s Intersection Number Conjecture and give the geometric and algebraic combinatorics background to our approach. The Faber-Hurwitz classes are defined in Section 3, and we obtain a Join-cut Recursion for them by *degeneration* of the target. This is the first form of the Degeneration Theorem, and the equivalent Join-cut (partial differential) Equation for the Faber-Hurwitz series is given as the second form of this theorem. In addition, using *localization*, we obtain

an expression for the Faber-Hurwitz classes as a weighted sum over (localization) trees, giving the first form of the Localization Tree Theorem. Section 4 addresses the weighted tree sum, using the combinatorics of rooted, labelled trees and exponential generating series in a countable set of indeterminates. This results in the second form of the Localization Tree Theorem, which is purely algebraic, giving an expression for the Faber-Hurwitz series in terms of Faber's intersection numbers and the unique solution to a functional equation. Section 5 introduces the *fundamental transformation*, as the composition of three operators. The first two operators are a symmetrization and an implicit change of variables, which give *polynomials* in the new indeterminates. The third operator restricts these polynomials to “top” terms — those of maximum total degree. Our strategy of proof for Faber's Intersection Number Conjecture is to apply the fundamental transformation to the Localization Tree Theorem and to the Degeneration Theorem, and eliminate the transformed Faber-Hurwitz series. The key to the proof is that only top intersection numbers remain when we do so, and they appear in enough linearly independent equations to uniquely determine them. Section 6 applies the strategy for the first time, to the one-part case (of Faber top intersection numbers) which has some non-trivial geometric consequences. The methodology to that point requires us to consider an equation for each genus g separately, so in Section 7 we refine the methodology by creating generating series in genus. This means that a single generating series equation suffices to prove Faber's Intersection Number Conjecture for each number of parts. In Section 8, we apply this refined methodology to establish Faber's (top) Intersection Number Conjecture for 2 and 3 parts, and we include remarks about the general case.

Appendix 10 is a glossary of notation for the reader's convenience.

2. Introduction

2.1. Geometric Background

We work over the complex numbers. Throughout, the genus g is at least 1. We assume some knowledge of the moduli space of curves. An overview is given in [45], which outlines the necessary background and ends with a sketch of many of the results of this paper; see also [44]. We also assume familiarity with Gromov-Witten theory, in particular the theory of relative stable maps, and with localization on their moduli space (“relative virtual localization”) [26, 33, 34, 37]. An introduction to many of the Gromov-Witten ideas we shall use may be found in [45] and [27]. We prefer to work in the Chow ring A^* rather than the cohomology ring H^{2*} , because our arguments apply in this more refined setting, but there is no loss should the reader wish to work in cohomology. Chow/homology classes will often be written in blackboard bold font (e.g., \mathbb{F}) in order to distinguish them from numbers.

Faber's conjectures on the topology of the moduli space of smooth curves \mathcal{M}_g (given in [7]) are a striking description of the “tautological” part of the cohomology ring, the part of the cohomology ring arising “naturally from geometry”. The “top intersections” in this ring have a particularly remarkable combinatorial structure. We begin by describing these conjectures.

On the moduli space of stable n -pointed genus g stable curves $\overline{\mathcal{M}}_{g,n}$ (or any open

subset thereof), let ψ_i ($1 \leq i \leq n$) be the first Chern class of the line bundle corresponding to the cotangent space of the universal curve at the i th marked point.

We shall denote all forgetful morphisms $\overline{\mathcal{M}}_{g,n} \rightarrow \overline{\mathcal{M}}_{g,n'}$ by π , where the source and target will be clear from the context. For example, if $\pi: \mathcal{M}_{g,1} \rightarrow \mathcal{M}_g$, then the i th Mumford-Morita-Miller “ κ -class” is defined by $\kappa_i := \pi_* \psi_1^{i+1}$.

Given an $(n_1 + 1)$ -pointed curve of genus g_1 , and an $(n_2 + 1)$ -pointed curve of genus g_2 , gluing the first curve to the second along the last point of each yields an $(n_1 + n_2)$ -pointed curve of genus $g_1 + g_2$. This gives a map

$$\overline{\mathcal{M}}_{g_1, n_1+1} \times \overline{\mathcal{M}}_{g_2, n_2+1} \rightarrow \overline{\mathcal{M}}_{g_1+g_2, n_1+n_2}.$$

Similarly, we can take a single $(n + 2)$ -pointed curve of genus g , and glue its last two points together to get an n -pointed curve of genus $g + 1$. This gives a map

$$\overline{\mathcal{M}}_{g, n+2} \rightarrow \overline{\mathcal{M}}_{g+1, n}.$$

We call these two types of maps *gluing morphisms*. We call the forgetful and gluing morphisms the *natural morphisms* between moduli spaces of curves.

2.2. The Tautological Ring

There are many equivalent definitions of the tautological ring in the literature. The following will be convenient for our purposes.

Definition 2.1. ([26, Def. 4.2]) *The system of tautological rings $(R^*(\overline{\mathcal{M}}_{g,n}) \subset A^*(\overline{\mathcal{M}}_{g,n}))_{g,n}$ is the smallest system of \mathbb{Q} -vector spaces closed under pushforwards by the natural morphisms, such that all monomials in ψ_1, \dots, ψ_n lie in $R^*(\overline{\mathcal{M}}_{g,n})$.*

If \mathcal{M} is an open subset of $\overline{\mathcal{M}}_{g,n}$, let $R^*(\mathcal{M}) := R^*(\mathcal{M}_{g,n})|_{\mathcal{M}}$ and $R_j(\mathcal{M}) := R^{\dim \mathcal{M} - j}(\mathcal{M})$. (Here $R^k(\mathcal{M})$ is the codimension k part of $R^*(\mathcal{M})$.) We shall often use the notation R_* instead of R^* , because we wish to think of classes as homology classes.

We take this opportunity to introduce a third sort of tautological class: Let $\mathbb{E}_{g,n}$ be the *Hodge bundle* on $\overline{\mathcal{M}}_{g,n}$. It has rank g , and

$$\pi^* \mathbb{E}_{g,0} = \mathbb{E}_{g,n},$$

where π is the forgetful morphism $\pi: \overline{\mathcal{M}}_{g,n} \rightarrow \overline{\mathcal{M}}_g$. Over a point $[(C, p_1, \dots, p_n)] \in \mathcal{M}_{g,n}$, the fiber of $\mathbb{E}_{g,n}$ is the vector space of differentials on C . The λ -classes are defined by $\lambda_i = c_i(\mathbb{E}_{g,n})$. By the above relation, they behave well with respect to pullback by forgetful morphisms (“ $\pi^* \lambda_i = \lambda_i$ ”). Note that $\lambda_0 = 1$.

It is not hard to show that the tautological ring of the moduli space of *smooth curves* \mathcal{M}_g is generated by the κ -classes, and indeed this is essentially the original definition of $R^*(\mathcal{M}_g)$ in [7].

We now describe three predictions of Faber, namely, the Vanishing Conjecture, the Perfect Pairing Conjecture, and the Intersection Number Conjecture, which is a central subject of this paper.

Vanishing Conjecture. $R^i(\mathcal{M}_g) = 0$ for $i > g - 2$, and $R^{g-2}(\mathcal{M}_g) \cong \mathbb{Q}$. This was proved by Looijenga and Faber. Looijenga’s Theorem [38] is that $R^{g-2}(\mathcal{M}_g)$ is generated as a vector space by a single element (this will also follow from our analysis, see Theorem 3.12). Faber proved the following “non-vanishing theorem”.

Theorem 2.2. (Faber [7, Thm. 2]) $R^{g-2}(\mathcal{M}_g) \neq 0$.

(For other proofs, see [2, Thm. 6.5], and [1, Thm. 0.2], which is based on [3]. More precisely, Faber described a linear functional $\int \cdot \lambda_{g-1} \lambda_g : R^{g-2}(\mathcal{M}_g) \rightarrow \mathbb{Q}$, and showed that the image was non-zero by computing a certain intersection number. Bryan-Pandharipande [2] and later Bertram-Cavalieri-Todorov [1] showed that the image is non-zero by different enlightening computations.)

Perfect Pairing Conjecture. The analogue of Poincaré duality holds: For $0 \leq i \leq g - 2$, the cup product $R^i(\mathcal{M}_g) \times R^{g-2-i}(\mathcal{M}_g) \rightarrow R^{g-2}(\mathcal{M}_g) \cong \mathbb{Q}$ is a perfect pairing. This conjecture is known only in special cases, and is essentially completely open.

Informally, these two conjectures state that “ $R^*(\mathcal{M}_g)$ behaves like the $((p, p)$ -part of the) cohomology ring of a $(g - 2)$ -dimensional complex projective manifold.” They imply that the entire structure of the ring is determined by the top intersections of the κ -classes, i.e., by $\prod \kappa_{d_i}$ (where $\sum_i d_i = g - 2$) in terms of some fixed generator of $R_{2g-1}(\mathcal{M}_g)$.

Faber’s Intersection Number Conjecture. This gives a combinatorial description of these top intersections. We note that this conjecture is useful even without knowing the perfect pairing conjecture; Faber’s algorithm [8] reduces all “top intersections” in the tautological ring to intersections of the form described in his Intersection Number Conjecture.

Faber reformulated his conjecture in the striking form given in Conjecture 2.4. In order to give this reformulation, we review the extension of Faber’s Conjecture to curves with “rational tails” (by Faber and Pandharipande, see for example, [41]). Recall that a nodal genus g nodal curve is said to be a *genus g curve with rational tails* if one component is a smooth curve of genus g , and hence the remaining components are genus 0 (spheres), see Figure 1. Then the dual graph is a tree, a fact which will prove crucial for us. This is the reason that the graphs arising from localization involve just tree combinatorics.




Figure 1: A pointed curve with “rational tails”. Notice that the dual graph is a tree.

Denote the corresponding open subset of $\overline{\mathcal{M}}_{g,n}$ (corresponding to stable curves

with rational tails) by $\mathcal{M}_{g,n}^{rt}$. If $\pi: \overline{\mathcal{M}}_{g,n} \rightarrow \overline{\mathcal{M}}_g$, then $\mathcal{M}_{g,n}^{rt} = \pi^{-1}(\mathcal{M}_g)$. In particular, the forgetful morphism $\mathcal{M}_{g,n}^{rt} \rightarrow \mathcal{M}_{g,n'}$ for $n \geq n'$ is projective, and $\mathcal{M}_{g,1}^{rt} = \mathcal{M}_{g,1}$.

We recall some background.

Remark 2.3. i) $R_j(\mathcal{M}_{g,n}^{rt}) = 0$ for $j < 2g - 1$ and all n ;
 ii) for $n \geq 1$,

$$\pi_*: R_{2g-1}(\mathcal{M}_{g,n}^{rt}) \rightarrow R_{2g-1}(\mathcal{M}_{g,1}^{rt}) = R_{2g-1}(\mathcal{M}_{g,1}) \tag{2.1}$$

(where $\pi: \mathcal{M}_{g,n}^{rt} \rightarrow \mathcal{M}_{g,1}$ is the forgetful morphism) is an isomorphism; and

iii) $\pi_*: R_{2g-1}(\mathcal{M}_{g,1}) \rightarrow R_{2g-1}(\mathcal{M}_g)$ (where $\pi: \mathcal{M}_{g,1} \rightarrow \mathcal{M}_g$ is the forgetful morphism) is surjective.

Statements (i) and (ii) are parts (a) and (b) of [26, Prop. 5.8]. Statement (iii) is immediate from an appropriate definition of the tautological ring, for example, Definition 2.1.

Conjecture 2.4. (Faber’s Intersection Number Conjecture, “ ψ -form”) For any n -tuple of **positive** integers (d_1, \dots, d_n) ,

$$\pi_* \left(\psi_1^{d_1} \cdots \psi_n^{d_n} \right) = \frac{(2g - 3 + n)!(2g - 1)!!}{(2g - 1)! \prod_{j=1}^n (2d_j - 1)!!} \psi_1^{g-1}, \quad \text{for } \sum_i d_i = g - 2 + n, \tag{2.2}$$

where π is the forgetful morphism $\mathcal{M}_{g,n}^{rt} \rightarrow \mathcal{M}_{g,1}$. (Recall that $(2a - 1)!! = 1 \cdot 3 \cdots (2a - 1) = (2a)! / (2^a a!)$.)

Note that Faber’s Intersection Number Conjecture is immediate for the case $n = 1$. We shall prove the following theorem.

Theorem 2.5. *Faber’s Intersection Number Conjecture 2.4 is true for up to 3 points (i.e., for $n \leq 3$).*

We shall need to consider more general intersection numbers involving one λ -class, arising in the virtual localization formula. Motivated by the isomorphism (2.1), and in analogy with the “Witten symbol” used in Gromov-Witten theory (e.g., [27, Sec. 26.2]), define the *Faber symbol* as

$$\langle \tau_{a_1} \cdots \tau_{a_n} \lambda_k \rangle_g^{\text{Fab}} := \pi_* \left(\psi_1^{a_1} \cdots \psi_n^{a_n} \lambda_k \right) \in R_{2g-1}(\mathcal{M}_{g,1}), \tag{2.3}$$

where the pushforward π_* is the isomorphism of (2.1) ($\pi: \mathcal{M}_{g,n}^{rt} \rightarrow \mathcal{M}_{g,1}$ is the forgetful morphism). *Caveat:* This symbol is a Chow class, *not* a number; we repeat that we will often indicate classes using blackboard bold font. The symbol is declared to be zero if some $a_i < 0$, or if the product lies in wrong degree, i.e.,

$$k + \sum_i a_i \neq \dim \mathcal{M}_{g,n}^{rt} - (2g - 1) = g - 2 + n.$$

Note that the case $k = 0$ is equivalent to there being no λ -factor, as $\lambda_0 = 1$. This symbol satisfies a version of the usual string and dilaton equations:

Proposition 2.6. *The following relations among the Faber symbols hold.*

$$\langle \tau_0 \tau_{a_1} \cdots \tau_{a_n} \lambda_k \rangle_g^{\mathbb{F}ab} = \sum_{i=1}^n \langle \tau_{a_1} \cdots \tau_{a_{i-1}} \cdots \tau_{a_n} \lambda_k \rangle_g^{\mathbb{F}ab} \quad (\text{string equation}), \quad (2.4)$$

$$\langle \tau_1 \tau_{a_1} \cdots \tau_{a_n} \lambda_k \rangle_g^{\mathbb{F}ab} = (2g - 2 + n) \langle \tau_{a_1} \cdots \tau_{a_n} \lambda_k \rangle_g^{\mathbb{F}ab} \quad (\text{dilaton equation}). \quad (2.5)$$

The proof is a variation of the proof of the usual string and dilaton equations, and is left as an exercise to the reader (see for example, [45, Sec. 3.13]).

Part II. Geometry

In Part II, we use degeneration and localization to obtain a topological recursion for the Faber-Hurwitz classes, and an expression for the Faber-Hurwitz classes as a sum over a class of weighted trees that are to be defined. The topological recursion is then transformed into a partial differential equation for the Faber-Hurwitz series.

3. Degeneration and Localization

For a partition α , we use $|\alpha|$ and $l(\alpha)$ for the sum of parts and number of parts, respectively, and write $\alpha \vdash |\alpha|$. If α has i_j parts equal to j , $j \geq 1$, then we also write $\alpha = (1^{i_1} 2^{i_2} \cdots)$, where convenient. The set of all non-empty partitions is denoted by \mathcal{P} .

3.1. Relative Stable Maps

We shall use the theory of stable relative maps to \mathbb{P}^1 , following J. Li’s algebro-geometric description in [33], and his description of their deformation-obstruction theory in [34]. (We point out earlier definitions of relative stable maps in the differentiable category due to A.-M. Li and Y. Ruan [32], and Ionel and Parker [28, 29], and Gathmann’s work [13] in the algebraic category in genus 0.) We need the algebraic category for several reasons, most importantly because we shall use virtual localization, and an explicit description of the moduli space’s deformation-obstruction theory.

3.1.1. Relative to One Point

The moduli space of genus g relative stable maps to \mathbb{P}^1 relative to one point ∞ , with branching above ∞ given by the partition $\alpha \vdash d$, is denoted by $\overline{\mathcal{M}}_{g, \alpha}(\mathbb{P}^1)$ where d is the degree of the cover.

A relative map to $X = \mathbb{P}^1$ is the following data:

- A morphism f_1 from a nodal n -pointed genus g curve (C, q_1, \dots, q_n) (where as usual the q_j are distinct non-singular points) to a chain of \mathbb{P}^1 ’s, $T = T_0 \cup T_1 \cup \cdots \cup T_l$ (where T_i and T_{i+1} meet), with a point $\infty \in T_i - T_{i-1}$, so there are two points named ∞ . We will call the one on X , ∞_X , and the one on T , ∞_T , whenever there is any ambiguity.

- A projection $f_2: T \rightarrow X$ contracting T_i to ∞_X (for $i > 0$) and giving an isomorphism from $(T_0, T_0 \cap T_1)$ ((T_0, ∞_T) , respectively) to (X, ∞_X) if $t > 0$ (if $t = 0$, respectively). Denote $f_2 \circ f_1$ by f .
- We have an equality of divisors on C : $f_1^* \infty_T = \sum \alpha_i q_i$. In particular, $f_1^{-1} \infty_T$ consists of non-singular marked points of C .
- The preimage of each node n of T is a union of nodes of C . At any such node n' of C , the two branches map to the two branches of n , and their orders of branching are the same. This is called the *predeformability* or *kissing* condition. See Figure 2 for a pictorial representation. Analytically, this map is of the following form. The node $uv = 0$ in the uv -plane maps to the node $xy = 0$ in the xy -plane by $(u, v) \mapsto (u^m, v^m) = (x, y)$. (The branching of the u -axis over the x -axis is the same as the branching of the v -axis over the y -axis.)




Figure 2: The “predeformability” or “kissing” condition on maps of nodes to nodes. The singularities of the source and target are nodes (analytically isomorphic to $xy = 0$ in \mathbb{C}^2), although it is impossible to depict them as such on the page.

An *isomorphism* of two such maps is a commuting diagram

$$\begin{array}{ccc}
 (C, p_1, \dots, p_m, q_1, \dots, q_n) & \xrightarrow{\sim} & (C', p'_1, \dots, p'_m, q'_1, \dots, q'_n) \\
 f_1 \downarrow & & \downarrow f_1 \\
 (T, \infty_T) & \xrightarrow{\sim} & (T, \infty_T) \\
 f_2 \downarrow & & \downarrow f_2 \\
 (X, \infty_X) & \xrightarrow{=} & (X, \infty_X)
 \end{array}$$

where all horizontal morphisms are isomorphisms, the bottom (although not necessarily the middle) is an equality, the top horizontal isomorphism sends p_i to p'_i and q_j to q'_j . Note that the middle isomorphism must preserve the isomorphism of T_0 with X ,

and is hence the identity on T_0 , but for $i > 0$, the isomorphism may not be the identity on T_i .

We say that f is *stable* if it has a finite automorphism group.

Let r_α^g be the “expected” number of branch points away from ∞ (the number if all the branching over points other than ∞ — “away from ∞ ” — were *simple*, i.e., corresponding to the partition $(1^{d-2}) \vdash d$). Then,

$$r_\alpha^g := d + l(\alpha) + 2g - 2, \tag{3.1}$$

by the Riemann-Hurwitz formula. Thus $\overline{\mathcal{M}}_{g,\alpha}(\mathbb{P}^1)$ supports a virtual fundamental class

$$[\overline{\mathcal{M}}_{g,\alpha}(\mathbb{P}^1)]^{\text{vir}} \in A_{r_\alpha^g}(\overline{\mathcal{M}}_{g,\alpha}(\mathbb{P}^1))$$

of dimension r_α^g .

There is a Fantechi-Pandharipande branch morphism [12]

$$br: \overline{\mathcal{M}}_{g,\alpha}(\mathbb{P}^1) \rightarrow \text{Sym}^{r_\alpha^g}(\mathbb{P}^1) \tag{3.2}$$

sending each map to the set of its branch points in \mathbb{P}^1 , excluding the ones that are “automatic” because of the branching over ∞ . Note that the total branching $C \rightarrow \mathbb{P}^1$ above a point in the target $p \neq \infty \in \mathbb{P}^1$ is 1 only if the map is simply branched above p .

3.1.2. Relative to Two Points

Similarly, let $\overline{\mathcal{M}}_{g,\alpha,\beta}(\mathbb{P}^1)$ be the moduli space of stable relative maps to \mathbb{P}^1 relative to two points 0 and ∞ , where branching above 0 is given by a partition $\alpha \vdash d$ and the branching above ∞ is given by a partition $\beta \vdash d$. By the Riemann-Hurwitz formula, the expected number of branch points away from 0 and ∞ is

$$r_{\alpha,\beta}^g := l(\alpha) + l(\beta) + 2g - 2, \tag{3.3}$$

and the moduli space supports a virtual fundamental class

$$[\overline{\mathcal{M}}_{g,\alpha,\beta}(\mathbb{P}^1)]^{\text{vir}} \in A_{r_{\alpha,\beta}^g}(\overline{\mathcal{M}}_{g,\alpha,\beta}(\mathbb{P}^1))$$

of this dimension. There is also a Fantechi-Pandharipande branch morphism

$$br: \overline{\mathcal{M}}_{g,\alpha,\beta}(\mathbb{P}^1) \rightarrow \text{Sym}^{r_{\alpha,\beta}^g}(\mathbb{P}^1).$$

A mild variation of this is the case of relative stable maps to a non-rigidified target (see [26, Sec. 2.4]; these are sometimes called “rubber maps”), where two relative maps to $(\mathbb{P}^1, 0, \infty)$ are considered the same if they differ by an automorphism of the target \mathbb{P}^1 preserving 0 and ∞ (an element of the group \mathbb{C}^*). This moduli space $\overline{\mathcal{M}}_{g,\alpha,\beta}(\mathbb{P}^1)_\sim$ of such objects supports a virtual fundamental class of dimension one less than that of the “unrigidified” usual space:

$$[\overline{\mathcal{M}}_{g,\alpha,\beta}(\mathbb{P}^1)_\sim]^{\text{vir}} \in A_{r_{\alpha,\beta}^g - 1}(\overline{\mathcal{M}}_{g,\alpha,\beta}(\mathbb{P}^1)_\sim).$$

There is an obvious map $\phi: \overline{\mathcal{M}}_{g,\alpha,\beta}(\mathbb{P}^1) \rightarrow \overline{\mathcal{M}}_{g,\alpha,\beta}(\mathbb{P}^1)_{\sim}$ “forgetting the rigidification of the target”. The two fundamental classes are related by the following result.

Proposition 3.1. ([26, Lem. 4.6])

$$\phi_* \left(br^*(L) \cap \left[\overline{\mathcal{M}}_{g,\alpha,\beta}(\mathbb{P}^1) \right]^{\text{vir}} \right) = r_{\alpha,\beta}^g \left[\overline{\mathcal{M}}_{g,\alpha,\beta}(\mathbb{P}^1)_{\sim} \right]^{\text{vir}}$$

where L is the class in $\text{Sym}^{r_{\alpha,\beta}^g}(\mathbb{P}^1)$ corresponding to those $r_{\alpha,\beta}^g$ -tuples of points containing a given fixed point p_0 .

Intuitively, this is because given any unrigidified map in $\overline{\mathcal{M}}_{g,\alpha,\beta}(\mathbb{P}^1)_{\sim}$, there should be $r_{\alpha,\beta}^g$ ways to “rigidify” it so that a branch point maps to $1 \in \mathbb{P}^1$, as there “should be” $r_{\alpha,\beta}^g$ branch points away from 0 and ∞ .

3.1.3. Relative Stable Maps with Rational Tails

We define the moduli spaces of relative stable maps with rational tails $\mathcal{M}_{g,\alpha}(\mathbb{P}^1)^{rt}$ as the analogous moduli spaces of maps where the source curve is a nodal curve with rational tails. This is an open substack of the space of relative stable maps. We define $\mathcal{M}_{g,\alpha,\beta}(\mathbb{P}^1)^{rt}$ similarly.

3.1.4. Relative Stable Maps with Possibly Disconnected Source

The above definitions of relative maps would all make sense without the requirement that the source curve be connected. The resulting moduli space of maps from “possibly-disconnected” curves to \mathbb{P}^1 relative to one point is denoted by $\overline{\mathcal{M}}_{g,\alpha}(\mathbb{P}^1)_{\bullet}$, and similarly for maps relative to two points.

3.1.5. Observations Using the Riemann-Hurwitz Formula

We make some crucial observations, which are straightforward to verify using the Riemann-Hurwitz formula. We shall use them repeatedly. A degree d cover is said to be *completely branched* over a point if the branching data is $(d) \vdash d$. Define a *trivial cover* of \mathbb{P}^1 to be a map $\mathbb{P}^1 \rightarrow \mathbb{P}^1$ of the form $[x; y] \mapsto [x^u; y^u]$, branched only over 0 and ∞ .

Remark 3.2. Suppose we have a map $C \rightarrow \mathbb{P}^1$ from a nodal (possibly disconnected) curve, unbranched away from 0 and ∞ , where C is non-singular over 0 and ∞ . Then it is a disjoint union of trivial covers.

Remark 3.3. a) Suppose we have a map from a nodal curve C to \mathbb{P}^1 , with no branching away from 0, 1, and ∞ , simple branching at 1, and non-singular over 0 and ∞ . Then it is a union of trivial covers, with one further component, that is non-singular of genus 0, completely branched over one of $\{0, \infty\}$, and with two preimages over the other (see Figure 3). We call this an *almost-trivial cover*.




Figure 3: Almost-trivial covers: All branched covers from non-singular curves with simple branching above 1 and no other branching away from 0 and ∞ must look like this.

b) More generally, suppose we have a map from a curve C to a chain of \mathbb{P}^1 's, satisfying the kissing or predeformability condition, unbranched except for two non-singular points 0 and ∞ on the ends of the chain, and simple branching at one more point. Then the map looks like a number of trivial covers glued together, with one further almost-trivial cover $\mathbb{P}^1 \rightarrow \mathbb{P}^1$ of the sort described in Remark 3.3(a).

Remark 3.4. a) If we have a map from a nodal curve C to \mathbb{P}^1 , with total branching away from 0 and ∞ of degree less than $2g$, and non-singular over 0 and ∞ , then C has no components of geometric genus g . If instead the total branching away from 0 and ∞ is precisely $2g$, and C has a component of geometric genus g , then the cover is a disjoint union of trivial covers, and one connected curve C' of arithmetic genus g , where the map $C' \rightarrow \mathbb{P}^1$ is completely branched over 0 and ∞ .

b) Suppose more generally we have a map from a curve C to a chain of \mathbb{P}^1 's, satisfying the kissing condition, with branching of less than $2g$ away from the nodes and two smooth points 0 and ∞ on the ends of the chain, then C has no component of geometric genus g . If the branching is precisely $2g$ away from the nodes and 0 and ∞ , then the map is a number of trivial covers glued together, plus one other cover of the sort described in Remark 3.4(a).

The following fact will prove essential.

Theorem 3.5. *Let $\mathbb{G}_{g,d,\sim} := \mu_* [\mathcal{M}_{g,(d),(d)}(\mathbb{P}^1, d)_{\sim}]^{\text{vir}}$ where μ is the moduli map to $\mathcal{M}_{g,2}^{\text{rt}}$, remembering the curve and the points over 0 and ∞ . Then $\mathbb{G}_{g,d,\sim} = d^{2g} \mathbb{G}_{g,1,\sim}$. Similarly, if $\mathbb{G}_{g,d} := \mu_* [br^*(L) \cap \mathcal{M}_{g,(d),(d)}(\mathbb{P}^1, d)^{\text{rt}}]^{\text{vir}}$, then $\mathbb{G}_{g,d} = d^{2g} \mathbb{G}_{g,1}$.*

Proof. The second statement follows from the first by Proposition 3.1. We now prove the first.

Let $v: \text{Jac} \rightarrow \mathcal{M}_{g,2}^{\text{rt}}$ be the universal Jacobian or Picard stack over $\mathcal{M}_{g,2}^{\text{rt}}$. Let $\text{Jac}[d] := \ker(\text{Jac} \xrightarrow{\times d} \text{Jac})$ be the d -torsion substack. (Points of $\text{Jac}[d]$ correspond to curves along with a d -torsion point.)

Let p and q be the names of the points of the genus g curve parametrized by $\mathcal{M}_{g,2}^{\text{rt}}$. Let $s_{p-q}: \mathcal{M}_{g,2}^{\text{rt}} \rightarrow \text{Jac}$ be the section corresponding to the line bundle $O(p-q)$. There

is a natural (stack-theoretic) isomorphism

$$\mathcal{M}_{g,(d),(d)}(\mathbb{P}^1)_{\sim}^{rt} \cong \text{Jac}[d] \cap s_{p-q}(\mathcal{M}_{g,2}^{rt}), \tag{3.4}$$

as follows. Given any family of such relative stable maps (with rational tails), let p and q be the preimages of 0 and ∞ , respectively. Note that the target \mathbb{P}^1 never degenerates (or sprouts) in such a family. Thus $O(dp - dq) \cong O$, and hence $O(p - q)$ is indeed d -torsion. Conversely, given any family of curves in $\mathcal{M}_{g,2}^{rt}$ where $O(p - q)$ is d -torsion, we obtain a unique family of relative stable maps with unrigidified target of the desired sort. (It is essential to note that this isomorphism holds over the boundary of $\mathcal{M}_{g,2}^{rt}$ as well.)

$$\begin{array}{ccc} \mathcal{M}_{g,(d),(d)}(\mathbb{P}^1)_{\sim}^{rt} = \text{Jac}[d] \cap s_{p-q}(\mathcal{M}_{g,2}^{rt}) & \longrightarrow & \text{Jac} \\ & \searrow \mu & \downarrow \nu \\ & & \mathcal{M}_{g,2}^{rt} \end{array} \quad \begin{array}{c} \uparrow s_{p-q} \\ \downarrow \nu \end{array}$$

Both sides of the isomorphism (3.4) have natural virtual fundamental classes, the latter by intersecting the section $s_{p-q}(\mathcal{M}_{g,2}^{rt})$ with the local complete intersection $\text{Jac}[d] \hookrightarrow \text{Jac}$. We claim that these virtual fundamental classes agree. (It is easy to show that away from the boundary of $\mathcal{M}_{g,2}^{rt}$, both will agree with the actual fundamental class, but we shall not need this fact.) We prove this by showing that they have the same deformation-obstruction theory over $\mathcal{M}_{g,2}^{rt}$.

We use the description of the deformation-obstruction theory of $\overline{\mathcal{M}}_{g,(d),(d)}(\mathbb{P}^1)$ over $\overline{\mathcal{M}}_{g,2}$ given in [26, Sec. 2.8]. On the locus, at a relative stable map $[f: C \rightarrow \mathbb{P}^1]$, the relative deformation space is identified with $H^0(C, f^*T_{\mathbb{P}^1}(-[0] - [\infty])) = H^0(C, f^*O_{\mathbb{P}^1})$ in [26, equ. (1)]. (The formula given in [26] is more involved, as it applies in more general situations. In particular, in the locus, the target never sprouts, so the notation f^\dagger in [26] agrees with f^* . Also $T_{\mathbb{P}^1}(-\log[0] - \log[\infty]) = T_{\mathbb{P}^1}(-[0] - [\infty])$, as \mathbb{P}^1 has dimension 1.) Thus the relative deformation space is canonically identified with $H^0(C, O_C)$, which is one-dimensional. But such deformations correspond precisely to the \mathbb{C}^* -action induced by its action on the target (keeping the map otherwise fixed), so the relative deformation space of maps to \mathbb{P}^1 with *unrigidified target* is 0.

The relative obstruction space $\text{RelOb}(f)$ at a relative stable map $[f: C \rightarrow \mathbb{P}^1]$ has a natural filtration [26, Equ. (2)]

$$\begin{aligned} 0 \rightarrow H^1(C, f^*T_{\mathbb{P}^1}(-[0] - [\infty])) &\rightarrow \text{RelOb}(f) \\ &\rightarrow H^0(C, f^{-1}\mathcal{E}xt^1(\Omega_{\mathbb{P}^1}(-[0] - [\infty]), O_{\mathbb{P}^1})) \rightarrow 0. \end{aligned}$$

Now $\mathcal{E}xt^1(O_{\mathbb{P}^1}, O_{\mathbb{P}^1}) = 0$, so this filtration gives a natural isomorphism

$$H^1(C, O_C) \xrightarrow{\sim} \text{RelOb}(f).$$

But this is the relative obstruction space of the intersection: The normal bundle to the d -torsion locus $\text{Jac}[d]$ is canonically $H^1(C, O_C)$. Thus we have shown that the isomorphism (3.4) indeed extends to an isomorphism of virtual fundamental classes.

We conclude by noting that $[\text{Jac}[d]] = d^{2g}[\text{Jac}[1]]$ in $A_*(\text{Jac})$. This follows from [4, Sec. 2.15], as observed in the proof of [38, Lem. 2.10]. ■

Remark. $\mathbb{G}_{g,1}$ will be our “natural generator” of $R_{2g-1}(\mathcal{M}_{g,1})$. The generator used by Looijenga in [38] is the hyperelliptic locus $\mathbb{H}yp$. They are related as follows: $(2g + 2)(2g + 1)\mathbb{H}yp = (2^{2g} - 1)\mathbb{G}_{g,1,\sim}$ (both count hyperelliptic covers with two fixed branch points), and $\mathbb{G}_{g,1} = 2g\mathbb{G}_{g,1,\sim}$ from Proposition 3.1; see Corollary 6.5 and its proof.

3.2. Faber-Hurwitz Classes

We define Hurwitz classes, following [26]. Our motivation is as follows. Suppose we are interested in dimension j classes on $\overline{\mathcal{M}}_{g,n}$. One way of producing a family of genus g curves with n marked points is by considering branched covers of \mathbb{P}^1 with fixed branching over ∞ corresponding to a partition $(\alpha_1, \dots, \alpha_n)$ of d with n parts. The curve in question will be the source of the map to \mathbb{P}^1 , and the n points will be the points over ∞ . This Hurwitz scheme (the moduli space of such maps) will have dimension r_α^g . In order to get a class of dimension j , we fix all but j branch points. We can then push forward this class to $\overline{\mathcal{M}}_{g,n}$. The definition of “Hurwitz class” involves doing this “virtually”:

$$\mathbb{H}_j^{g,\alpha} := \mu_* \left(br^*(L)^{r_\alpha^g - j} \cap \left[\overline{\mathcal{M}}_{g,\alpha}(\mathbb{P}^1) \right]^{\text{vir}} \right),$$

where μ is the forgetful map $\overline{\mathcal{M}}_{g,\alpha}(\mathbb{P}^1) \rightarrow \overline{\mathcal{M}}_{g,n}$ and br is the branch morphism (3.2). Here, as in Proposition 3.1, L is the class in $\text{Sym}^{r_\alpha^g}(\mathbb{P}^1)$ corresponding to unordered tuples of points containing a given fixed point p_0 .

Define the *Faber-Hurwitz* class $\mathbb{F}^{g,\alpha}$ by

$$\mathbb{F}^{g,\alpha} := \mathbb{H}^{g,\alpha} \Big|_{\mathcal{M}_{g,n}^{rt}} \in A_{2g-1}(\mathcal{M}_{g,n}^{rt}). \tag{3.5}$$

Equivalently, we can consider the space of relative stable maps with rational tails $\mathcal{M}_{g,\alpha}(\mathbb{P}^1)^{rt}$, along with its virtual fundamental class; fix all but $2g - 1$ branch points; and push forward to the moduli space $\mathcal{M}_{g,n}^{rt}$. Let

$$r_{g,\alpha}^{\text{Fab}} := r_\alpha^g - (2g - 1) = d + l(\alpha) - 1 \tag{3.6}$$

be the number of *fixed* branch points on \mathbb{P}^1 in this construction.

We shall now understand this class in two ways, by degeneration and localization. The first will connect us to a Hurwitz-type problem (and join-cut type recursion) that we can solve. The second will connect us to the tautological ring.

Readers less comfortable with these ideas from Gromov-Witten theory may go directly to the recursions that are obtained by these means without losing the thread of the paper. These recursions are treated quite cleanly through transforms of the corresponding partial differential equations (see Section 4).

3.3. Degeneration of Faber-Hurwitz Classes

We now describe a join-cut type recursive formula for the Faber-Hurwitz classes $\mathbb{F}^{g,\alpha}$. This will allow us to compute $\mathbb{F}^{g,\alpha}$ in terms of the putative generator $\mathbb{G}_{g,1}$, first recursively, and later, in closed form. We use Li's degeneration formula [33, 34], which in our case states that, if we degenerate $[\overline{\mathcal{M}}_{g,\alpha}(\mathbb{P}^1)]_{\bullet}^{\text{vir}}$ by degenerating the target into two components $\mathbb{P}_L^1 \cup \mathbb{P}_R^1$, where the point corresponding to α is on the right component \mathbb{P}_R^1 , then

$$[\overline{\mathcal{M}}_{g,\alpha}(\mathbb{P}^1)]_{\bullet}^{\text{vir}} = \sum_{g_1, g_2, \gamma} \left(\prod_i \gamma_i \right) [\overline{\mathcal{M}}_{g_1, \gamma}(\mathbb{P}^1)]_{\bullet}^{\text{vir}} \boxtimes [\overline{\mathcal{M}}_{g_2, \gamma, \alpha}(\mathbb{P}^1)]_{\bullet}^{\text{vir}}, \quad (3.7)$$

where the sum is over all “splitting of the data”: $|\gamma| = |\alpha|$, $g_1 + g_2 + l(\gamma) - 1 = g$, and the genus g_1 cover C_1 is glued to the genus g_2 cover C_2 over the node $\mathbb{P}_L^1 \cap \mathbb{P}_R^1$ by gluing the point corresponding to γ_i on C_1 to the point corresponding to γ_i on C_2 . There is the obvious variation for $[\overline{\mathcal{M}}_{g,\alpha,\beta}(\mathbb{P}^1)]_{\bullet}^{\text{vir}}$ for stable maps relative to two points. If we are interested in spaces with connected source, i.e., $\overline{\mathcal{M}}_{g,\alpha}(\mathbb{P}^1)$ rather than $\overline{\mathcal{M}}_{g,\alpha}(\mathbb{P}^1)_{\bullet}$, then we include only the summands where the union $C_1 \cup C_2$ is connected. We shall “cap” these equalities with pullbacks under the branch morphism, and interpret them as considering maps with given branch points, where we shall specify how many branch points degenerate to \mathbb{P}_L^1 and \mathbb{P}_R^1 , respectively.

Lemma 3.6. *If $j < 2g - 1$, then the restriction of $\mu_*(\mathbb{H}_j^{g,\alpha})$ to the rational-tails locus $\mathcal{M}_{g,n}^{rt}$ is 0.*

Proof. Degenerate the target \mathbb{P}^1 into a chain of $(r_{\alpha}^g - j)$ \mathbb{P}^1 's, where each of the $r_{\alpha}^g - j$ fixed branch points lies in a different component of the target. Then by the degeneration formula, this class will be obtained as a sum (over all choices of splittings of the data) of classes glued together from virtual fundamental classes of various spaces of relative stable maps to each component. But any such relative stable map to one component, with points 0 and ∞ connecting it to adjacent elements of the chain (with the obvious variation if it is the end of the chain) has at most $j + 1 < 2g$ branch points away from 0 and ∞ . Hence by Remark 3.4, there is no curve of geometric genus g mapping to this component. Thus, after degeneration, the source curve can have no irreducible component of genus g , and hence the class is 0 in $\mathcal{M}_{g,n}^{rt}$. ■

Theorem 3.7. [Degeneration Theorem — Join-cut recursion] *If $r_{g,\alpha}^{\text{Fab}} > 0$, then*

$$\mathbb{F}^{g,\alpha} = \sum_{i+j=\alpha_k} ij H_{\alpha'}^0 \mathbb{F}^{g,\alpha'} \binom{r_{g,\alpha}^{\text{Fab}} - 1}{r_{g,\alpha'}^{\text{Fab}}} + \sum_{\alpha_i, \alpha_j} (\alpha_i + \alpha_j) \mathbb{F}^{g,\alpha'} + \sum_{i=1}^{l(\alpha)} \alpha_i^{2g+1} H_{\alpha}^0 \mathbb{G}_{g,1}.$$

where α' , α'' , etc., are as defined in the proof below.

This recursion inductively determines $\mathbb{F}^{g,\alpha}$ in terms of $\mathbb{G}_{g,1}$ (Theorem 3.8). There is no “base case” necessary. This result is best stated in terms of generating series (see Corollary 3.9).

Proof. We obtain a recursion for Faber-Hurwitz classes by a similar argument to that used in Lemma 3.6 above. We degenerate the target into two pieces $\mathbb{P}_L^1 \cup \mathbb{P}_R^1$, where ∞ and one of the $r_{g,\alpha}^{\text{Fab}} = r_\alpha^g - (2g - 1) = d + l(\alpha) - 1$ fixed branch points are on the right component \mathbb{P}_R^1 , and the remaining $r_{g,\alpha}^{\text{Fab}} - 1$ fixed branch points are on the left component \mathbb{P}_L^1 . By the degeneration formula (3.7), this class will be obtained as a sum (over all choices of splittings of the data) of classes glued together from virtual fundamental classes of various spaces of relative stable maps to each component. We discard every term that does not have a smooth component of genus g .

There are two cases: The genus g curve maps to the left component \mathbb{P}_L^1 or the right component \mathbb{P}_R^1 .

Case 1. If the genus g curve maps to the right component \mathbb{P}_R^1 , then by Remark 3.4, all the moving branch points must also map to \mathbb{P}_R^1 (we need $2g$ branching total away from $\mathbb{P}_L^1 \cap \mathbb{P}_R^1$ and ∞ in order to have a genus g curve in the cover). Also by Remark 3.4(a), the cover of \mathbb{P}_R^1 must be a union of trivial covers, together with one cover by a connected arithmetic genus g curve that is completely branched over ∞ and $\mathbb{P}_L^1 \cap \mathbb{P}_R^1$. In particular, the partition over the node $\mathbb{P}_L^1 \cap \mathbb{P}_R^1$ must be the same as the partition α over ∞ , and the genus g component can be on the component of the source corresponding to any α_j . Also, the cover of \mathbb{P}_L^1 must be genus 0 and connected, and the number of such covers is the genus 0 Hurwitz number

$$H_\alpha^0 = (d - 2 + l(\alpha))! d^{l(\alpha)-3} \prod_{i=1}^{l(\alpha)} \frac{\alpha_i \alpha_i}{\alpha_i!}. \tag{3.8}$$

(This celebrated formula was first given by Hurwitz, and has now been proved in many ways, see for example, [18].) This case is depicted pictorially in Figure 4; here the genus g curve corresponds to α_1 . We thus get a contribution to $\mathbb{F}^{g,d}$ of

$$\sum_{i=1}^{l(\alpha)} \alpha_i H_\alpha^0 \mathbb{G}_{g,\alpha_i} = \sum_{i=1}^{l(\alpha)} \alpha_i^{2g+1} H_\alpha^0 \mathbb{G}_{g,1}. \tag{3.9}$$

The equality in (3.9) is through Theorem 3.5. The contributions to the left side of (3.9) are as follows: We have a contribution of \mathbb{G}_{g,α_i} from the map to \mathbb{P}_L^1 . We have a contribution of $\mathbb{G}_{g,\alpha_i} \prod_{j \neq i} (1/\alpha_j)$ from the map to \mathbb{P}_R^1 (since the trivial cover contributes the order α_k of the automorphism group and we are therefore counting objects weighted by $1/\alpha_k$). We have a multiplicity of $\prod \alpha_i$ from the kissing condition.

Case 2a. If otherwise the genus g curve maps to the left component \mathbb{P}_L^1 , then by Lemma 3.6, all the non-fixed branch points must also map to \mathbb{P}_L^1 (as we need at least $2g - 1$ moving branch points in order to get a non-zero contribution), so the only branching over \mathbb{P}_R^1 away from its 0 and ∞ is simple branching over one (fixed) point. By Remark 3.3, the cover of \mathbb{P}_R^1 is a union of trivial covers, with one almost-trivial cover. If the almost-trivial cover of \mathbb{P}_R^1 is completely branched over $\mathbb{P}_L^1 \cap \mathbb{P}_R^1$, and has two preimages over ∞ (e.g., as shown in Figure 3), then it can connect any two of the points over ∞ (corresponding to two parts of the partition α , say, α_i and α_j). The virtual fundamental class of this space of relative stable maps to \mathbb{P}_R^1 is then $\alpha_i \alpha_j / \prod \alpha_i$: Any trivial cover corresponding to α_k ($k \neq i, j$) is weighted by $1/\alpha_k$. The space of




Figure 4: First case in degeneration argument in the proof of Theorem 3.7.

relative maps to \mathbb{P}_L^1 corresponds to maps from a connected curve of genus g , with branching over $\mathbb{P}_L^1 \cap \mathbb{P}_R^1$ corresponding to the partition α' obtained by removing α_i and α_j from α , and adding $\alpha_i + \alpha_j$, with all but $2g - 1$ branch points fixed. In other words, the contribution is precisely $\mathbb{F}^{g, \alpha'}$. Finally, the multiplicity in the degeneration formula is the product of the multiplicities of the kissing over the node $\mathbb{P}_L^1 \cap \mathbb{P}_R^1$, i.e., $\prod \alpha'_i = (\prod \alpha_i) (\alpha_i + \alpha_j) / (\alpha_i \alpha_j)$. Thus we obtain a contribution to $\mathbb{F}^{g, \alpha}$ of

$$\sum_{\alpha_i, \alpha_j} (\alpha_i + \alpha_j) \mathbb{F}^{g, \alpha'}$$

Case 2b. Finally, if the almost-trivial cover over \mathbb{P}_R^1 has one preimage over ∞ (corresponding to α_k , say), and two preimages over $\mathbb{P}_L^1 \cap \mathbb{P}_R^1$ (q_1 and q_2 , say, branching with multiplicity i and j , respectively, where $i + j = \alpha_k$), then we have contributions of $\alpha_k / \prod \alpha_i$ from the cover of \mathbb{P}_R^1 , and $\prod \alpha_i \times ij / \alpha_k$ from the kissing multiplicities. We next examine the contribution from the stable relative maps to \mathbb{P}_L^1 . The cover must have two connected components, one containing the point q_1 and one containing the point q_2 . As the map has a component of geometric genus g , one of these two components must be arithmetic genus g , and the other must be arithmetic genus 0. Suppose the genus 0 curve contains q_1 , and the genus g curve contains q_2 . Suppose the partition corresponding to the genus 0 curve is α' (i.e., its ramification above the node $\mathbb{P}_L^1 \cap \mathbb{P}_R^1$), and the partition corresponding to the genus g curve is α'' , so $\alpha' + \alpha'' = \alpha - \alpha_k + \{i, j\}$, and $i \in \alpha', j \in \alpha''$. In order for the contribution to be non-zero, by Lemma 3.6, all of the $(2g - 1)$ “moving” branching must belong to the genus g component, so all of the genus 0 curve’s branch points must be fixed. There are $\binom{r_{g, \alpha}^{\text{Fab}} - 1}{r_{g, \alpha'}^{\text{Fab}}}$ ways of choosing which fixed branch point on \mathbb{P}_L^1 belongs to the genus 0 curve, and which belongs to the genus g curve. Hence the contribution of the cover of \mathbb{P}_L^1 is $H_{\alpha'}^0 \mathbb{F}^{g, \alpha''} \binom{r_{g, \alpha}^{\text{Fab}} - 1}{r_{g, \alpha'}^{\text{Fab}}}$, so the combined contribution to $\mathbb{F}^{g, \alpha}$ from this case is

$$\sum_{i+j=\alpha_k} ij H_{\alpha'}^0 \mathbb{F}^{g, \alpha''} \binom{r_{g, \alpha}^{\text{Fab}} - 1}{r_{g, \alpha'}^{\text{Fab}}}$$

Summing the three types of contribution, we obtain the result. ■

Theorem 3.8. *For each fixed g , $\mathbb{F}^{g,\alpha}$ is a rational multiple of $\mathbb{G}_{g,1}$, as determined by the recursion of Theorem 3.7.*

Proof. By Remark 3.4(a), if $r_{g,\alpha}^{Fab} = 0$, then $\mathbb{F}^{g,\alpha} = 0$; this is the trivial base case of the recursion in Theorem 3.7. The result follows by induction. ■

For convenience, we define the *Faber-Hurwitz number* $F_\alpha^g \in \mathbb{Q}$ to be this multiple of $\mathbb{G}_{g,1}$, to remind us that these classes are all commensurate:

$$F_\alpha^g \mathbb{G}_{g,1} := \mathbb{F}^{g,\alpha}. \tag{3.10}$$

(Corollary 6.6 is a sign that F_α^g is a well-behaved quantity to consider.)

Theorem 3.7 is best formulated in terms of generating series. A natural generating series for genus 0 Hurwitz numbers is

$$\widehat{H}^0(z; \mathbf{p}) := \sum_{\alpha \in \mathcal{P}} z^{|\alpha|} \frac{p_\alpha}{|\text{Aut}\alpha|} \frac{H_\alpha^0}{r_\alpha^0!}, \tag{3.11}$$

where $\mathbf{p} = (p_1, p_2, \dots)$, and a natural generating series for Faber-Hurwitz numbers, which we shall call the *Faber-Hurwitz series*, is

$$F^g(z; \mathbf{p}) := \sum_{\alpha \in \mathcal{P}} z^{|\alpha|} \frac{p_\alpha}{|\text{Aut}\alpha|} \frac{F_\alpha^g}{r_{g,\alpha}^{Fab!}}. \tag{3.12}$$

Then the recursion of Theorem 3.7, for *classes*, becomes the following (linear) partial differential equation for F^g .

Corollary 3.9. [Degeneration Theorem — Join-Cut Equation for Faber-Hurwitz Series] *For $g \geq 1$, $F^g(z; \mathbf{p})$ is the unique formal power series solution to*

$$\begin{aligned} \left(z \frac{\partial}{\partial z} - 1 + \sum_{i \geq 1} p_i \frac{\partial}{\partial p_i} \right) F^g &= \sum_{i,j \geq 1} p_i p_j \left(i \frac{\partial}{\partial p_i} \widehat{H}^0 \right) \left(j \frac{\partial}{\partial p_j} F^g \right) \\ &+ \frac{1}{2} \sum_{i,j \geq 1} p_i p_j (i+j) \frac{\partial}{\partial p_{i+j}} F^g + \sum_{i \geq 1} i^{2g+1} p_i \frac{\partial}{\partial p_i} \widehat{H}^0. \end{aligned}$$

In Corollary 3.9, we have changed from classes to numbers by erasing $\mathbb{G}_{g,1}$ from Theorem 3.7 via (3.10). To obtain a generating series for Faber-Hurwitz *classes*, we would simply multiply this series by $\mathbb{G}_{g,1}$.

3.4. Localization of Faber-Hurwitz Classes

This section contains the Localization Tree Theorem for Faber-Hurwitz classes. It is to be thought of in conjunction with the Degeneration Theorem. The theorem gives the fundamental relationship between Faber-Hurwitz classes on the one hand, and intersection numbers and (genus 0) single and double Hurwitz numbers on the other hand. Implicitly, it allows us to determine the Faber-Hurwitz numbers. It involves a sum over a set $\mathcal{T}_{g,m}$, which is one of three sets of trees defined as follows.

Definition 3.10. (Localization Trees) Consider rooted trees with the following properties. For each tree \mathfrak{t} there are three classes of vertices: $\mathcal{V}_0(\mathfrak{t})$, the 0-vertices; $\mathcal{V}_\infty(\mathfrak{t})$, the ∞ -vertices; $\mathcal{V}_t(\mathfrak{t})$, the t -vertices. All t -vertices are monovalent (here t stands for “tail”). The number of non-root 0-vertices in \mathfrak{t} is denoted by $\eta_0(\mathfrak{t})$. The non-root 0-vertices are labelled (each receives one of $\eta_0(\mathfrak{t})$ labels in all possible ways). The ∞ -vertices are not labelled. There are two classes of edges: $\mathcal{E}_{0\infty}(\mathfrak{t})$, the 0∞ -edges, each of which joins a 0-vertex to an ∞ -vertex; $\mathcal{E}_\infty(\mathfrak{t})$, the ∞t -edges, each of which joins a non-root ∞ -vertex to a t -vertex.

There is at least one edge. Each edge is assigned a positive integer weight, and the integer weight on an 0∞ -edge e is denoted by $\varepsilon(e)$. The t -vertices incident with an edge of weight k are labelled among themselves, for each $k \geq 1$. The list of the weights on all 0∞ -edges incident with a 0-vertex v is specified by the partition $\delta^v(\mathfrak{t})$, the list of the weights on all 0∞ -edges incident with an ∞ -vertex v is specified by the partition $\beta^v(\mathfrak{t})$, and the list of the weights on all ∞t -edges incident with a non-root ∞ -vertex v is specified by the partition $\gamma^v(\mathfrak{t})$. For each non-root ∞ -vertex v , we impose the condition that

$$|\beta^v(\mathfrak{t})| = |\gamma^v(\mathfrak{t})|. \quad (3.13)$$

The root-vertex may be either a 0-vertex or an ∞ -vertex, and is denoted by \bullet . For $m \geq 1$, we define $\mathcal{T}_{g,m}$ to be the set of rooted trees above in which the root-vertex is a 0-vertex of degree m . For $j \geq 1$, we define $\mathcal{T}_{0,j}$ to be the subset of $\mathcal{T}_{g,1}$ in which the edge incident with the root-vertex has weight j . For $j \geq 1$, we define $\mathcal{T}_{\infty,j}$ to be the set of rooted trees above in which the root-vertex is a monovalent ∞ -vertex, and the edge incident with the root-vertex has weight j .

We refer to the trees in any of the sets $\mathcal{T}_{g,m}$, $\mathcal{T}_{0,j}$, and $\mathcal{T}_{\infty,j}$ as “localization trees”.

Hereinafter for any localization tree \mathfrak{t} we shall subsume the dependence on \mathfrak{t} of the above sets of vertices and edges, and the partitions, by suppressing the occurrence of \mathfrak{t} as an argument. We also define some more notation. Let

$$r_\infty := \sum_{v \in \mathcal{V}_\infty} r_{\gamma^v, \beta^v}^0, \quad (3.14)$$

and let

$$\alpha := \prod_{v \in \mathcal{V}_\infty} \gamma^v \quad (3.15)$$

be the partition formed by all the parts of each of the γ^v 's for non-root ∞ -vertices v , and let $d = |\alpha|$. Examples of such trees *without a specified root* are given in Figure 5 (these examples correspond to the geometric picture of Figure 6).

We now define some terminology that will allow us to write the relative virtual localization calculation cleanly. Let $\mathbb{P}_m^g(\alpha_1, \dots, \alpha_m)$ be the dimension $2g - 1$ portion of

$$\frac{1 - \lambda_1 + \dots + (-1)^g \lambda_g}{(1 - \alpha_1 \Psi_1) \cdots (1 - \alpha_m \Psi_m)}$$

on $\mathcal{M}_{g,m}^r$, viz.,

$$\mathbb{P}_m^g(\alpha_1, \dots, \alpha_m) := \sum_{\substack{a_1, \dots, a_m, k \geq 0, \\ a_1 + \dots + a_m + k = g - 2 + m}} (-1)^k \langle \tau_{a_1} \cdots \tau_{a_m} \lambda_k \rangle_g^{\mathbb{F}^{\text{ab}}} \alpha_1^{a_1} \cdots \alpha_m^{a_m}. \quad (3.16)$$




Figure 5: Sample decorated trees (Definition 3.10) corresponding roughly to the examples of Figure 6. Omitted are further combinatorial labelling of vertices.

Note that \mathbb{P}_m^g is a (Chow-valued) polynomial in the numbers $\alpha_1, \dots, \alpha_m$, symmetric of degree between $m - 2$ and $g - 2 + m$, and its leading coefficients (the portion of homogeneous degree $g - 2 + m$) are precisely the subject of Faber’s Intersection Number Conjecture. (It is easy to show that the homogeneous degree $m - 2$ and $m - 1$ portions of this polynomial vanish, but we shall not need this fact.) We shall refer to \mathbb{P}_m^g as the *Faber polynomial*.

For any localization tree \mathfrak{t} , let

$$B(\mathfrak{t}) := \prod_{e \in \mathcal{E}_{\infty}} \epsilon(e), \quad C(\mathfrak{t}) := \prod_{v \in \mathcal{V}_0^1} \frac{H_{\delta^v}^0}{r_0^v \delta^v!}, \quad D(\mathfrak{t}) := \prod_{v \in \mathcal{V}_\infty^1} \frac{H_{\gamma^v, \beta^v}^0}{r_\infty^v \gamma^v! \beta^v!} \quad (3.17)$$

and, for $\mathfrak{t} \in \mathcal{T}_{g, m}$, let

$$A(\mathfrak{t}) := \mathbb{P}_m^g(\delta^\bullet) \prod \frac{\epsilon(e)^{\epsilon(e)}}{\epsilon(e)!}, \quad (3.18)$$

where the product is over all edges e incident with the root-vertex \bullet of \mathfrak{t} . Let “ \ddagger ” as a superscript on a product denote the removal of the contribution of the root-vertex from that product.

Theorem 3.11. (Localization Tree Theorem — Tree Summation) *For $g \geq 1$ and $\alpha = (\alpha_1, \dots, \alpha_m) \vdash d$,*

$$\mathbb{F}^{g, \alpha} = \sum_{m \geq 1} \sum_{\mathfrak{t} \in \mathcal{T}_{g, m}} (-1)^{r_\infty} r_{g, \alpha}^{\text{Fab}1} \binom{r_\alpha^g - r_\infty}{r_{g, \alpha}^{\text{Fab}}} \frac{1}{\eta_0(\mathfrak{t})!} A(\mathfrak{t}) B(\mathfrak{t}) C^\ddagger(\mathfrak{t}) D(\mathfrak{t}), \quad (3.19)$$

where the sum is subject to (3.15).

Note that the sum in (3.19) is finite, and that the binomial coefficient in it is zero unless r_∞ is small (at most $r_\alpha^g - r_{g, \alpha}^{\text{Fab}} = 2g - 1$).

Before proving Theorem 3.11, we digress to observe that just the “shape” of the formula (3.19) quickly yields the result **(I)** promised in Section 1.

Theorem 3.12. (Socle Statement for $\mathcal{M}_{g,n}^{rt}$) $R_{2g-1}(\mathcal{M}_{g,n}^{rt}) \cong \mathbb{Q}$.

Proof. We shall show that any monomial in the ψ -classes (pushed forward to $\mathcal{M}_{g,1}$) is a multiple of $\mathbb{G}_{g,1}$. This (with Faber’s Non-vanishing Theorem 2.2 and Remark 2.3 (iii)) implies that $R_{2g-1}(\mathcal{M}_{g,n}^{rt})$ is generated by a single element. *In particular, $\mathbb{G}_{g,1} \neq 0$ and*

$$\mathbb{Q} \xrightarrow{\times \mathbb{G}_{g,1}} R_{2g-1}(\mathcal{M}_{g,1}) \tag{3.20}$$

is an isomorphism.

We show first that $\mathbb{P}_n^g(\alpha)$ is a multiple of $\mathbb{G}_{g,1}$ for each partition α , by induction on $|\alpha| = \sum_i \alpha_i$. Now $\mathbb{P}_n^{g,\alpha}$ is a multiple of $\mathbb{G}_{g,1}$ (Theorem 3.8), and the contribution of the unique graph with $r_\infty = 0$ (“the simplest graph in $\mathcal{T}_{g,m}$ ”) to Theorem 3.11 is a non-zero multiple of $\mathbb{P}_n^g(\alpha)$. The contribution of any other graph is a multiple of $\mathbb{P}_{n'}^g(\alpha')$ for some smaller $|\alpha'|$. Thus, by induction, $\mathbb{P}_n^g(\alpha)$ is a multiple of $\mathbb{G}_{g,1}$ as desired.

Next, we apply the “polynomiality trick” used in [24] and [26]. We fix g and n , and hence the polynomial $\mathbb{P}_n^g(\cdot)$. For arbitrary choices of $\alpha_1, \dots, \alpha_n$, $\mathbb{P}_n^g(\alpha)$ is a multiple of $\mathbb{G}_{g,1}$. But by knowing enough values of a polynomial of known degree, we can determine its coefficients as linear combinations of these values. Hence all coefficients of $\mathbb{P}_n^g(\alpha)$ are multiples of $\mathbb{G}_{g,1}$, and in particular, the monomials in ψ -classes (of degree $g - 2 + n$) are multiples of $\mathbb{G}_{g,1}$. ■

Proof of Theorem 3.11. We apply relative virtual localization as developed in [26] (based on the foundational [23]). As with many virtual localization calculations, Faber classes will be expressed as sums over certain graphs. We show that the set of trees $\mathcal{T}_{g,m}$ is precisely the set of graphs that is required for this purpose by supplying a geometric meaning to the vertices, edges, partitions, weights, and constants associated with $t \in \mathcal{T}_{g,m}$ through a geometry-combinatorics lexicon.

Classification of torus-fixed loci. We shall have a contribution from each torus-fixed locus of stable relative maps. Recall that Faber-Hurwitz classes are defined by considering the pullback of a linear space under the branch map, applied to the virtual fundamental class of the moduli space of stable relative maps (Equation (3.5)). As in the proof of the “tautological vanishing theorem” of [26], we choose a linearization on the branch-class that corresponds to requiring the $r_{g,\alpha}^{\text{Fab}} = r_\alpha^g - 2g + 1$ fixed branch points to go to 0. Hence in any contributing torus-fixed locus, the amount of branching over ∞ is at most $2g - 1$.

We now classify the fixed loci which can appear in the “rational tails” case, where we must have a smooth irreducible component of genus g , and then consider the evaluation of their contributions (**L1** to **L5** below), although some require further elaboration. Fixed loci correspond to maps of the following sort (see Figure 6), and in each case we shall see that the same statements hold for both the left hand side (the simple case) and the right hand side (the composite case) of Figure 6, although the arguments differ slightly. The components mapping surjectively onto \mathbb{P}^1 are trivial covers (see **L1**). Over 0, there can be smooth points (as in Figure 6(a)) (see **L3**), nodes (Figure 6(b)), or contracted components (Figure 6(c)) (see **L2**). Over ∞ , either there is no “sprouting”, and the preimage of ∞ consists of smooth points (the “simple” case, see

the left side of Figure 6), or the target “sprouts”, and we obtain a relative stable map to an unrigidified target (that we denote by T_R), with possibly-disconnected source (the “*composite*” case, see the right side of Figure 6) (see **L4**). In the composite case, let z be the node where the “sprouted” portion T_R of the target meets the “unsprouted” portion. If T_L denotes the “unsprouted” portion of T , then ∞_{T_L} and 0_{T_R} are identified. Note that in the composite case, the sprouted portion of T need not be a single \mathbb{P}^1 ; it could be a chain of \mathbb{P}^1 ’s.




Figure 6: Sketches of two examples of torus-fixed relative stable maps to $X = (\mathbb{P}^1, \infty)$. T is the “sprouted target”.

Over ∞ , we can have no genus g components by Remark 3.4 — the only possible branching is over the two endpoints (z and ∞_T), together with at most $2g - 1$ more. Thus the genus g component must lie over 0, and all irreducible curves over ∞ must be genus 0. (In particular, in the second figure in Figure 6, there cannot be any contracted components mapping to T_R , so the picture is misleading.)

Geometry-combinatorics lexicon. With the following combinatorial elements, we associate the following geometrical information.

Vertices and edges of $\mathfrak{t} \in \mathcal{T}_{g,m}$:

- 0-vertex: \leftrightarrow connected component of the preimage of 0 (contracted curve, node, or smooth point).
- ∞ -vertex: \leftrightarrow connected component of pre-images of ∞_X .
- t -vertex: \leftrightarrow preimage of ∞_T .
- 0∞ -edge: \leftrightarrow trivial cover of \mathbb{P}^1 ; edge joins vertices corresponding to the loci it meets.
- root \bullet : \leftrightarrow genus g (contracted) curve.

Edge weights:

- weight $\varepsilon(e)$ on the 0∞ -edge e : \leftrightarrow degree of corresponding trivial cover.
- weight on an ∞t -edge: \leftrightarrow contribution to ramification over ∞_X on component specified by the ∞ -vertex.

Partitions:

- δ^v for a 0-vertex v : *Comb.* formed by weights on 0∞ -edges incident with v . *Geom.* formed by the degrees of the trivial covers meeting that component.
- β^v for an ∞ -vertex v : *Comb.* formed by weights on 0∞ -edges incident with v . *Geom.* formed by the degrees of the trivial covers meeting that component.
- γ^v for an ∞ -vertex v : *Comb.* formed by weights on ∞t -edges incident with v . *Geom.* specifies ramification over ∞_X on that component.
- α : *Comb.* see Equation (3.15). *Geom.* ramification over ∞ .

Conditions:

- $|\beta^v| = |\gamma^v|$: *Comb.* see Equation (3.13). *Geom.* the degree of the map from this component to T_R is $|\beta^v|$ (by examining the preimage of z) and $|\gamma^v|$ (from the preimage of ∞_T).

Constants:

- $\eta_0(t)$: *Comb.* number of non-root 0-vertices. *Geom.* number of connected components of the preimage of 0 excluding the component containing the contracted genus g curve.
- n : *Comb.* $l(\alpha)$. *Geom.* number of pre-images of ∞_T .
- m : *Comb.* degree of the root-vertex \bullet (note $m \leq n$). *Geom.* number of trivial covers of \mathbb{P}^1 meeting the genus g contracted curve.
- r_∞ : *Comb.* see Equation (3.14). *Geom.* the total branching over ∞_X .
- $r_{\delta^v}^0$ for a 0-vertex v : *Comb.* see Equation (3.1). *Geom.* the total branching contributed by the component corresponding to v .
- r_{γ^v, β^v}^0 for an ∞ -vertex v : *Comb.* see Equation (3.3). *Geom.* the total branching contributed by the component corresponding to v .
- d : *Comb.* $d = |\alpha|$. *Geom.* degree of the relative stable map.

Relative virtual localization. Relative virtual localization ([26, Thm. 3.6], see [26, Sec. 3.7] for the special case of target \mathbb{P}^1 , and [23, Sec. 4] for the non-relative case) tells us that the contribution of a fixed locus can be deduced by looking at the various parts of Figure 6. In what follows, t is the generator of the equivariant cohomology (or Chow) ring of a point, $t \in A_{\mathbb{C}^*}^1(pt) = \mathbb{Q}[t]$, although we will quickly forget the t . The relative virtual localization formula (abbreviated to **L** below) gives the following contributions associated with the salient “parts” of the graph. Each of the factors in the summand of (3.19) will be readily derivable, with the exception of the sign and $D(t)$ which will require more attention.

Items **L1** to **L4** below come from the description of the fixed loci, and **L5** arises from the cohomology class corresponding to fixing some branch points. Moreover, **L1** to **L3** hold for both the simple and the composite case.

The results for the simple case and the composite case are the same, but the arguments are slightly different.

L1. For each trivial cover of the target \mathbb{P}^1 of degree a , we have contribution $a^a / (a!t^a)$. Hence we obtain the product $\prod \frac{a^a}{q!}$ in (3.19) by collecting those contributions associated with the (genus g) root 0-vertex.

L2. For each contracted curve above 0 of genus h (Figure 6(c)), meeting trivial covers (components mapping surjectively onto \mathbb{P}^1) of degree $\alpha_1, \dots, \alpha_m$, respectively, we have a contribution

$$t^{-1} \left(t^h - \lambda_1 t^{h-1} + \dots + (-1)^h \lambda_h \right) \prod_{i=1}^m \frac{t}{t/\alpha_i - \psi_i}. \tag{3.21}$$

This contribution is on the factor $\overline{\mathcal{M}}_{h,m}$ corresponding to the contracted curve. We will have $h = 0$ or $h = g$, as all components have one of these two genera.

L3. For each node above 0 (Figure 6(b)) joining trivial covers of degrees α_1 and α_2 , we get a contribution of $t^{-1} \alpha_1 \alpha_2 / (\alpha_1 + \alpha_2)$. For each smooth point above 0 (Figure 6(a)), on a trivial cover of degree α_1 , we get t^{-1} / α_1 . Using the contribution $\prod (\alpha_i^{\alpha_i} / \alpha_i!)$ from **L1**, we obtain the factor $\boxed{H_{\delta_j^0 / r_{\delta_j^0}^0}!}$ in (3.19) for each (non-root) 0-vertex of degree 1 or 2, using the formula (3.8) for genus 0 single Hurwitz numbers. We also obtain the product of the $\boxed{\varepsilon(e)}$ in (3.19) corresponding to 0_∞ -edges e meeting a degree 1 or 2 (genus 0, non-root) 0-vertex.

L4. In the composite case (Figure 6(d)), we have a contribution of $1 / (-t - \psi_z)$, where ψ_z is the first Chern class of the line bundle corresponding to the cotangent space of T_R at z .

L5. From the pullback of the linear space by the branch morphism (informally, requiring $r_{g,\alpha}^{\text{Fab}}$ branch points to map to 0), we have a contribution of $((2g - r_\infty)t) ((2g + 1 - r_\infty)t) \dots ((r_\alpha^g - r_\infty)t)$ (there are $r_{g,\alpha}^{\text{Fab}}$ factors), from which we obtain $\boxed{r_{g,\alpha}^{\text{Fab}}! \binom{r_\alpha^g - r_\infty}{r_{g,\alpha}^{\text{Fab}}}}$ in (3.19).

So **L1–L4** arise from the fixed loci, and **L5** arises from the cohomology class corresponding to fixing some branch points. We take the product of these contributions, and read off the constant (t^0) term to obtain the contribution of this fixed locus.

The result is a $(2g - 1)$ -dimensional class on $\mathcal{M}_{g,n}^{rt}$. One of the ingredients (from **L3**) is a (tautological) class on $\overline{\mathcal{M}}_{g,m}$ corresponding to the contracted genus g curve mapping to 0. Now all tautological classes of dimension *less* than $2g - 1$ vanish on $\overline{\mathcal{M}}_{g,m}^{rt}$ by Remark 2.3(i). Thus a non-zero contribution is possible only by taking the contribution of a class of dimension *precisely* $2g - 1$ on $\mathcal{M}_{g,m}^{rt}$, and thus the contributions from every other ingredient must have dimension 0. In light of this observation, we list the contributions from each of the parts of Figure 6, ignoring the equivariant parameter t . Also, from **L3**, the contribution by the contracted genus g curve is $\boxed{\mathbb{P}_m^g(\delta^0)}$ so, with the contribution from **L1**, we obtain the term $\boxed{A(t)}$, a term in (3.19). In addition, we obtain the product of the $\boxed{\varepsilon(e)}$ over those edges e meeting the (genus g) root 0-vertex.

From (3.21) (using $h = 0$), any contracted genus 0 component over $0 \in \mathbb{P}^1$ meeting trivial covers of degrees $\delta_1^j, \dots, \delta_p^j$ ($p \geq 3$ by the stability condition) gives the dimension 0 contribution

$$\left(\prod_{i=1}^p \delta_i^j \right) \int_{\overline{\mathcal{M}}_{0,p}} \frac{1}{(1 - \delta_1^j \psi_1) \dots (1 - \delta_p^j \psi_p)}.$$

Combining this with $\prod_{i=1}^p \frac{\delta_i^{\delta_i^j}}{\delta_i!}$ from **L1**, using the ELSV formula [5, 6, 25] we obtain $\left(\prod \delta_i^j\right) H_{\delta_j}^0 / r_{\delta_j}^0!$. (We could have bypassed the ELSV formula, using instead the formula (3.8) for genus 0 Hurwitz numbers and the string equation, given in Proposition 2.6.) Thus we obtain the factor $\boxed{H_{\delta_j}^0 / r_{\delta_j}^0!}$ for each non-root 0-vertex of degree at least three. We also obtain the product of the $\boxed{\varepsilon(e)}$ corresponding to 0_∞ -edges e meeting all (genus 0, non-root) 0-vertices of degree at least 3. **L3** gives the same values for degree 1 and 2 so, combining the three sources for the ε 's, we now obtain the entire product $\boxed{B(t)}$, and combining the two sources for the $H_{\delta_j}^0 / r_{\delta_j}^0!$, we obtain the entire product $\boxed{C(t)}$.

The contributions from **L1–L3**, and **L5** are now exhausted.

It remains to obtain the sign $(-1)^{r_\infty}$, the term $D(t)$, and the division by $\eta_0(t)!$. To do so, we now appeal to **L4**. In the composite case (if the target ‘‘sprouts’’, Figure 6(d)), suppose $\overline{\mathcal{M}} := \overline{\mathcal{M}}_{\beta, \alpha}$ is the moduli space of relative stable maps to the unrigidified T_R , where β is the kissing partition above z . It is the moduli space of relative stable maps, where the source is a disjoint union of genus 0 curves. As always for maps to unrigidified targets, the virtual dimension of this space is one less than the number of ‘‘moving branch points’’ $r_{\overline{\mathcal{M}}}$. Then the contribution is the dimension 0 portion of $\frac{1}{-1-\psi_z} [\overline{\mathcal{M}}]^{\text{vir}}$ which is

$$(-1)^{r_{\overline{\mathcal{M}}}} \psi_z^{r_{\overline{\mathcal{M}}}-1} [\overline{\mathcal{M}}]^{\text{vir}}. \tag{3.22}$$

The sign gives us the factor $\boxed{\prod (-1)^{r^i, \beta^i}}$ which, from (3.14), is equal to $(-1)^{r_\infty}$, the sign in (3.19).

The proof of [26, Lem. 4.8] (see also [26, Fig. 2]) shows that ψ^a applied to $\overline{\mathcal{M}}$ can be interpreted as requiring that the target break into $a + 1$ components. More precisely, ψ^a applied to $[\overline{\mathcal{M}}]^{\text{vir}}$ is the same as gluing virtual fundamental classes of relative stable maps to the $a + 1$ components of T_R (in the same sense as the degeneration formula, with kissing multiplicities arising for each node of the target T_R), divided by $(a + 1)!$. This latter term gives a factor of $\boxed{1/r_\infty!}$.

In particular, $\psi^{r_{\overline{\mathcal{M}}}-1}$ corresponds to the target breaking into $r_{\overline{\mathcal{M}}}$ components. Because the resulting map must be stable, there must be some branching on each of these components (away from the nodes of T_R , and z and ∞_T). Thus as the total amount of branching is $r_{\overline{\mathcal{M}}}$ away from the nodes of T_R , and there is precisely this number of components, we must have branching number 1 on each irreducible component of T_R . By Remark 3.3(b), above each component of the T_R , we must have precisely one almost-trivial cover, along with some trivial covers.

Thus the contribution of (3.22) is the size of a discrete set (counted modulo automorphisms). This set counts the number of branched covers of T_R (a chain of $r_{\overline{\mathcal{M}}}$ \mathbb{P}^1 's), with one simple branching on each component, and given branching ε over z (a point at the end of the chain) and α over ∞_T (a point at the other end of the chain), satisfying the kissing condition over each node of T_R . By the gluing formula (or in-

deed, the much older technique of just studying the degeneration), this is the number of branched covers of \mathbb{P}^1 by a union of genus 0 curves with branching given by ε and α over two points z and ∞_T , and simple branching over $r_{\overline{\mathcal{M}}}$ other given fixed points.

We shall now see that this is (up to a combinatorial factor) a product of genus 0 double Hurwitz numbers. Recall that a genus 0 double Hurwitz number

$$H_{\alpha, \varepsilon}^0 \tag{3.23}$$

(where α and ε are partitions of some number e) counts the number of degree e covers of \mathbb{P}^1 by \mathbb{P}^1 , with branching α at one fixed point, ε at another, and simple branching at $r_{\alpha, \varepsilon}^0$ other fixed points. Suppose we are considering covers by N \mathbb{P}^1 ’s, where component i corresponds to the subpartition β^i of ε (over z) and the subpartition γ^i of α (over ∞_T). (Thus $|\beta^i| = |\gamma^i|$ is the degree of that subcover, $\varepsilon = \coprod_i \beta^i$, and $\alpha = \coprod_i \gamma^i$.) Then component i has simple branching over $r_{\infty}^i := r_{\beta^i, \gamma^i}^0$ of the fixed simple branch points. There are $\binom{\sum_i r_{\infty}^i}{r_{\infty}^1, \dots, r_{\infty}^N}$ ways of partitioning the branch points into these N sets. Once this partition is chosen, there are $\prod_{i=1}^N H_{\beta^i, \gamma^i}^0$ such branched covers. (One caution: We have cavalierly described H_{β^i, γ^i}^0 as enumerating a set. In reality, each cover is counted with multiplicity equal to the inverse of the size of its automorphism group, so H_{β^i, γ^i}^0 need not be integral, and in fact is not precisely for $\beta^i = \gamma^i = (d_i)$; trivial covers of degree d_i “count for” $1/d_i$.) We have obtained the factors $\boxed{H_{\beta^k, \gamma^k}^0}$ and $\boxed{\binom{r_{\infty}}{r_{\infty}^1, \dots, r_{\infty}^N}}$ in (3.19). (The numerator $r_{\infty}!$ in the multinomial coefficient cancels the $1/r_{\infty}!$ from earlier.) This is the term $D(t)$.

Finally, the $\boxed{1/\eta_0(t)!}$ in (3.19) is present because the trees have labelled non-root 0-vertices (Definition 3.10). ■

Part III. Algebraic Combinatorics

At this point, we have defined the Faber-Hurwitz classes $\mathbb{F}^{g, \alpha}$, which “virtually” correspond to “rational tail” curves admitting a branched cover of \mathbb{P}^1 with branching at ∞ corresponding to α , and “all but $2g - 1$ branching fixed”. Such classes are a multiple of a basic class $\mathbb{G}_{g, 1}$; this multiple is the Faber-Hurwitz number $F^{g, \alpha}$. By degeneration, we have obtained Corollary 3.9, the Degeneration Theorem for the generating series F^g for these numbers, involving the genus 0 Hurwitz series \widehat{H}^0 . By localization, we have also obtained Theorem 3.11, the Localization Tree Theorem, which describes these classes (or numbers) as a sum over certain rooted, labelled trees, involving genus 0 Hurwitz numbers, double Hurwitz numbers and the desired intersection numbers (of ψ -classes). Theorem 3.12 shows us that we can “invert” this expression, to determine intersection numbers in terms of Hurwitz numbers and double Hurwitz numbers. Our goal is to formalize this. The strategy is to show that the Localization Tree Theorem and the Degeneration Theorem, taken together, give a non-singular system of linear equations for the top Faber intersection numbers, so it has a unique solution, and that the conjectural values satisfy it.

We accomplish this by a sequence of transformations, which yield a number of refined versions of the Localization Tree Theorem and the Degeneration Theorem.

These versions of the Localization Tree Theorem (we say that these are results for the “localization side”) are given by the sequence

$$\boxed{\text{Theorem 3.11} \rightsquigarrow \text{Theorem 4.4} \rightsquigarrow \text{Corollary 5.1} \rightsquigarrow \text{Corollary 7.1}} .$$

These versions of the Degeneration Theorem (we say that these are results for the “degeneration side”) are given by the sequence

$$\boxed{\text{Theorem 3.7} \rightsquigarrow \text{Corollary 3.9} \rightsquigarrow \text{Corollary 5.2} \rightsquigarrow \text{Lemma 7.5}} .$$

4. Exponential Generating Series for Localization Trees

The purpose of this section is to “evaluate” the sum over localization trees that arises from the localization arguments in Theorem 3.11. Localization trees are a class of rooted, labelled trees, and we use the standard multivariate exponential generating series for combinatorial structures with many sets of labels, as well as variants of the standard branch decomposition for rooted trees.

4.1. Exponential Generating Series and the \star -Product

For a localization tree t , let $\eta_k(t)$ denote the number of t -vertices in t that are incident with an edge of weight k , $k \geq 1$.

Definition 4.1. *Let \mathcal{A} be a set of localization trees, with weight function wt . Then the exponential generating series for \mathcal{A} with respect to wt is*

$$[\mathcal{A}, \text{wt}]_\eta := \sum_{t \in \mathcal{A}} \frac{p_1^{\eta_1(t)} p_2^{\eta_2(t)} \dots}{\eta_0(t)! \eta_1(t)! \eta_2(t)! \dots} \text{wt}(t).$$

For now, we shall allow the range of the weight function to be any ring, or even a vector space, and in particular we allow geometric classes. Note that, as a formal power series in p_1, p_2, \dots , $[\mathcal{A}, \text{wt}]_\eta$ is always well-formed because of the balance condition (3.13), which ensures that there is only a finite number of localization trees t with $\eta_k(t) = i_k, k \geq 1$, for each i_1, i_2, \dots (so the coefficients are finite sums of weight function values).

Since localization trees are labelled objects, we consider a particular version of the standard \star -product for them, which is the Cartesian product together with a “label-distribution” operation. We define this \star -product as follows. Consider two localization trees t_1 and t_2 . Suppose that $\eta_k(t_1) = i_k$, and $\eta_k(t_2) = j_k$, for $k \geq 0$. Now choose subsets $\alpha_k \subseteq [i_k + j_k]$, with $|\alpha_k| = i_k$, for $k \geq 0$, and let $\beta_k = [i_k + j_k] \setminus \alpha_k$ (so $|\beta_k| = j_k$) (we use the notation $[n] = \{1, \dots, n\}$). Let $\alpha(t_1)$ be the tree obtained from t_1 by relabelling the labelled vertices as follows: Replace the label m on a non-root 0-vertex by the m th smallest element of α_0 , for $m = 1, \dots, i_0$; replace the label m on a t -vertex incident with an edge of weight k by the m th smallest element of α_k , for $m = 1, \dots, i_k, k \geq 1$. Let $\beta(t_2)$ be the tree obtained from t_2 by relabelling the labelled vertices with the elements of β_0, β_1, \dots , in the analogous manner. We call (α, β) a *compatible relabelling* (of (t_1, t_2)). Where convenient, we also refer to t_1 and t_2 as *canonically labelled*.

Definition 4.2. Let \mathcal{A}, \mathcal{B} be sets of localization trees. Then,

$$\mathcal{A} \star \mathcal{B} := \{(\alpha(t_1), \beta(t_2)) : (t_1, t_2) \in \mathcal{A} \times \mathcal{B}, (\alpha, \beta) \text{ a compatible relabelling of } (t_1, t_2)\}.$$

The reason for using exponential generating series for labelled combinatorial objects is the Product Lemma, given in the following result. For a proof, see, e.g., Goulden and Jackson [17, Lem. 3.2.11].

Lemma 4.3. [Product Lemma (for localization tree generating series)] Let $\mathcal{S}, \mathcal{A}, \mathcal{B}$ be sets of localization trees, with weight functions $\text{wt}, \text{wt}_1, \text{wt}_2$, respectively. Suppose there is a bijection

$$\mathcal{S} \xrightarrow{\sim} \mathcal{A} \star \mathcal{B} : t \mapsto (\alpha(t_1), \beta(t_2)),$$

subject to $\text{wt}(t) = \text{wt}_1(t_1)\text{wt}_2(t_2)$ and $\eta_k(t) = \eta_k(t_1) + \eta_k(t_2), k \geq 0$. Then

$$[\mathcal{S}, \text{wt}]_\eta = [\mathcal{A}, \text{wt}_1]_\eta [\mathcal{B}, \text{wt}_2]_\eta.$$

For any set \mathcal{S} of localization trees \mathcal{S} , let $\mathcal{S}^{\star k} := \mathcal{S} \star \dots \star \mathcal{S}$, where there are k \mathcal{S} ’s in this k -fold \star -product. We define $\mathcal{U}_k(\mathcal{S}) := \mathcal{S}^{\star k} / \mathfrak{S}_k$, under the natural action of \mathfrak{S}_k , for $k \geq 0$.

4.2. Branch Decompositions for Localization Trees

In order to decompose localization trees, we require three variants of the standard branch decomposition, described below. These are formalized as combinatorial mappings, called $\Omega^{\text{br}}, \Omega_0^{\text{br}}$, and $\Omega_\infty^{\text{br}}$.

First, we give Ω^{br} : If the root-vertex \bullet and incident edges are deleted from a localization tree t , then we obtain a list (t_1, \dots, t_m) of rooted trees on mutually distinct sets of vertices, each inheriting as root vertex its unique vertex that was adjacent to \bullet in t , where m is the degree of the root-vertex of t . Now let t'_i be obtained from t_i by joining the root vertex of t_i to a new copy of \bullet (which becomes the new root-vertex of t'_i), joined by an edge whose weight is equal to the weight of the edge joining \bullet to the root-vertex of t_i in t . Since \bullet is unlabelled in t , then (t'_1, \dots, t'_m) is equal to $(\alpha_1(t''_1), \dots, \alpha_m(t''_m))$, a compatible relabelling of canonical localization trees (t''_1, \dots, t''_m) . Then we define $\Omega^{\text{br}}(t) = (\alpha_1(t''_1), \dots, \alpha_m(t''_m))$. This corresponds to removing the root from the tree, and describing the remainder as a list of trees.

Second, we give Ω_0^{br} : Suppose that t is a localization tree whose root-vertex \bullet is a monovalent 0-vertex. Let the ∞ -vertex adjacent to \bullet be u , and let t' be the tree obtained by deleting \bullet and incident edge from t , and deleting all t -vertices adjacent to u , together with their incident edges, and rooting the resulting tree at u . Now form a new graph G , containing these deleted t -vertices (labelled as in t), joined to a new ∞ -vertex w by an edge of the same weight as the deleted incident edge in t . Then we define $\Omega_0^{\text{br}}(t) = (G, \Omega^{\text{br}}(t'))$. Now consider partition $\gamma = (1^{a_1} 2^{a_2} \dots)$, and let \mathfrak{w}_γ be the graph consisting of a single ∞ -vertex, joined by an edge of weight k to a_k canonically labelled monovalent t -vertices, $k \geq 1$. Then note that $\Omega_0^{\text{br}}(t) = (\alpha_0(\mathfrak{w}_\gamma), \alpha_1(t''_1), \dots, \alpha_m(t''_m))$, a compatible relabelling of the canonical $(\mathfrak{w}_\gamma, t''_1, \dots, t''_m)$, where $\gamma = \gamma^u(t), |\gamma| = |\beta^u(t)|$ (because of the balance condition (3.13) at u), and $\beta^u(t)$ has $m + 1$ parts, $m \geq 0$.

Third, we give $\Omega_{\infty}^{\text{br}}$: Suppose that \mathfrak{t} is a localization tree whose root-vertex \bullet is a monovalent ∞ -vertex. Let the 0-vertex adjacent to \bullet be v , with label i , then let \mathfrak{t}' be the tree obtained by removing \bullet from \mathfrak{t} , and removing the label from the vertex v , and rooting the resulting tree at v . Let v_i be the graph consisting of a single, 0-vertex, labelled i . Then we define $\Omega_{\infty}^{\text{br}}(\mathfrak{t}) = (v_i, \Omega_{\infty}^{\text{br}}(\mathfrak{t}'))$, and note that $\Omega_{\infty}^{\text{br}}(\mathfrak{t}) = (\alpha_0(v_1), \alpha_1(\mathfrak{t}'_1), \dots, \alpha_m(\mathfrak{t}'_m))$, a compatible relabelling of the canonical $(v_1, \mathfrak{t}'_1, \dots, \mathfrak{t}'_m)$, where $\delta^v(\mathfrak{t})$ has $m + 1$ parts, $m \geq 0$.

4.3. Generating Series Form for the Localization Tree Theorem

In the next result, Theorem 4.4, which is the second form of the Localization Tree Theorem, we introduce a generating series, \mathbb{T}^g , for the tree summation that arose in Theorem 3.11, which was the first form of the Localization Tree Theorem. We also introduce two ancillary classes of generating series $f_j, g_j, j \geq 1$. These are all exponential generating series with respect to particular weight functions, and the proofs of the equations that relate them are combinatorial, applying the variants of the branch decomposition, together with the \star -product and the Product Lemma. However, the statement of Theorem 4.4 is purely algebraic (without reference to trees), and the series \mathbb{T}^g, f_j, g_j are all uniquely defined by the equations in the statement. Note that the coefficients in \mathbb{T}^g are geometric *classes*, while the coefficients in f_j, g_j, F^g are *rationals*.

The statement of the theorem involves the generating series \widehat{H}^0 for genus 0 Hurwitz numbers, defined in (3.11), and the generating series for genus 0 double Hurwitz numbers, given by

$$H^0(z, u; \mathbf{p}; \mathbf{q}) := \sum_{\substack{\alpha, \beta \in \mathcal{P} \\ |\alpha| = |\beta|}} z^{|\beta|} p_{\alpha} q_{\beta} u^{l(\beta)} \frac{H_{\alpha, \beta}^0}{r_{\alpha, \beta}^0 |\text{Aut}\alpha| |\text{Aut}\beta|}, \tag{4.1}$$

where $\mathbf{p} = (p_1, p_2, \dots)$, $\mathbf{q} = (q_1, q_2, \dots)$, and $r_{\alpha, \beta}^0$ is defined in (3.3).

Theorem 4.4. (Localization Tree Theorem — Functional Equations) *For $g \geq 1$, we have*

$$(a) \quad F^g(z; \mathbf{p}) \mathbb{G}_{g,1} = [u^{2g-1}] \mathbb{T}^g \left(\frac{z}{1-u}, -u; \frac{-u}{1-u} \mathbf{p} \right),$$

where

$$(b) \quad \mathbb{T}^g(z, u; \mathbf{p}) = \sum_{n \geq 1} \frac{1}{n!} \sum_{\substack{a_1, \dots, a_n, k \geq 0, \\ a_1 + \dots + a_n + k = g - 2 + n}} (-1)^k \langle \tau_{a_1} \dots \tau_{a_n} \lambda_k \rangle_g^{\text{Fab}} \prod_{j=1}^n \xi^{(a_j)}(z, u; \mathbf{p}),$$

with

$$\xi^{(i)}(z, u; \mathbf{p}) := \sum_{j \geq 1} \frac{j^{j+i}}{j!} f_j(z, u; \mathbf{p}), \quad i \geq 0, \tag{4.2}$$

and for $j \geq 1$, (c) $f_j = u^{-2} \left(j \frac{\partial}{\partial q_j} H^0(z, u; \mathbf{p}; \mathbf{q}) \right) \Big|_{\substack{q_i = g_i \\ i \geq 1}}$, (d) $g_j = \left(j \frac{\partial}{\partial q_j} \widehat{H}^0(1; \mathbf{q}) \right) \Big|_{\substack{q_i = f_i \\ i \geq 1}}$,

with $f_j = f_j(z, u; \mathbf{p})$, $g_j = g_j(z, u; \mathbf{p})$.

Proof. As special cases of weight functions for localization trees \mathfrak{t} , consider $\text{wt}_g(\mathfrak{t}) := A(\mathfrak{t})\text{wt}_0(\mathfrak{t})$, $\text{wt}_0(\mathfrak{t}) := B(\mathfrak{t})C^\ddagger(\mathfrak{t})D(\mathfrak{t}) \prod_{v \in \mathcal{V}_\infty} u^{l(\beta^v)-2} z^{|\beta^v|}$, and $\text{wt}_\infty(\mathfrak{t}) := B(\mathfrak{t})C(\mathfrak{t})D^\ddagger(\mathfrak{t}) \prod_{v \in \mathcal{V}_\infty} u^{l(\beta^v)-2} z^{|\beta^v|}$, where A, B, C, D were defined in (3.18), (3.17). Now define generating series

$$\mathbb{T}^g(z, u; \mathbf{p}) := [\cup_{m \geq 1} \mathcal{T}_{g,m}, \text{wt}_g]_\eta, \quad g \geq 1, \quad f_j := [\mathcal{T}_{0,j}, \text{wt}_0]_\eta, \quad g_j := [\mathcal{T}_{\infty,j}, \text{wt}_\infty]_\eta, \quad j \geq 1.$$

For part (a), in \mathbb{T}^g we have $p_1^{\eta_1(\mathfrak{t})} p_2^{\eta_2(\mathfrak{t})} \dots \eta_1(\mathfrak{t})! \eta_2(\mathfrak{t})! \dots = p_\alpha / |\text{Aut}_\alpha|$, from (3.15). Thus, for $g \geq 1$ and $|\alpha| = d \geq 1$, we have

$$\mathbb{F}^{g,\alpha} = r_{g,\alpha}^{\text{Fab}}! |\text{Aut}_\alpha| \sum_{r_\infty} \binom{r_\alpha^g - r_\infty}{r_{g,\alpha}^{\text{Fab}}} (-1)^{r_\infty} \left[p_\alpha u^{r_\infty - l(\alpha)} z^d \right] \mathbb{T}^g(z, u; \mathbf{p}),$$

from Theorem 3.11. But $\binom{r_\alpha^g - r_\infty}{r_{g,\alpha}^{\text{Fab}}} = [u^{2g-1-r_\infty}] (1-u)^{-(d+l(\alpha))}$, so

$$\mathbb{F}^{g,\alpha} = r_{g,\alpha}^{\text{Fab}}! |\text{Aut}_\alpha| \left[p_\alpha u^{2g-1} z^d \right] \mathbb{T}^g \left(\frac{z}{1-u}, -u; \frac{-u}{1-u} \mathbf{p} \right).$$

Then part (a) follows from (3.12) and (3.10).

For part (b), define $\mathcal{T}_{g,\delta}$ to be the set of localization trees \mathfrak{t} in $\mathcal{T}_{g,m}$ with $\delta^\bullet(\mathfrak{t}) = \delta$, for any partition δ with m parts, $m \geq 1$. Then variant Ω^{br} of the branch decomposition gives a bijection

$$\Omega^{\text{br}} : \mathcal{T}_{g,\delta} \xrightarrow{\sim} \mathcal{U}_{a_1}(\mathcal{T}_{0,1}) \star \mathcal{U}_{a_2}(\mathcal{T}_{0,2}) \star \dots : \mathfrak{t} \mapsto (\alpha_1(t'_1), \dots, \alpha_m(t'_m)),$$

where $\delta = (1^{a_1} 2^{a_2} \dots)$ with $a_1 + a_2 + \dots = m$. It is straightforward to check in this bijection that $\eta_k(\mathfrak{t}) = \eta_k(t'_1) + \dots + \eta_k(t'_m)$, for $k \geq 0$. Moreover, $\text{wt}_g(\mathfrak{t}) = \mathbb{P}_m^g(1^{a_1} 2^{a_2} \dots) \left(\prod_{j \geq 1} (j^j / j!)^{a_j} \right) \text{wt}_0(t'_1) \dots \text{wt}_0(t'_m)$, so from the Product Lemma we obtain

$$[\mathcal{T}_{g,\delta}, \text{wt}_g]_\eta = \mathbb{P}_m^g(1^{a_1} 2^{a_2} \dots) \prod_{j \geq 1} \frac{(j^j f_j / j!)^{a_j}}{a_j!},$$

and part (b) follows by summing this over all $a_1, a_2, \dots \geq 0$ with $a_1 + a_2 + \dots = m$, and $m \geq 1$, and applying (3.16).

For part (c), for $j \geq 1$, variant Ω_0^{br} of the branch decomposition gives a bijection

$$\Omega_0^{\text{br}} : \mathcal{T}_{0,j} \xrightarrow{\sim} \bigcup \{ \mathfrak{w}_\gamma \} \star \mathcal{U}_{b_1}(\mathcal{T}_{\infty,1}) \star \mathcal{U}_{b_2}(\mathcal{T}_{\infty,2}) \star \dots : \mathfrak{t} \mapsto (\alpha_0(\mathfrak{w}_\gamma), \alpha_1(t'_1), \dots, \alpha_m(t'_m)),$$

where the union is over all $\gamma = (1^{a_1} 2^{a_2} \dots)$, $\beta = (1^{b_1} \dots (j-1)^{b_{j-1}} j^{b_j+1} (j+1)^{b_{j+1}} \dots)$ with $|\gamma| = |\beta| \geq j$, and $l(\beta) = m+1$, $m \geq 0$ (here, $\gamma = \gamma^u(\mathfrak{t})$ and $\beta = \beta^u(\mathfrak{t})$, where u is the ∞ -vertex adjacent to \bullet in \mathfrak{t} , as given in the description of Ω_0^{br} above). It is straightforward to check in this bijection that $\eta_k(\mathfrak{t}) = \eta_k(\mathfrak{w}_\gamma) + \eta_k(t'_1) + \dots + \eta_k(t'_m)$, for $k \geq 0$. Moreover, $\text{wt}_0(\mathfrak{t}) = j (H_{\gamma,\beta}^0 u^{l(\beta)-2} z^{|\beta|} / r_{\gamma,\beta}^0!) \text{wt}_\infty(t'_1) \dots \text{wt}_\infty(t'_m)$, so from the Product Lemma we obtain

$$f_j = u^{-2} j \sum \frac{H_{\gamma,\beta}^0 u^{l(\beta)-2} z^{|\beta|}}{r_{\gamma,\beta}^0!} \prod_{i \geq 1} \frac{p_i^{a_i} g_i^{b_i}}{a_i! b_i!},$$

since $[\mathfrak{w}_\gamma, 1]_\eta = \prod_{i \geq 1} \frac{p_i^{a_i}}{a_i!}$, and part (c) follows immediately from (4.1).

For part (d), variant $\Omega_\infty^{\text{br}}$ of the branch decomposition gives a bijection

$$\Omega_\infty^{\text{br}}: \mathcal{T}_{\infty, j} \xrightarrow{\sim} \bigcup \{ \mathfrak{v}_1 \} \star \mathcal{U}_{a_1}(\mathcal{T}_{0,1}) \star \mathcal{U}_{a_2}(\mathcal{T}_{0,2}) \star \cdots : \mathfrak{t} \mapsto (\alpha_0(\mathfrak{v}_1), \alpha_1(\mathfrak{t}'_1), \dots, \alpha_m(\mathfrak{t}'_m)),$$

where the union is over all $\delta = (1^{a_1} \cdots (j-1)^{a_{j-1}} j^{a_j+1} (j+1)^{a_{j+1}} \cdots)$ with $|\delta| \geq j$, and $l(\delta) = m+1$, $m \geq 0$. (Here, $\delta = \delta^v(\mathfrak{t})$, where v is the 0-vertex adjacent to \bullet in \mathfrak{t} , as given in the description of $\Omega_\infty^{\text{br}}$ above.) It is straightforward to check in this bijection that $\eta_k(\mathfrak{t}) = \eta_k(\mathfrak{v}_1) + \eta_k(\mathfrak{t}'_1) + \cdots + \eta_k(\mathfrak{t}'_m)$, for $k \geq 0$, and that $\text{wt}_\infty(\mathfrak{t}) = j (H_\delta^0 / r_\delta^0!) \text{wt}_0(\mathfrak{t}'_1) \cdots \text{wt}_0(\mathfrak{t}'_m)$. Then from the Product Lemma we obtain

$$g_j = j \frac{1}{1!} \sum \frac{H_\delta^0}{r_\delta^0!} \prod_{i \geq 1} \frac{f_i^{a_i}}{a_i!},$$

since $[\{\mathfrak{v}_1\}, 1]_\eta = 1/1!$, and part (d) follows immediately from (3.11). ■

5. Symmetrization and a Polynomial Transformation

At this stage, we have established the two generating series results that we need to prove Faber’s Conjecture. The first of these results, on the degeneration side, is Corollary 3.9, the second form of the Degeneration Theorem, which gives a linear partial differential equation for the Faber-Hurwitz series F^g , in terms of the genus 0 Hurwitz series \widehat{H}^0 . The second of these results, on the localization side, is Theorem 4.4, the second form of the Localization Tree Theorem, which expresses Faber-Hurwitz classes as a linear combination of the intersections $\langle \tau_{a_1} \cdots \tau_{a_n} \lambda_k \rangle_g^{\mathbb{F}^{\text{ab}}}$, in terms of \widehat{H}^0 and the genus 0 double Hurwitz series H^0 .

In this section, in order to prove Faber’s Intersection Number Conjecture, we introduce three operators, giving us a three-step transformation that will enable us to apply Corollary 3.9 and Theorem 4.4 conveniently, so that we can extract the intersection numbers of top degree, that are the subject of Faber’s Conjecture. The first step is a symmetrization operator Ξ_m , and the second step is a change of variables C . The composition $C\Xi_m$ yields polynomials in the new variables, and the third step is the operator T' , that restricts to terms of maximum total degree. We shall refer to $T'C\Xi_m = T\Xi_m$ as the *fundamental transformation*.

Note that our fundamental transformation is only a slight modification of the three-step transformation that was used in [22] to give another proof of Getzler and Pandharipande’s λ_g -Conjecture. There the transformation was applied to the Hurwitz number generating series in arbitrary genus, and the first two steps, of symmetrizing and changing variables, were identical. In that Hurwitz case, polynomiality also held, but the third step was to restrict to (full) terms of *minimum* total degree, so that the intersection numbers of bottom degree, that are the subject of the λ_g -Conjecture, could be extracted. Changes of variables for similar purposes arise in work of Kazarian and Lando [30] and Shadrin and Zvonkine [43], as well as in [18–21].

5.1. Symmetrization

The first step of the fundamental transformation is the linear symmetrization operator Ξ_m .

Definition of Ξ_m (Step 1): Following [21], we define

$$\Xi_m \left(p_\alpha z^{|\alpha|} \right) := \sum_{\sigma \in \mathfrak{S}_m} x_{\sigma(1)}^{\alpha_1} \cdots x_{\sigma(m)}^{\alpha_m}, \quad m \geq 1, \tag{5.1}$$

if $l(\alpha) = m$ (with $\alpha = (\alpha_1, \dots, \alpha_m)$), and zero otherwise.

In order to provide more compact expressions when applying Ξ_m , some notation is required. For $\rho = \{\rho_1, \dots, \rho_l\} \subseteq \{1, \dots, m\}$, let $x_\rho = \{x_{\rho_1}, \dots, x_{\rho_l}\}$. Given $i_1, \dots, i_k \geq 1$, with $i_1 + \dots + i_k = m$, let $s_j = i_1 + \dots + i_j$, for $j = 1, \dots, k$, and $s_0 = 0$. Also let

$$\rho_j = \{s_{j-1} + 1, \dots, s_j\}, \tag{5.2}$$

for $j = 1, \dots, k$. Then $\text{sym}_{i_1, \dots, i_k}$ is a summation operator, over the set of ordered set partitions (ν_1, \dots, ν_k) of $\{1, \dots, m\}$, in which, in the summand, x_{ρ_j} is replaced by x_{ν_j} , for $j = 1, \dots, k$.

We require the symmetrized series

$$\begin{cases} F_m^g(x_1, \dots, x_m) & := \Xi_m F^g(z; \mathbf{p}), & \widehat{\mathbf{H}}_m^0(x_1, \dots, x_m) & := \Xi_m \widehat{H}^0(z; \mathbf{p}), \\ \mathbb{T}_m^g(x_1, \dots, x_m, u) & := \Xi_m \mathbb{T}^g(z, u; \mathbf{p}), & \xi_m^{(i)}(x_1, \dots, x_m, u) & := \Xi_m \xi^{(i)}(z, u; \mathbf{p}), \end{cases} \tag{5.3}$$

for $m \geq 1$. When the arguments of these series are suppressed, they are the ones stated above.

5.2. Symmetrizing on the Localization and Degeneration Sides

On the localization side, the following result is the third form of the Localization Tree Theorem. It is the symmetrized form of Theorem 4.4, and uses the substitution operator

$$\Lambda: u \mapsto -u, x_i \mapsto (1 - u)^{-1} x_i, \quad i = 1, \dots, m. \tag{5.4}$$

Corollary 5.1. (Localization Tree Theorem — Symmetrized Functional Equations)
For $m, g \geq 1$,

$$(a) \quad F_m^g \mathbb{G}_{g,1} = [u^{2g-1}] \left(\frac{-u}{1-u} \right)^m \Lambda \mathbb{T}_m^g,$$

where

$$(b) \quad \mathbb{T}_m^g = \sum_{n \geq 1} \frac{1}{n!} \sum_{\substack{a_1, \dots, a_n, k \geq 0, \\ a_1 + \dots + a_n + k = g - 2 + n, \\ i_1, \dots, i_n \geq 1, \\ i_1 + \dots + i_n = m}} (-1)^k \langle \tau_{a_1} \cdots \tau_{a_n} \lambda_k \rangle_g^{\text{Fab}} \text{sym}_{i_1, \dots, i_n} \prod_{j=1}^n \xi_{i_j}^{(a_j)}(x_{\rho_j}).$$

Proof. The result follows by applying Ξ_m to Theorem 4.4(a) and (b), for $m \geq 1$, and using the properties of Ξ_m given in Lemmas 4.1–4.3 of [19]. ■

On the degeneration side, the following result is the symmetrized form of Corollary 3.9. For each $1 \leq i < j \leq m$, note that there are two terms in the result with denominator $x_i - x_j$, and that the numerator is antisymmetric in x_i, x_j , so these terms combine to give a formal power series. To account for each individual term $(x_i - x_j)^{-1}$ that arises in the statement, we adopt the total ordering $x_i \prec x_j$ if $i < j$, and then define $(x_i - x_j)^{-1}$ by $(x_i - x_j)^{-1} := x_j^{-1} (1 - x_i x_j^{-1})^{-1}$. This gives us an ordered Laurent series ring in x_1, \dots, x_m .

Corollary 5.2. (Degeneration Theorem — Symmetrized Join-Cut Equation) *For $m, g \geq 1$,*

$$\begin{aligned} & \left(\sum_{i=1}^m x_i \frac{\partial}{\partial x_i} + m - 1 \right) F_m^g \\ &= \sum_{\substack{i_2, i_3 \geq 0, \\ i_2 + i_3 = m - 1}} \text{sym}_{1, i_2, i_3} \left(x_1 \frac{\partial}{\partial x_1} \widehat{\mathbf{H}}_{1+i_2}^0(x_1, x_{p_2}) \right) \left(x_1 \frac{\partial}{\partial x_1} F_{1+i_3}^g(x_1, x_{p_3}) \right) \\ & \quad + \text{sym}_{1, 1, m-2} \frac{x_2}{x_1 - x_2} x_1 \frac{\partial}{\partial x_1} F_{m-1}^g(x_1, x_{p_3}) + \sum_{i=1}^m \left(x_i \frac{\partial}{\partial x_i} \right)^{2g+1} \widehat{\mathbf{H}}_m^0. \end{aligned}$$

Proof. The result follows by applying Ξ_m to Corollary 3.9 for $m \geq 1$, and using the properties of Ξ_m given in Lemmas 4.1–4.3 of [19]. ■

Note that the second term on the right hand side of Corollary 5.2 vanishes in the case $m = 1$.

5.3. Polynomiality and Terms of Top Degree

The second step of the fundamental transformation is the change of variables C .

Definition of C (Step 2). Let

$$a) \quad w(x) := \sum_{n \geq 1} \frac{n^{n-1}}{n!} x^n, \quad b) \quad y(x) := \frac{1}{1 - w(x)} = \sum_{n \geq 0} \frac{n^n}{n!} x^n, \quad (5.5)$$

where the second equality in the expression for $y(x)$ is well known (see, e.g., [18, Prop. 3.2.1]). Let $w_i := w(x_i)$ and $y_i = y(x_i)$, for $i = 1, \dots, m$, and let C be an operator, applied to a formal power series in x_1, \dots, x_m , that changes variables from the indeterminates x_1, \dots, x_m to y_1, \dots, y_m . Thus, a direct way to apply C is to substitute $x_i = G(y_i - 1)$, where $G(z)$ is the compositional inverse of the formal power series $\sum_{n \geq 1} n^n z^n / n!$. In this paper, we do *not* apply this substitution directly to the symmetrized series $F_m^g, \Lambda \mathbb{T}_m^g, \Lambda \xi_m^{(i)}$, but instead quite indirectly. The details are intricate, but the key to our method is that in all cases the result is a *polynomial* in y_1, \dots, y_m (we say that a Laurent series in another indeterminate, either u or t in this paper, is *polynomial* if each of its coefficients is polynomial).

This key fact is recorded in the following result. The proof involves a number of transformations for implicitly defined series, and is deferred until Appendix A to

avoid interrupting the present development. Note that part (b) follows from part (a) by Theorem 4.4(a), (b), and by (3.10) and Theorem 3.12 (which allows us to “divide” $\langle \tau_{a_1} \cdots \tau_{a_n} \lambda_k \rangle_g^{\text{Fab}}$ by $\mathbb{G}_{g,1}$). Consequently, in the Appendix it suffices to prove part (a) of Theorem 5.3 only.

Theorem 5.3. *For $m = 1, 2, 3$, we have*

$$(a) \ C \Lambda \xi_m^{(i)} \in \mathbb{Q}[y_1, \dots, y_m]((u)), \quad i \geq 1, \quad (b) \ C F_m^g \in \mathbb{Q}[y_1, \dots, y_m], \quad g \geq 1.$$

Theorem 5.3 is essential to our proof of Faber’s Intersection Conjecture for at most 3 parts. In order for us to extend our proof to, say, n parts, we would first need to prove Theorem 5.3 for all $m \leq n$. We are presently able to prove this result for $m \leq 5$, but not for larger values of m , since we require the symmetrized double Hurwitz series in our method of proof, and we only have explicit expressions for this when $m \leq 5$, as given in [21]. We conjecture that Theorem 5.3 holds for all positive integers m .

The third and final step of the fundamental transformation is the operator T' , that restricts a polynomial to terms of maximum total degree (these are referred to as the “top” terms).

Definition of T' (Step 3). Let S_i be the operator that restricts a polynomial in y_1, \dots, y_m to the terms of total degree i . Then we define T' , when applied to $C F_m^g$, to denote $S_{4g+3m-5}$. It turns out that there are no terms of higher degree in $C F_m^g$, so we say that T' restricts to the terms of top degree, though we understand that this is informal, since it assumes that the terms of total degree $4g + 3m - 5$ are not all zero.

If $C \Lambda \xi_m^{(i)} = \sum_k a_k u^k$, we define $T' C \Lambda \xi_m^{(i)} := \sum_k (S_{2g+2m-2+k} a_k) u^k$. Again, it turns out that there are no terms in any of the a_k of higher total degree, so in this case also T' restricts to terms of top degree. We define T' , when applied to $C \widehat{H}_m^0$, to denote S_{3m-6} . Again there are no terms of higher total degree, so in this case also T' restricts to terms of top degree.

Finally, in all other cases, we define T' as a homomorphism, and we define $T := T' C$.

5.4. Polynomiality and Terms of Top Degree on the Localization and Degeneration Sides

On the localization side, we can apply the fundamental transformation via Theorem 4.4, because of the polynomiality of $\Lambda \xi_m^{(i)}$ that was established in Theorem 5.3(a) for $m = 1, 2, 3$. This requires the top terms, $T \Lambda \xi_m^{(i)}$, which are given in the following result for $m = 1, 2, 3$. Again, the proof is intricate, and is deferred until Appendix A to avoid interrupting the present development. The result uses the notation

$$Y_i(u) := \frac{y_i}{1 - uy_i}, \quad i \geq 1. \tag{5.6}$$

Theorem 5.4. For $i \geq 0$,

- (a) $u \tau \Lambda \xi_1^{(i)} = -(2i - 1)!! Y_1(u)^{2i+1},$
- (b) $u^2 \tau \Lambda \xi_2^{(i)} = -(2i + 1)!! u \operatorname{sym}_{1,1} \frac{y_1^2 y_2}{y_1 - y_2} Y_1(u)^{2i+3},$
- (c) $\frac{(-u)^3 \tau \Lambda \xi_3^{(i)}}{(2i + 1)!!} = u^2 \operatorname{sym}_{1,1,1} \frac{y_1^3 y_2^4 y_3}{(y_2 - y_3)(y_1 - y_2)^2} Y_1(u)^{2i+3} Y_2(u)$
 $- \operatorname{sym}_{1,2} \left(u y_1 Y_1(u)^{2i+5} Y_2(u) Y_3(u) - u^2 y_1^3 \frac{\partial}{\partial y_1} \frac{y_1^3 y_2 y_3}{(y_2 - y_1)(y_3 - y_1)} Y_1(u)^{2i+4} \right).$

On the degeneration side, we can apply the fundamental transformation via Corollary 5.2 if we can apply the change of variables C to the partial differential operator $x_i \frac{\partial}{\partial x_i}$ and to the symmetrized generating series \widehat{H}_m^0 for Hurwitz numbers, genus 0. This is straightforward in both cases.

For the partial differential operator, from a) and b) of (5.5) we have the functional equations $y_i = 1/(1 - w_i)$, $y_i = 1 + x_i \frac{dw_i}{dx_i}$, and together these imply $x_i \frac{dy_i}{dx_i} = y_i^2 (y_i - 1)$. This immediately implies the operator identities

$$a) C x_i \frac{\partial}{\partial x_i} = y_i^2 (y_i - 1) \frac{\partial}{\partial y_i} C, \quad b) \tau x_i \frac{\partial}{\partial x_i} = y_i^3 \frac{\partial}{\partial y_i} \tau. \tag{5.7}$$

For the symmetrized generating series \widehat{H}_m^0 , (3.8) and (5.5) b) give

$$\widehat{H}_m^0 = \left(\sum_{j=1}^m x_j \frac{\partial}{\partial x_j} \right)^{m-3} \prod_{i=1}^m (y_i - 1), \tag{5.8}$$

for $m \geq 1$ (note that (5.7) applied to (5.8) immediately identifies $C \widehat{H}_m^0$ as having degree $3m - 6$, consistent with the definition of T' above). Also, we have

$$a) x_1 \frac{\partial}{\partial x_1} \widehat{H}_1^0 = 1 - y_1^{-1}, \quad b) \widehat{H}_2^0 = \log \left(\frac{y_1 - y_2}{y_1 y_2 (x_1 - x_2)} \right) + y_1^{-1} + y_2^{-1} - 2, \tag{5.9}$$

where the first of these expressions arises by applying $x_1 \frac{d}{dx_1}$ to (5.8) when $m = 1$, and using (5.7) a), and the second is given in [19, p. 38].

6. Intersection Numbers with One Part

In this section, we use our strategy for the first time. This involves applying the fundamental transformation to give results for the symmetrized Faber-Hurwitz series F_m^g on both the degeneration and localization sides, to obtain results for Faber’s intersection numbers. Here, these results are given in Section 6.1 for the case of a single ($m = 1$) part. Faber’s Intersection Number Conjecture, giving $\langle \tau_{g-1} \rangle_g^{\text{Fab}} = \Psi_1^{g-1}$, is immediate in this case, but it gives us an equation relating the two generators Ψ_1^{g-1} and $\mathbb{G}_{g,1}$ of $A_{2g-1}(\mathcal{M}_{g,n}^{rt})$. There is non-trivial geometric information to be gained by relating these generators, which is described in Section 6.2.

6.1. Intersection Numbers with One Part

We begin with a result for the localization side.

Theorem 6.1. *For $g \geq 1$, we have $\mathbb{T}F_1^g(x_1)\mathbb{G}_{g,1} = (2g - 3)!! \binom{4g-3}{2g-1} y_1^{4g-2} \Psi_1^{g-1}$.*

Proof. From Corollary 5.1 (b) we have $\mathbb{T}_1^g = \sum_{k=0}^{g-1} (-1)^k \langle \tau_{g-1-k} \lambda_k \rangle_g^{\mathbb{F}ab} \xi_{\mathbb{S}_1}^{(g-1-k)}$. Together with Theorem 5.4 (a), this gives

$$\begin{aligned} [u^{2g-1}] (-u)^1 \mathbb{T} \Lambda \mathbb{T}_1^g(x_1, u) &= [u^{2g-1}] \langle \tau_{g-1} \rangle_g^{\mathbb{F}ab} (2g - 3)!! Y_1(u)^{2g-1} \\ &= [u^{2g-1}] (2g - 3)!! Y_1(u)^{2g-1} \Psi_1^{g-1}, \end{aligned}$$

where, for the second equality, we have used the immediate fact $\langle \tau_{g-1} \rangle_g^{\mathbb{F}ab} = \Psi_1^{g-1}$ for $g \geq 1$. But $Y_1(u)^{2g-1} = \sum_{i \geq 0} \binom{-(2g-1)}{i} (-1)^i y_1^{2g-1+i} u^i = \sum_{i \geq 0} \binom{2g-2+i}{i} y_1^{2g-1+i} u^i$, and the result follows immediately. ■

Now we turn to the corresponding result for the degeneration side.

Theorem 6.2. *For $g \geq 1$, we have $\mathbb{T}F_1^g(x_1) = (4g - 3)!! \frac{y_1^{4g-2}}{4g-2}$.*

Proof. Let $m = 1$ in Corollary 5.2, to obtain $x_1 \frac{\partial}{\partial x_1} F_1^g = (x_1 \frac{\partial}{\partial x_1} \widehat{\mathbf{H}}_1^0) (x_1 \frac{\partial}{\partial x_1} F_1^g) + (x_1 \frac{\partial}{\partial x_1})^{2g+1} \widehat{\mathbf{H}}_1^0$. Solving for $x_1 \frac{\partial}{\partial x_1} F_1^g$, and using (5.9) a), we obtain

$$x_1 \frac{\partial}{\partial x_1} F_1^g = y_1 \left(x_1 \frac{\partial}{\partial x_1} \right)^{2g-1} (y_1 - 1). \tag{6.1}$$

The polynomiality of F_1^g follows immediately from (5.7) a) (or we can use Theorem 5.3 (b), with $m = 1$), so from (5.7) b) we have $y_1^3 \frac{\partial}{\partial y_1} \mathbb{T}F_1^g = y_1 (y_1^3 \frac{\partial}{\partial y_1})^{2g-1} y_1 = (4g - 3)!! y_1^{4g}$, where the second equality follows by induction. The result follows by integrating in y_1 . ■

6.2. Some Immediate Geometric Consequences

Comparing the above two results, we obtain the following result, which relates the generator $\mathbb{G}_{g,1}$ of $A_{2g-1}(\mathcal{M}_{g,n}^{rt})$ with the generator Ψ_1^{g-1} suggested by Faber’s Intersection Number Conjecture.

Corollary 6.3. *For $g \geq 1$, $\mathbb{G}_{g,1} = \frac{2^g}{(g-1)!} \Psi_1^{g-1}$ on $A_{2g-1}(\mathcal{M}_{g,1})$.*

In particular, $\Psi_1^{g-1} = \langle \tau_{g-1} \rangle_g^{\mathbb{F}ab}$ is another non-zero element (=basis) of the one-dimensional vector space $A_{2g-1}(\mathcal{M}_{g,1})$, and this describes the change of basis from $\mathbb{G}_{g,1}$ to Ψ_1^{g-1} .

Proof. Comparing Theorems 6.1 and 6.2, we obtain that $(2g - 3)!! \binom{4g-3}{2g-1} \Psi_1^{g-1} = \frac{(4g-3)!!}{4g-2} \mathbb{G}_{g,1}$, and the result follows immediately. ■

Corollary 6.4. ([40, Thm. 4]) $\sum_{i=0}^{g-2} (-1)^i \lambda_i \Psi_1^{g-1-i} = 2^{g-1} \Psi_1^{g-1} / g!$.

Proof. Applying the usual (non-relative) virtual localization formula to $\mathbb{G}_{g,1,\sim}$, we obtain $\mathbb{G}_{g,1,\sim} = \psi_1^{2g-1} - \psi_1^{2g-2}\lambda_1 + \dots + (-1)^g \psi_1^{g-1}\lambda_g$. The result then follows from Corollary 6.3 and the fact that $\mathbb{G}_{g,1} = 2g\mathbb{G}_{g,1,\sim}$ (Proposition 3.1). ■

Corollary 6.5. *The class of the hyperelliptic curves in \mathcal{M}_g is $\frac{(2^{2g}-1)2^{g-1}}{g!(2g+1)(2g+2)}\kappa_{g-2}$.*

(This was stated, for example, in the concluding remarks of [7].) The argument carries through for the locus of degree d covers fully ramified over two points.

Sketch of Proof. The class $\mathbb{G}_{g,2,\sim}$ corresponds to the points $[C, p, q] \in \mathcal{M}_{g,2}^{rt}$ where $O_C(p - q)$ is a 2-torsion point (along with a virtual class). The locus where $O_C(p - q)$ is trivial corresponds to $\mathbb{G}_{g,1}$. Thus the locus where $O_C(p - q)$ is 2-torsion but non-trivial has virtual class $(2^{2g} - 1)\mathbb{G}_{g,1}$ by Theorem 3.5. It is straightforward to check the virtual fundamental class on this locus is the actual fundamental class (as mentioned in the proof of Theorem 3.5). This locus corresponds to hyperelliptic curves with two marked Weierstrass points. Thus the locus in $\mathcal{M}_{g,1}$ of hyperelliptic curves with choice of one Weierstrass point has class $\frac{2^{2g}-1}{2g+1}\mathbb{G}_{g,1} = \frac{(2^{2g}-1)2^{g-1}}{g!(2g+1)}\psi_1^{g-1}$. Pushing forward to \mathcal{M}_g , and forgetting the choice of one of the $2g + 2$ Weierstrass points, we get the desired result. ■

Corollary 6.6. *For $d, g \geq 1$, we have $F_{(d)}^g = \frac{1}{d} \sum_{i=1}^d \binom{d}{i} i^{2g+i-1} (d-i)^{d-i}$.*

This is a corollary of the proof of Theorem 6.2, not of the result itself.

Proof. From (6.1) and (5.5) b), we have $x_1 \frac{\partial}{\partial x_1} F_1^g = \sum_{n \geq 0} \frac{n^n}{n!} x_1^n \sum_{i \geq 1} \frac{i^{2g+i-1}}{i!} x_1^i$, and the result follows from (5.3) and (3.12). ■

Thus, for example, the class of non-singular genus g curves C admitting a degree d cover of \mathbb{P}^1 via $O(dp)$ for some $p \in C$, ramified over an appropriate number of points is

$$\frac{1}{d} \sum_{i=1}^d \binom{d}{i} i^{2g+i-1} (d-i)^{d-i} \mathbb{G}_{g,1} = \frac{1}{d} \sum_{i=1}^d \binom{d}{i} i^{2g+i-1} (d-i)^{d-i} \frac{2^g \kappa_{g-2}}{(g-1)!} \in A_{2g-1}(\mathcal{M}_g).$$

7. A Strategy for an Inductive Proof of Faber’s Intersection Number Conjecture

In this section, we consider an inductive strategy for proving the Conjecture.

7.1. Final Form of the Localization Tree Theorem

First, we define two generating series, in which the genus g is marked by the indeterminate t .

The localization tree generating series is

$$\Psi_m(y_1, \dots, y_m, t) := \sum_{g \geq 1} \frac{t^{2g}}{(2g-1)!!} [u^{2g-1}] (-u)^m \mathbb{T} \wedge \mathbb{T}_m^g(x_1, \dots, x_m, u) / \psi_1^{g-1}, m \geq 1. \tag{7.1}$$

The Faber generating series is

$$\Phi_m(y_1, \dots, y_m, t) := \sum_{g \geq 1} 2^{2g-1} \frac{t^{2g}}{(2g-1)!} \top F_m^g(x_1, \dots, x_m), \quad m \geq 1. \quad (7.2)$$

For Ψ_m , thanks to the fact that $R_{2g-1}(\mathcal{M}_{g,1}) \cong \mathbb{Q}$ (Thm. 3.12), we “divide by ψ_1^{g-1} ” using the isomorphism

$$\mathbb{Q} \xrightarrow{\times \psi_1^{g-1}} R_{2g-1}(\mathcal{M}_{g,1})$$

(combining (3.20) and Cor. 6.3), and thus we define the *Faber intersection number*

$$\langle \dots \rangle_g^{\text{Fab}} := \langle \dots \rangle_g^{\mathbb{F}\text{ab}} / \psi_1^{g-1} = \langle \dots \rangle_g^{\mathbb{F}\text{ab}} / \langle \tau_{g-1} \rangle_g^{\mathbb{F}\text{ab}}. \quad (7.3)$$

Note that Ψ_m and Φ_m , since they require the operator \top , are only defined when the operator \mathbb{C} yields a polynomial. So far, we have proved this polynomiality only for $m = 1, 2, 3$, in Theorem 5.3. Subject to this, Ψ_m and Φ_m are both generating series with rational coefficients, and the next result shows that they are equal.

Corollary 7.1. (Localization Tree Theorem — for Genus Generating Series) *For $m \geq 1$,*

$$\Psi_m(y_1, \dots, y_m, t) = \Phi_m(y_1, \dots, y_m, t).$$

Proof. The result follows from Corollary 5.1 (a) by noting that $\frac{\mathbb{G}_{g,1}}{(2g-1)!!} = \frac{2^g \psi_1^{g-1}}{(g-1)!(2g-1)!!}$
 $= \frac{2^{2g-1} \psi_1^{g-1}}{(2g-1)!}$ for $g \geq 1$, from Corollary 6.3. ■

Corollary 7.1 is the final form of Theorem 3.11. It summarizes the relationship between the localization side (dealing with Ψ_m) and the degeneration side (dealing with Φ_m) as the equality of Ψ_m and Φ_m . Our strategy for exploiting this relationship in order to prove the Faber Conjecture for a fixed number of parts is indirect. We prove in the following lemma that Corollary 7.1 implies sufficiently many linearly independent linear equations for Faber’s intersection numbers to uniquely identify them. This means that we can prove the Faber Conjecture iteratively on the number of parts, by simply verifying the equality $\Psi_m = \Phi_m$ of Corollary 7.1 with Faber’s conjectured values substituted for the intersection numbers. In fact, as we shall see, with these substituted values, we are able to verify the equality $\Psi_m = \Phi_m$ for particular m simply by equating polynomials.

Lemma 7.2. *For any $n \geq 2$, if the Faber Conjecture is true for up to $n - 1$ parts, and*

$$\Psi_n(y_1, \dots, y_n, t) = \Phi_n(y_1, \dots, y_n, t)$$

with Faber’s conjectured values substituted for the intersection numbers with up to n parts, then the Faber Conjecture is true for up to n parts.

Proof. Consider the equality $\Psi_n(y_1, \dots, y_n, t) = \Phi_n(y_1, \dots, y_n, t)$, and equate coefficients of t^{2g} for each fixed $g \geq 2$. This gives an equation involving Faber’s intersection numbers with at most n parts, and from Corollary 5.1(b), (7.1), and Theorem 5.4(a), we deduce that the only terms involving the intersection numbers with exactly n , positive, parts are given by

$$\begin{aligned} & \sum_{\substack{a_1, \dots, a_n \geq 1 \\ a_1 + \dots + a_n = g + n - 2}} \langle \tau_{a_1} \cdots \tau_{a_n} \rangle_g^{\text{Fab}} \frac{1}{(2g-1)!!} [u^{2g-1}] \prod_{j=1}^n (2a_j - 1)!! Y_j(u)^{2a_j+1} \\ &= \sum_{\substack{a_1, \dots, a_n \geq 1 \\ a_1 + \dots + a_n = g + n - 2}} \sum_{\substack{m_1, \dots, m_n \geq 0 \\ m_1 + \dots + m_n = 2g-1}} \langle \tau_{a_1} \cdots \tau_{a_n} \rangle_g^{\text{Fab}} \frac{\prod_{j=1}^n (2a_j - 1)!! \binom{2a_j + m_j}{m_j} y_j^{2a_j + m_j + 1}}{(2g-1)!!}. \end{aligned} \tag{7.4}$$

Now under the hypothesis that the Faber Conjecture has been proved for up to $n - 1$ parts, we are able to evaluate all Faber intersection numbers in the above equation with up to $n - 1$ parts, as well as those with n parts, at least one of which is zero (by use of the string equation), and we consider Faber’s intersection numbers with exactly n , positive, parts as unknowns. Now equate coefficients of $y_1^{i_1} \cdots y_n^{i_n}$ in this equation for each $i_1, \dots, i_n \geq 3$ with $i_1 + \dots + i_n = 4g - 5 + 3n$, to obtain a linear equation that we shall refer to as B_{i_1, \dots, i_n} . Then we thus obtain a system of $\binom{4g-6+n}{n-1}$ equations B_{i_1, \dots, i_n} in the $\binom{g-3+n}{n-1}$ “unknowns” $\langle \tau_{a_1} \cdots \tau_{a_n} \rangle_g^{\text{Fab}}$, where $a_1, \dots, a_n \geq 1$, with $a_1 + \dots + a_n = g - 2 + n$. Let the coefficient matrix for this system be denoted by \mathbf{M} , and note that $\binom{4g-6+n}{n-1} \geq \binom{g-3+n}{n-1}$, since $g, n \geq 2$. (We will not use the symmetry of these unknowns.)

Now we prove that this system has at most one solution. To do so, suppose that we consider the unknowns in lexicographic order of the $a_1 \cdots a_n$, and the equations in lexicographic order of the $i_1 \cdots i_n$ (so we have ordered the rows and columns of \mathbf{M}). Note that, from (7.4), the entry in row $i_1 \cdots i_n$ and column $a_1 \cdots a_n$ is non-zero if and only if $i_j \geq 2a_j + 1$ for all $j = 1, \dots, n$. Then it is straightforward to check that the submatrix of \mathbf{M} consisting of all columns, and rows i_1, \dots, i_n with $i_1, \dots, i_{n-1} \geq 3$ and odd, and $i_1 + \dots + i_{n-1} \leq 2g - 7 + 3n$, is lower triangular, with non-zero coefficients on the diagonal. Thus these rows of \mathbf{M} are linearly independent, so the system has at most one solution. The result follows since, if Faber’s conjectured values satisfy this system, then they are uniquely the correct values for these intersection numbers. ■

7.2. A Lemma for the Localization Tree Generating Series

We consider first the localization side. In order to determine a compact rational expression for the localization tree generating series Ψ_m , we shall need the following lemma, where Ef denotes the *even* subseries of the formal power series f in the indeterminate t . Also, for $i \geq 1$ we let

$$B_i := \sqrt{1 - 4y_i^2 t}, \quad A_i := \frac{1 - B_i}{2y_i}. \tag{7.5}$$

Lemma 7.3. *For a formal power series f , we have $\sum_{g \geq 1} t^{2g} [u^{2g-1}] f(u) Y_1(u)^{2g-1} = \frac{1}{2} \text{E}t (1 + B_1^{-1}) f(A_1)$.*

Proof. Now $[u^{2g-1}] f(u)Y_1(u)^{2g-1} = [t^{2g-1}] \frac{f(s)}{1-t\frac{d}{ds}Y_1(s)}$, from Lagrange’s Implicit Function Theorem (see, e.g., [17, Thm. 1.2.4]), where s is the unique formal power series solution (in t) of $s = tY_1(s)$, which is the equation $y_1s^2 - s + y_1t = 0$, and so $s = A_1$. But $\frac{d}{ds}Y_1(s) = Y_1(s)^2$, so with $s = A_1$, we have $1 - t\frac{d}{ds}Y_1(s) = \frac{2B_1}{1+B_1}$. Then $[u^{2g-1}] f(u)Y_1(u)^{2g-1} = [t^{2g-1}] \frac{1}{2} (1 + B_1^{-1}) f(A_1)$. The result follows. ■

7.3. Final Form of the Degeneration Theorem

For the degeneration side, we describe the general results that will allow us to obtain an explicit rational expression for the Faber generating series Φ_m . First we give a compact expression for Φ_1 .

Corollary 7.4. $y_1 \frac{\partial}{\partial y_1} \Phi_1(y_1, t) = EtB_1^{-1}$.

Proof. From (7.2) with $m = 1$, and Theorem 6.2, we obtain

$$y_1 \frac{\partial}{\partial y_1} \Phi_1(y_1, t) = \sum_{g \geq 1} 2^{2g-1} \frac{(4g-3)!!}{(2g-1)!} y_1^{4g-2} t^{2g} = t \sum_{g \geq 1} \binom{-\frac{1}{2}}{2g-1} (-4y_1^2 t)^{2g-1},$$

and the result follows immediately. ■

In the next result, we consider the top terms in Corollary 5.2, to obtain a simple iterative equation for the Faber generating series Φ_m , $m \geq 2$. Similarly to Corollary 5.2, there are two terms in this result with denominator $y_i - y_j$, and the numerator is antisymmetric in y_i, y_j , so these terms combine to give a formal power series. Also, to account for each individual term $(y_i - y_j)^{-1}$, we adopt the total ordering $y_i \prec y_j$ if $i < j$, to obtain an ordered Laurent series ring in y_1, \dots, y_m .

The statement of the result uses the operator

$$\Delta_k := \sum_{i \geq 1} y_i^k \frac{\partial}{\partial y_i}, \quad k \geq 2, \tag{7.6}$$

and the generating series

$$\Upsilon_m(y_1, \dots, y_m, t) := \sum_{g \geq 1} 2^{2g-1} \frac{t^{2g}}{(2g-1)!} \top \sum_{i=1}^m \left(x_i \frac{\partial}{\partial x_i} \right)^{2g+1} \widehat{\mathbf{H}}_m^0(x_1, \dots, x_m), \quad m \geq 1. \tag{7.7}$$

Lemma 7.5. (Degeneration Theorem — Join-Cut Equation for Φ_m) *For $m \geq 2$, we have*

$$\begin{aligned} & \Delta_2 \Phi_m(y_1, \dots, y_m, t) \\ &= \sum_{\substack{i_2 \geq 2, i_3 \geq 0, \\ i_2 + i_3 = m-1}} \text{sym}_{1, i_2, i_3} \left(y_1^3 \frac{\partial}{\partial y_1} \top \widehat{\mathbf{H}}_{1+i_2}^0(x_1, x_{p_2}) \right) \left(y_1^3 \frac{\partial}{\partial y_1} \Phi_{1+i_3}(y_1, y_{p_3}, t) \right) \\ &+ \text{sym}_{1, 1, m-2} \frac{y_1^5 y_2}{y_1 - y_2} \frac{\partial}{\partial y_1} \Phi_{m-1}(y_1, y_{p_3}, t) + \Upsilon_m(y_1, \dots, y_m, t). \end{aligned}$$

Proof. From the symmetrized Join-cut equation (Corollary 5.2), and moving the contribution for $i_2 = 0$ from the summation on the right hand side to the left hand side, and using (5.9a) and (5.7a), we have $\left(\sum_{i=1}^m \frac{x_i}{y_i} \frac{\partial}{\partial x_i} + m - 1\right) F_m^g$ on the left hand side. But, from (5.9b) and (5.7a), we have

$$x_1 \frac{\partial}{\partial x_1} \widehat{\mathbf{H}}_2^0 = \frac{y_1^2(y_1 - 1)}{y_1 - y_2} - y_1(y_1 - 1) - (y_1 - 1) - \frac{x_1}{x_1 - x_2} = \frac{y_1^2(y_2 - 1)}{y_1 - y_2} - \frac{x_2}{x_1 - x_2},$$

and we apply this result to combine the contribution for $i_2 = 1$ in the summation on the right hand side with the $\text{sym}_{1,1,m-2}$ term on the right hand side, to give

$$\begin{aligned} & \sum_{\substack{i_2 \geq 2, i_3 \geq 0, \\ i_2 + i_3 = m-1}} \text{sym}_{1, i_2, i_3} \left(x_1 \frac{\partial}{\partial x_1} \widehat{\mathbf{H}}_{1+i_2}^0(x_1, x_{p_2}) \right) \left(x_1 \frac{\partial}{\partial x_1} F_{1+i_3}^g(x_1, x_{p_3}) \right) \\ & + \text{sym}_{1,1,m-2} \frac{y_1^2(y_2 - 1)}{y_1 - y_2} x_1 \frac{\partial}{\partial x_1} F_{m-1}^g(x_1, x_{p_3}) + \sum_{i=1}^m \left(x_i \frac{\partial}{\partial x_i} \right)^{2g+1} \widehat{\mathbf{H}}_m^0, \end{aligned}$$

on the right hand side. Now, multiply on both sides by $2^{2g-1} \frac{t^{2g}}{(2g-1)!}$, sum over $g \geq 1$, and apply \mathbb{T} , to obtain the result, from (7.2) and (5.7) b). \blacksquare

Lemma 7.5 is the final form of Theorem 3.7. In order to apply this result, we require a technical result about the invertibility of the partial differential operator Δ_k .

Proposition 7.6. *For $m \geq 1, k \geq 2$, the operator Δ_k is invertible for formal power series in y_1, \dots, y_m in which every monomial has positive exponent for each of y_1, \dots, y_m .*

Proof. Let $A := \sum_{n_1, \dots, n_m \geq 1} a_{(n_1, \dots, n_m)} y_1^{n_1} \cdots y_m^{n_m}$ and $B := \sum_{n_1, \dots, n_m \geq 1} b_{(n_1, \dots, n_m)} y_1^{n_1} \cdots y_m^{n_m}$, and suppose that $\Delta_k A = \Delta_k B$. We prove that $a_{(n_1, \dots, n_m)} = b_{(n_1, \dots, n_m)}$ for all $n_1, \dots, n_m \geq 1$ by putting a partial order \prec_ℓ on the (n_1, \dots, n_m) : We let $(n_1, \dots, n_m)_\leq$ be the vector of length m containing the n_i in weakly increasing order, and we define $(n_1, \dots, n_m)_\leq \prec_\ell (i_1, \dots, i_m)_\leq$ if and only if $(n_1, \dots, n_m)_\leq$ precedes $(i_1, \dots, i_m)_\leq$ in lexicographic order.

To prove the result for (n_1, \dots, n_m) , suppose that $n_i = \max\{n_1, \dots, n_m\}$ (make an arbitrary choice of i if there is more than one, equal, maximum element), and equate coefficients of $y_1^{n_1} \cdots y_i^{n_i+k-1} \cdots y_m^{n_m}$ in $\Delta_k A = \Delta_k B$, to obtain

$$\begin{cases} n_i a_{(n_1, \dots, n_m)} + \sum_{\substack{j \neq i, \\ n_j \geq k}} (n_j - k + 1) a_{(n_1, \dots, n_j - k + 1, \dots, n_i + k - 1, \dots, n_m)} \\ = n_i b_{(n_1, \dots, n_m)} + \sum_{\substack{j \neq i, \\ n_j \geq k}} (n_j - k + 1) b_{(n_1, \dots, n_j - k + 1, \dots, n_i + k - 1, \dots, n_m)}. \end{cases} \quad (7.8)$$

Now consider any fixed linear extension of \prec_ℓ , and note that, for any term in the summations in (7.8), we have $(n_1, \dots, n_j - k + 1, \dots, n_i + k - 1, \dots, n_m) \prec_\ell (n_1, \dots, n_m)$. The result follows from (7.8), by induction on the position in this linear extension. \blacksquare

8. Proof of Faber’s Intersection Number Conjecture for 2 and 3 Parts, with Comments about the General Case of m Parts

8.1. Faber’s Intersection Number Conjecture for two Parts

In this section, we prove Faber’s Conjecture for the case of two parts. It will be convenient to “smooth” the result to allow zeros, by applying the string equation. For cases of (2.2) when some $d_i = 0$, we can use the string equation to deduce a consistent value. For example, when $n = 2$, $d_1 = g$, $d_2 = 0$, the string equation gives $\langle \tau_g \tau_0 \rangle_g^{\text{Fab}} = \langle \tau_{g-1} \rangle_g^{\text{Fab}} = 1$, which agrees with (2.2), so Faber’s Conjecture for $n = 2$ becomes

$$\langle \tau_{d_1} \tau_{d_2} \rangle_g^{\text{Fab}} = \frac{(2g - 1)!!}{(2d_1 - 1)!!(2d_2 - 1)!!}, \quad d_1 + d_2 = g, \quad d_1, d_2 \geq 0, \quad g \geq 1. \quad (8.1)$$

For the localization side, we have the following result. In the proof, we use the notation

$$\widehat{Y}_m := Y_m(A_1), \quad m \geq 1, \quad (8.2)$$

where Y_m, A_1 are defined in (7.5), (5.6), respectively.

Theorem 8.1. *If the Faber Conjecture is true for $n = 2$, then*

$$\Psi_2(y_1, y_2, t) = \text{E sym}_{1,1} \left(\frac{y_1^3 y_2 t}{y_1 - y_2} - \frac{y_1^4 y_2^2 t}{(y_1 - y_2)(y_1^2 - y_2^2 + 4y_1^2 y_2^2 t)} \right) B_1^{-1}.$$

Proof. From Corollary 5.1 (b), we have

$$\begin{aligned} \mathbb{T}_2^g &= \sum_{\substack{a_1, a_2 \geq 0 \\ a_1 + a_2 \leq g}} (-1)^{g - a_1 - a_2} \langle \tau_{a_1} \tau_{a_2} \lambda_{g - a_1 - a_2} \rangle_g^{\text{Fab}} \xi_1^{(a_1)}(x_1) \xi_1^{(a_2)}(x_2) \\ &\quad + \sum_{k=0}^{g-1} (-1)^k \langle \tau_{g-1-k} \lambda_k \rangle_g^{\text{Fab}} \xi_2^{(g-1-k)}, \end{aligned}$$

where we have used the symmetry of the Faber symbol in the first summation. Together with Theorem 5.4 (a), (b), and (7.1), this gives

$$\begin{aligned} \Psi_2(y_1, y_2, t) &= \sum_{g \geq 1} \frac{t^{2g}}{(2g - 1)!!} [u^{2g-1}] \sum_{\substack{a_1, a_2 \geq 0, \\ a_1 + a_2 = g}} \langle \tau_{a_1} \tau_{a_2} \rangle_g^{\text{Fab}} \prod_{j=1}^2 (2a_j - 1)!! Y_j(u)^{2a_j+1} \\ &\quad - \sum_{g \geq 1} \frac{t^{2g}}{(2g - 1)!!} [u^{2g-1}] \langle \tau_{g-1} \rangle_g^{\text{Fab}} (2g - 1)!! u \text{sym}_{1,1} \frac{y_1^2 y_2}{y_1 - y_2} Y_1(u)^{2g+1}. \end{aligned}$$

Now, if the Faber Conjecture is true for $n = 2$, then from (8.1), we have

$$\begin{aligned} \Psi_2(y_1, y_2, t) &= \sum_{g \geq 1} t^{2g} [u^{2g-1}] \left(\sum_{\substack{a_1, a_2 \geq 0, \\ a_1 + a_2 = g}} Y_1(u)^{2a_1+1} Y_2(u)^{2a_2+1} \right. \\ &\quad \left. - \text{sym}_{1,1} \frac{y_1^2 y_2}{y_1 - y_2} u Y_1(u)^{2g+1} \right), \end{aligned}$$

But $\sum_{\substack{a_1, a_2 \geq 0 \\ a_1 + a_2 = g}} Y_1(u)^{2a_1+1} Y_2(u)^{2a_2+1} = \text{sym}_{1,1} \frac{Y_1(u)^{2g+3} Y_2(u)}{Y_1(u)^2 - Y_2(u)^2}$, and from Lemma 7.3 we conclude that

$$\Psi_2(y_1, y_2, t) = \frac{1}{2} \text{Esym}_{1,1} t (1 + B_1^{-1}) \left(\frac{\widehat{Y}_1^4 \widehat{Y}_2}{\widehat{Y}_1^2 - \widehat{Y}_2^2} - \frac{y_1^2 y_2}{y_1 - y_2} A_1 \widehat{Y}_1 \right).$$

From the proof of Lemma 7.3, we have $A_1 = ty_1/(1 - y_1 A_1)$, so routinely simplifying, we obtain

$$\begin{aligned} \Psi_2(y_1, y_2, t) &= \text{Esym}_{1,1} \left(\frac{y_1^3 y_2 t A_1 (1 - y_2 A_1) B_1^{-1}}{(y_1 - y_2)(y_1 + y_2 B_1)} - \frac{y_1^3 y_2 A_1^2 B_1^{-1}}{y_1 - y_2} \right) \\ &= \text{Esym}_{1,1} \left(\frac{y_1^3 y_2 t B_1^{-1}}{y_1 - y_2} + \frac{y_1^3 y_2^3 t - y_1^4 y_2^2 t B_1^{-1}}{(y_1 - y_2)(y_1^2 - y_2^2 B_1^2)} \right). \end{aligned}$$

Now it is straightforward to verify that $\text{Esym}_{1,1} \frac{y_1^3 y_2^3 t}{(y_1 - y_2)(y_1^2 - y_2^2 B_1^2)} = 0$, and the result follows with $y_1^2 - y_2^2 B_1^2 = y_1^2 - y_2^2 + 4y_1^2 y_2^2 t$. ■

For the degeneration side, we have the following result.

Theorem 8.2. $\Delta_3 \Delta_2 \Phi_2(y_1, y_2, t) = \text{Esym}_{1,1} \left(\Delta_3 \frac{y_1^4 y_2}{y_1 - y_2} t B_1^{-1} + 3y_1^5 y_2 t B_1^{-5} \right)$.

Proof. We begin by using (5.8) to obtain $\Delta_3 \mathbb{T} \widehat{\mathbf{H}}_2^0 = y_1 y_2$, so, from (5.7) b), we have $\mathbb{T} \left(x_1 \frac{\partial}{\partial x_1} \right)^{2g+1} \Delta_3 \widehat{\mathbf{H}}_2^0 = \left(y_1^3 \frac{\partial}{\partial y_1} \right)^{2g+1} y_1 y_2 = (4g + 1)!! y_1^{4g+3} y_2$, where the second equality follows by induction. This gives

$$\begin{aligned} \Delta_3 \Upsilon_2 &= \text{sym}_{1,1} \sum_{g \geq 1} 2^{2g-1} \frac{(4g + 1)!!}{(2g - 1)!} y_1^{4g+3} y_2 t^{2g} \\ &= \text{sym}_{1,1} y_1^5 y_2 t \sum_{g \geq 1} 3 \binom{-\frac{5}{2}}{2g - 1} (-4y_1^2 t)^{2g-1} \\ &= 3 \text{Esym}_{1,1} y_1^5 y_2 t B_1^{-5}. \end{aligned}$$

Now, consider Lemma 7.5 with $m = 2$, and apply Δ_3 to both sides, to obtain $\Delta_3 \Delta_2 \Phi_2 = \Delta_3 \text{sym}_{1,1} \frac{y_1^5 y_2}{y_1 - y_2} \frac{\partial}{\partial y_1} \Phi_1(y_1, t) + \Delta_3 \Upsilon_2$. The result follows from Corollary 7.4 and the expression for $\Delta_3 \Upsilon_2$ above, since Δ_3 and $\text{sym}_{1,1}$ commute. ■

Now we compare the results from the localization and degeneration sides, to obtain a proof of Faber’s Conjecture for two parts.

Theorem 8.3. *Faber’s Intersection Number Conjecture is true for $n = 2$.*

Proof. By applying $\Delta_3 \Delta_2$ to Theorem 8.1, and from Theorem 8.2, we routinely simplify to prove that $\Delta_3 \Delta_2 \Psi_2(y_1, y_2, t)$ and $\Delta_3 \Delta_2 \Phi_2(y_1, y_2, t)$ (under the condition that the Faber Conjecture is true for $n = 2$) are both equal to $\text{Esym}_{1,1} \frac{2y_1^5 y_2 t B_1^{-5}}{y_1 - y_2} (3y_1 - 2y_2 -$

$10y_1^3t + 4y_1^2y_2t + 16y_1^5t^2 - 8y_1^4y_2t^2$). Thus we have proved that if the Faber Conjecture is true for $n = 2$, then $\Delta_3\Delta_2\Psi_2(y_1, y_2, t) = \Delta_3\Delta_2\Phi_2(y_1, y_2, t)$. Then Proposition 7.6 implies that $\Psi_2(y_1, y_2, t) = \Phi_2(y_1, y_2, t)$, and the result follows from Lemma 7.2. ■

8.2. Faber's Intersection Number Conjecture for Three Parts

The proof for three parts follows the steps and strategy of the proof for two parts, with no further technical difficulties. However, the full details of some of the expressions are somewhat longer, and we omit many below.

For the localization side, we determine that, if Faber's Conjecture is true for up to 3 parts, then

$$\begin{aligned} \Psi_3(y_1, y_2, y_3, t) &= \frac{1}{2}\text{Esym}_{1,2}t \frac{d}{dt}t (1 + B_1^{-1}) \frac{\widehat{Y}_1^8\widehat{Y}_2\widehat{Y}_3}{(\widehat{Y}_1^2 - \widehat{Y}_2^2)(\widehat{Y}_1^2 - \widehat{Y}_3^2)} \\ &\quad - \frac{1}{2}\text{Esym}_{1,1,1}t^{-2} \frac{d}{dt}t^4 (1 + B_1^{-1}) \frac{y_1^2y_2}{y_1 - y_2} A_1 \frac{\widehat{Y}_1^6\widehat{Y}_3}{\widehat{Y}_1^2 - \widehat{Y}_3^2} \\ &\quad + \frac{1}{2}\text{Esym}_{1,2}t^{-2} \frac{d}{dt}t^3 (1 + B_1^{-1}) \frac{y_1^6y_2y_3}{(y_2 - y_1)(y_3 - y_1)} \widehat{Y}_1^4 \\ &\quad + \frac{1}{2}\text{Esym}_{1,1,1}t (1 + B_1^{-1}) \alpha + \frac{1}{2}\text{Esym}_{1,2}t (1 + B_1^{-1}) \beta, \end{aligned} \quad (8.3)$$

where

$$\begin{aligned} \alpha &= -\frac{y_3^2y_2}{y_3 - y_2} A_1 \left(\frac{\widehat{Y}_1^4\widehat{Y}_3^3}{\widehat{Y}_1^2 - \widehat{Y}_3^2} + 2 \frac{\widehat{Y}_1^4\widehat{Y}_3^5}{(\widehat{Y}_1^2 - \widehat{Y}_3^2)^2} \right) + 2 \frac{y_1^2y_2}{y_1 - y_2} A_1 \frac{\widehat{Y}_1^8\widehat{Y}_3}{(\widehat{Y}_1^2 - \widehat{Y}_3^2)^2} \\ &\quad + \frac{y_1^3y_2^4y_3}{(y_2 - y_3)(y_1 - y_2)^2} A_1^2 \widehat{Y}_1^2 \widehat{Y}_2 + \frac{y_1^6y_2y_3}{(y_2 - y_1)^2(y_3 - y_1)} A_1^2 \widehat{Y}_1^3, \\ \beta &= (1 - y_1) \widehat{Y}_1^4 \widehat{Y}_2 \widehat{Y}_3 + 3 \frac{y_1^5y_2y_3}{(y_2 - y_1)(y_3 - y_1)} A_1^2 \widehat{Y}_1^3. \end{aligned}$$

This is a large expression for Ψ_3 , but the method of proof follows exactly that of Theorem 8.1, and is routine: We apply Corollary 5.1 (b) with $m = 3$, and parts (a)–(c) of Theorem 5.4, to obtain an expression for $\Lambda \mathbb{T}_3$, and use a smoothing of Faber's Conjecture for three parts to account for parts that are zero. The above expression then follows from (7.1).

For the degeneration side, we determine that

$$\begin{aligned} \Delta_2\Phi_3(y_1, y_2, y_3, t) &= \text{Esym}_{1,2}y_1^5y_2y_3tB_1^{-1} + 3\text{Esym}_{1,2}y_1^5y_2y_3tB_1^{-5} \\ &\quad + \frac{1}{2}\text{Esym}_{1,1,1} \left(\frac{y_1^5y_3}{y_1 - y_3} \frac{\partial}{\partial y_1} + \frac{y_2^5y_3}{y_2 - y_3} \frac{\partial}{\partial y_2} \right) t \\ &\quad (1 + B_1^{-1}) \left(\frac{\widehat{Y}_1^4\widehat{Y}_2}{\widehat{Y}_1^2 - \widehat{Y}_2^2} - \frac{y_1^2y_2}{y_1 - y_2} A_1 \widehat{Y}_1^2 \right). \end{aligned} \quad (8.4)$$

The method of proof of this result follows exactly that of Theorem 8.2: We apply Lemma 7.5 with $m = 3$, using the expression in Theorem 8.1 for $\Phi_2 = \Psi_2$ and the fact that $\widehat{\mathbf{TH}}_3 = y_1 y_2 y_3$, from (5.8), which implies that $\Upsilon_3(y_1, y_2, y_3, t) = 3\text{Esym}_{1,2} y_1^5 y_2 y_3 t B_1^{-5}$, giving the result.

Now we compare the results from the localization and degeneration sides, to obtain a proof of Faber’s Conjecture for three parts.

Theorem 8.4. *Faber’s Intersection Number Conjecture is true for $n = 3$.*

Proof. Applying Δ_2 to (8.3), and from (8.4), with routine reductions (using Maple), we prove that $\Delta_2 \Psi_3(y_1, y_2, y_3, t) = \Delta_2 \Phi_3(y_1, y_2, y_3, t)$, under the condition that the Faber Conjecture is true for $n \leq 3$. Then Proposition 7.6 implies that $\Psi_3(y_1, y_2, y_3, t) = \Phi_3(y_1, y_2, y_3, t)$, and the result follows from Lemma 7.2 and Theorem 8.3. ■

8.3. Intersection Numbers with Four or More Parts

For the case of n parts, on the localization side, we need to determine $\Lambda \xi_n^{(i)}$, for which we need an expression for the double Hurwitz series \mathbf{H}_n^0 . Explicit expressions for the latter are known for $n = 4$ [21, Cor. 5.7] and $n = 5$ [21, Cor. 5.8], and with the help of Maple, we could in principle thus obtain expressions for Ψ_4 and Ψ_5 analogous to (8.3).

For the degeneration side, we apply Lemma 7.5, and note that Φ_n is obtained recursively from Φ_j , $j < n$, once we have $\widehat{\mathbf{H}}_n^0$, which is given explicitly for all n in (5.8). Thus we can obtain a rational expression for Φ_n for any n , by using Maple to help with the size of the expressions.

Together, via Lemma 7.2, these would enable us to prove Faber’s Conjecture for four and five parts. However, we are prevented from proving the result for larger values of n because we do not have explicit expressions for the double Hurwitz series \mathbf{H}_n^0 , and we do not see how to obtain an explicit formula that holds for all n . In order to use our methodology to prove the result for arbitrary n , we would need to use our functional equations for the various tree series given in Theorem 4.4, and then use the Join-cut Equation for the double Hurwitz series.

Appendix

9. Proofs of Theorems 5.3 and 5.4

In this Appendix we develop the substantial amount of material that is required to determine $\Lambda \xi_m^{(i)}$ for $m = 1, 2, 3$, and is not required elsewhere in the paper. A key ingredient is the symmetrization of the double Hurwitz generating series.

9.1. Preliminaries

(a) Completing the symmetrization on the localization side: In order to determine $\Lambda \xi_m^{(i)}$ we have the following result, which together with Corollary 5.1 completes the

symmetrization of Theorem 4.4 (the third form of the Localization Tree Theorem). It requires the symmetrized series

$$\begin{cases} f_{j,m}(x_1, \dots, x_m, u) & := \Xi_m f_j(z, u; \mathbf{p}), \\ g_{j,m}(x_1, \dots, x_m, u) & := \Xi_m g_j(z, u; \mathbf{p}), \\ \mathbf{H}_m^0(x_1, \dots, x_m, u; \mathbf{q}) & := \Xi_m H^0(z, u; \mathbf{p}, \mathbf{q}), \end{cases} \quad (9.1)$$

for $m \geq 1$, and the substitution operator

$$\Omega: q_i \mapsto \frac{i^{i-1}}{i!}, \quad i \geq 1. \quad (9.2)$$

Corollary 9.1. For $m \geq 1$,

$$a) \xi_m^{(i)} = \sum_{j \geq 1} \frac{j^{j+i}}{j!} f_{j,m}, \quad i \geq 0,$$

where, for $j \geq 1$,

$$b) f_{j,m} = \sum_{\substack{k \geq 0, \\ j_1, \dots, j_k \geq 1, \\ i_0, \dots, i_k \geq 1, \\ i_0 + \dots + i_k = m}} \text{sym}_{i_0, \dots, i_k} \frac{u^{-2}}{k!} \left(\Omega j \frac{\partial^{k+1}}{\partial q_j \partial q_{j_1} \dots \partial q_{j_k}} \mathbf{H}_{i_0}^0(x_{p_0}, u; \mathbf{q}) \right) \prod_{a=1}^k g_{j_a, i_a}(x_{p_a}, u),$$

$$c) g_{j,m} = \sum_{\substack{k \geq 0, j_1, \dots, j_k \geq 1, \\ i_1, \dots, i_k \geq 1, \\ i_1 + \dots + i_k = m}} \text{sym}_{i_1, \dots, i_k} \frac{(j + j_1 + \dots + j_k)^{k-2}}{k!} \frac{j^{j+1}}{j!} \prod_{a=1}^k \frac{j_a^{j_a}}{j_a!} f_{j_a, i_a}(x_{p_a}, u).$$

Proof. The result follows by applying Ξ_m to (4.2), Theorem 4.4 (c) and (d), for $m \geq 1$, and using the properties of Ξ_m given in [19, Lemmas 4.1–4.3]. ■

We shall determine $\Lambda \xi_m^{(i)}$, $m = 1, 2, 3$, by applying Corollary 9.1 directly. The results that are needed for the series \mathbf{H}_m^0 and its partial derivatives are given in the next two sections.

(b) The symmetrized double Hurwitz series, genus 0: Now let $Q(t) := \sum_{j \geq 1} q_j t^j$, and v be the unique formal power series solution of the functional equation

$$v = x e^{uQ(v)}. \quad (9.3)$$

Let

$$v_i := v(x_i), \quad Q_i := Q(v_i), \quad \mu(t) := (1 - utQ'(t))^{-1}, \quad \mu_i := \mu(v_i). \quad (9.4)$$

Then in [21], we proved that

$$x_1 \frac{\partial}{\partial x_1} \mathbf{H}_1^0 = uQ_1, \quad \mathbf{H}_2^0 = \log \left(\frac{v_1 - v_2}{x_1 - x_2} \right) - uQ_1 - uQ_2, \quad \mathbf{H}_3^0 = \sum_{i=1}^3 (\mu_i - 1) \prod_{\substack{1 \leq j \leq 3, \\ j \neq i}} \frac{v_j}{v_i - v_j}. \quad (9.5)$$

(c) Partial derivatives of the symmetrized double Hurwitz series: Differentiating (9.3) gives

$$x_1 \frac{\partial}{\partial x_1} v_1 = v_1 \mu_1, \quad \frac{\partial}{\partial q_j} v_1 = uv_1^{j+1} \mu_1, \tag{9.6}$$

and from (9.5) and (9.6) we obtain

$$\frac{\partial}{\partial q_j} \mathbf{H}_2^0 = \frac{uv_1^{j+1} \mu_1 - uv_2^{j+1} \mu_2}{v_1 - v_2} - uv_1^j \mu_1 - uv_2^j \mu_2 = \text{sym}_{1,1} \frac{uv_1^j \mu_1 v_2}{v_1 - v_2}. \tag{9.7}$$

For partial derivatives of \mathbf{H}_1^0 in general, we have the following result.

Lemma 9.2. (a) For $j \geq 1$, $\frac{\partial}{\partial q_j} \mathbf{H}_1^0 = \frac{u}{j} v_1^j$.

(b) For $k \geq 1$, $j_1, \dots, j_k \geq 1$, $\frac{\partial^k}{\partial q_{j_k} \dots \partial q_{j_1}} \mathbf{H}_1^0 = u^k \left(\mu_1 v_1 \frac{\partial}{\partial v_1} \right)^{k-2} \left(\mu_1 v_1^{j_1 + \dots + j_k} \right)$.

Proof. Differentiating (9.5), we obtain

$$\frac{\partial}{\partial q_j} x_1 \frac{\partial}{\partial x_1} \mathbf{H}_1^0 = uQ'(v_1) \frac{\partial v_1}{\partial q_j} + uv_1^j = uQ'(v_1) uv_1^{j+1} \mu_1 + uv_1^j = uv_1^j \mu_1,$$

where we have used (9.6) for the second equality. Dividing by x_1 and integrating, we obtain $\frac{\partial}{\partial q_j} \mathbf{H}_1^0 = \int_0^{x_1} uv_1^j \mu_1 \frac{dx_1}{x_1} = \int_0^{v_1} uv_1^{j-1} dv_1$, where we have used (9.6) to change variables for the second equality. Part (a) follows immediately.

We prove part (b) by induction on k . For the base case, note that the result is true for $k = 1$, since this is equivalent to part (a) above. Now assume that the result holds for $k - 1 \geq 1$, and note that (9.6) implies the operator identity $\mu_1 v_1 \partial / \partial v_1 = x_1 \partial / \partial x_1$, so we have

$$\begin{aligned} \frac{\partial^k}{\partial q_{j_k} \dots \partial q_{j_1}} \mathbf{H}_1^0 &= \frac{\partial}{\partial q_{j_k}} u^{k-1} \left(x_1 \frac{\partial}{\partial x_1} \right)^{k-2} \left(\mu_1 v_1 \frac{\partial}{\partial v_1} \right)^{-1} \left(\mu_1 v_1^{j_1 + \dots + j_{k-1}} \right) \\ &= u^{k-1} \left(x_1 \frac{\partial}{\partial x_1} \right)^{k-2} \frac{\partial}{\partial q_{j_k}} \left(\frac{v_1^{j_1 + \dots + j_{k-1}}}{j_1 + \dots + j_{k-1}} \right), \end{aligned}$$

and part (b) follows from (9.6) and the operator identity. ■

(d) Polynomiality and the symmetrized double Hurwitz series: From Corollary 9.1 (a),

(b), to evaluate $\Lambda \xi_m^{(i)}$ we need to apply the substitution $\Lambda \Omega$ to partial derivatives of the symmetrized double Hurwitz series \mathbf{H}_m^0 , which we have shown above can be expressed in terms of v_i and μ_i . Thus define

$$V_i := \Lambda \Omega v_i. \tag{9.8}$$

In the following result, the action of the change of variables C , the second step in our fundamental transformation, is given for expressions in V_i . Note how indirectly we proceed in the proof, especially for part (a), as referred to in the discussion following (5.5). This result is the key to polynomiality on the localization side.

Lemma 9.3. For $i \geq 1$,

- a) $C y(V_i) = \frac{(1-u)y_i}{1-uy_i}$,
- b) $C \Lambda \Omega \mu_i = \frac{1-uy_i}{1-u}$,
- c) $C V_i \frac{\partial}{\partial V_i} = y_i^2(y_i-1) \frac{1-u}{1-uy_i} \frac{\partial}{\partial y_i} C$.

Proof. For part (a), the series $w = w(x)$ defined in (5.5) a) is the unique formal power series solution to $w = xe^w$, which we shall call the *tree equation* (see, e.g., [17, Ex. 1.2.5]). Now $V_i = V(x_i)$, where, from (9.3), $V = V(x)$ is the unique formal power series solution to

$$V = (1-u)^{-1}x \exp\left(-u \sum_{j \geq 1} \frac{j^{j-1}}{j!} V^j\right) = (1-u)^{-1}xe^{-uw(V)}, \tag{9.9}$$

where the second equality follows from (5.5a). But the tree equation gives $w(V) = Ve^{w(V)}$, and eliminating V between this equation and (9.9), we obtain $(1-u)w(V) = xe^{(1-u)w(V)}$. Comparing this to the tree equation, we conclude that $(1-u)w(V(x)) = w(x)$, and by (5.5) b), this implies

$$y(V_i) = \frac{(1-u)y_i}{1-uy_i}, \tag{9.10}$$

where here $y_i = y(x_i)$ is a series in x_i . Part (a) follows immediately.

For part (b), from (9.4) and (5.5) b) we have

$$\Lambda \Omega \mu_i = \frac{1}{1+u \sum_{j \geq 1} \frac{j^j}{j!} V_i^j} = \frac{1}{1+u(y(V_i)-1)},$$

and part (b) follows immediately from part (a).

For part (c), from (5.7) a), we have

$$V_i \frac{\partial}{\partial V_i} = y(V_i)^2(y(V_i)-1) \frac{\partial}{\partial y(V_i)}. \tag{9.11}$$

But, by differentiating (9.10), we obtain

$$\frac{\partial y(V_i)}{\partial y_i} = \frac{1-u}{(1-uy_i)^2} = \frac{y(V_i)^2}{(1-u)y_i^2},$$

which implies $C y(V_i)^2 \frac{\partial}{\partial y(V_i)} = (1-u)y_i^2 \frac{\partial}{\partial y_i} C$, and part (c) follows immediately from (9.11). ■

(e) Two results: In the proofs of Section 9.2, we make use of the identity

$$\sum_{j,k \geq 1} \frac{1}{j+k} \frac{j^{j+1}}{j!} V_1^j \frac{k^k}{k!} V_2^k = \frac{V_2}{V_2-V_1} - y(V_1)^2 \frac{y(V_2)-1}{y(V_2)-y(V_1)}, \tag{9.12}$$

whose proof follows from [18], and the fact (an easy induction) that

$$\left(\frac{y_i^3}{1-uy_i} \frac{\partial}{\partial y_i}\right)^i \frac{y_i}{1-uy_i} = (2i-1)!! \left(\frac{y_i}{1-uy_i}\right)^{2i+1}, \quad i \geq 1. \tag{9.13}$$

9.2. The Proofs

Proofs of Theorem 5.3 (a) ($m = 1$), and Theorem 5.4 (a): From Corollary 9.1 (b) with $m = 1$, Lemma 9.2 (a) and (9.8), we have

$$\Lambda f_{j,1} = u^{-2} \Lambda \Omega j \frac{\partial}{\partial q_j} \mathbf{H}_1^0 = -u^{-1} \Lambda \Omega v_1^j = -u^{-1} V_1^j. \tag{9.14}$$

Then from Corollary 9.1 (a) with $m = 1$, we obtain

$$\Lambda \xi_1^{(i)} = \sum_{j \geq 1} \frac{j^{j+i}}{j!} \Lambda f_{j,1} = -u^{-1} \sum_{j \geq 1} \frac{j^{j+i}}{j!} V_1^j = -u^{-1} \left(V_1 \frac{\partial}{\partial V_1}\right)^i (y(V_1) - 1),$$

from (5.5) b). But Lemma 9.3 (a) and (c) give

$$C \Lambda \xi_1^{(i)} = -u^{-1} (1-u)^i \left(y_1^2 \frac{y_1-1}{1-uy_1} \frac{\partial}{\partial y_1}\right)^i \left(\frac{(1-u)y_1}{1-uy_1} - 1\right),$$

and Theorem 5.3 (a) ($m = 1$) follows immediately. Then $T \Lambda \xi_1^{(i)} = T u^{-1} (1-u)^{i+1} \left(\frac{y_1^3}{1-uy_1} \frac{\partial}{\partial y_1}\right)^i \frac{y_1}{1-uy_1}$, and Theorem 5.4 (a) follows immediately from (9.13). ■

Proofs of Theorem 5.3 (a) ($m = 2$), and Theorem 5.4 (b): From Corollary 9.1 (c) with $m = 1$, and (9.14), we obtain

$$\begin{aligned} \Lambda g_{j_1,1}(x_1, u) &= \sum_{j_2 \geq 1} \frac{1}{j_1 + j_2} \frac{j_1^{j_1+1}}{j_1!} \frac{j_2^{j_2}}{j_2!} \Lambda f_{j_2,1}(x_1, u) \\ &= -u^{-1} \frac{j_1^{j_1+1}}{j_1!} \sum_{j_2 \geq 1} \frac{1}{j_1 + j_2} \frac{j_2^{j_2}}{j_2!} V_1^{j_2}, \end{aligned} \tag{9.15}$$

and from Corollary 9.1 (b) with $m = 2$, we obtain $\Lambda f_{j,2} = S_1 + S_2$, where

$$S_1 = u^{-2} \Lambda \Omega j \frac{\partial}{\partial q_j} \mathbf{H}_2^0, \quad S_2 = u^{-2} \text{sym}_{1,1} \sum_{j_1 \geq 1} \Lambda g_{j_1,1}(x_2, u) \Omega j \frac{\partial^2}{\partial q_j \partial q_{j_1}} \mathbf{H}_1^0(x_1, u).$$

We can determine S_1 immediately from (9.7), giving $S_1 = -u^{-1} j \text{sym}_{1,1} \frac{V_1^{j_1} V_2}{V_1 - V_2} \Lambda \Omega \mu_1$. To determine S_2 , using Lemma 9.2 with $k = 2$ and (9.15), we can obtain $S_2 = -u^{-1} j \text{sym}_{1,1} V_1^j (\Lambda \Omega \mu_1) \sum_{j_1, j_2 \geq 1} \frac{1}{j_1 + j_2} \frac{j_1^{j_1+1}}{j_1!} V_1^{j_1} \frac{j_2^{j_2}}{j_2!} V_2^{j_2}$. Adding S_1 and S_2 , and using (9.12),

we get $f_{j,2} = u^{-1} j \operatorname{sym}_{1,1} V_1^j y(V_1)^2 \frac{y(V_2)-1}{y(V_2)-y(V_1)} \Lambda \Omega \mu_1$. Then from Corollary 9.1 (a) with $m = 2$, we obtain

$$\begin{aligned} \Lambda \xi_2^{(i)} &= \sum_{j \geq 1} \frac{j^{j+i}}{j!} \Lambda f_{j,2} = u^{-1} \operatorname{sym}_{1,1} y(V_1)^2 \frac{y(V_2)-1}{y(V_2)-y(V_1)} (\Lambda \Omega \mu_1) \sum_{j \geq 1} \frac{j^{j+i+1}}{j!} V_1^j \\ &= u^{-1} \operatorname{sym}_{1,1} y(V_1)^2 \frac{y(V_2)-1}{y(V_2)-y(V_1)} (\Lambda \Omega \mu_1) \left(V_1 \frac{\partial}{\partial V_1} \right)^{i+1} (y(V_1) - 1), \end{aligned}$$

from (5.5) b). Now Lemma 9.3 (a), (b), (c) give

$$C \Lambda \xi_2^{(i)} = u^{-1} (1-u)^{i+1} \operatorname{sym}_{1,1} \frac{y_1^2 (y_2 - 1)}{y_2 - y_1} \left(y_1^2 \frac{y_1 - 1}{1 - uy_1} \frac{\partial}{\partial y_1} \right)^{i+1} \left(\frac{(1-u)y_1}{1 - uy_1} - 1 \right),$$

and Theorem 5.3 (a) ($m = 2$) follows immediately. Then we have

$$T \Lambda \xi_2^{(i)} = T u^{-1} (1-u)^{i+2} \operatorname{sym}_{1,1} \frac{y_1^2 y_2}{y_2 - y_1} \left(\frac{y_1^3}{1 - uy_1} \frac{\partial}{\partial y_1} \right)^{i+1} \frac{y_1}{1 - uy_1},$$

and Theorem 5.4 (b) follows immediately from (9.13). ■

Outline of proofs of Theorem 5.3 (a) ($m = 3$), and Theorem 5.4 (c): The method of proof of these results follows those for the cases $m = 1, 2$ above, with no further technical difficulties. We give only an outline. We first determine $\Lambda f_{j,3}$ from Corollary 9.1 (b) This requires $\Lambda g_{j,2}$, which we determine from Corollary 9.1 (c), and the partial derivatives $\frac{\partial^3}{\partial q_j \partial q_{j_1} \partial q_{j_2}} \mathbf{H}_1^0$, $\frac{\partial^2}{\partial q_j \partial q_{j_1}} \mathbf{H}_2^0$ and $\frac{\partial}{\partial q_j} \mathbf{H}_3^0$. But rational expressions for these partials are routinely obtained, from Lemma 9.2 (b) for \mathbf{H}_1^0 , and by applying the chain rule and (9.6) to (9.7) and (9.5). We then use Corollary 9.1 (a) with $m = 3$ to obtain an explicit expression for $\Lambda \xi_3^{(i)}$, and then apply C to prove Theorem 5.3 (a) ($m = 3$). Then Theorem 5.4 (c) follows immediately. ■

10. Glossary of Notation

If a symbol in column 1 has an argument, it appears as the first item of the entry in column 3. In column 2, “C” denotes “Corollary”, “Cn” denotes “Conjecture”, “D” denotes “Definition”, “P” denotes “Proposition”, “T” denotes “Theorem”, “b” denotes “immediately before”, “f” denotes “immediately following” and “I” denotes “Introduction to”.

\ddagger	f(3.18)	Removal of contribution of root from corresponding product
\bullet	D3.10	Root-vertex of tree t
\star	D4.2	Combinatorial product
$[x^k]f$	-	Coefficient of x^k in a formal power series f
$[\cdot, \cdot]_{\eta}$	D4.1	\mathcal{A} , wt : Exponential generating series
$ \alpha $	I§3	Sum of parts of α
$\langle \dots \rangle_g^{\text{Fab}}$	(2.3)	$\tau_{a_1} \cdots \tau_{a_n} \lambda_k$: Faber symbol for π_* ($\Psi_1^{a_1} \cdots \Psi_n^{a_n} \lambda_k$)
$\langle \dots \rangle_g^{\text{Fab}}$	(7.3)	$\tau_{a_1} \cdots \tau_{a_n} \lambda_k$: Faber number
α	I§3	Ptn. of d : branching over ∞
α	(3.15)	$\coprod_{v \in \mathcal{V}_{\infty}} \gamma^v$
$\alpha(t)$	bD4.2	Relabelled localization tree
β	§3.1.2	Ptn. of d : branching over 0
β^v	D3.10	List of weights on 0∞ -edges incident with ∞ -vertex v
γ^v	D3.10	List of weights on all ∞t -edges incident with ∞ -vertex v
Δ_k	(7.6)	Partial differential operator
δ^v	D3.10	List of weights on 0∞ -edges incident with 0-vertex v
$\varepsilon(e)$	D3.10	Weight on a 0∞ -edge of t
$\eta_0(t)$	D3.10	No. of non-root 0-vertices in t
$\eta_k(t)$	bD4.1	No. of t -vertices in t incident with edge of weight k
Υ_m	(7.7)	(y_1, \dots, y_m)
κ_i	§2.2	“ κ -class”
Λ	(5.4)	Substitution operator $u \mapsto -u, x_i \mapsto (1-u)^{-1}x_i$
λ_i	§2.2	λ -class
μ, μ_i	(9.4)	Used for double Hurwitz series
Ξ_m	(5.1)	Symmetrization operator
$\xi^{(i)}$	(4.2)	$(z, u; \mathbf{p})$: Localization tree series
$\xi_m^{(i)}$	(5.3)	(x_1, \dots, x_m) : Symmetrized localization tree series
Φ_m	(7.2)	(y_1, \dots, y_m, t) : Generating series for $\text{T} F_m^g(x_1, \dots, x_m)$
ρ_j	(5.2)	Used in Ξ_m
Ψ_m	(7.1)	(y_1, \dots, y_m, t) : Generating series for $\text{T} \Lambda \mathbb{T}_m^g / \mathbb{G}_{g,1}$
ψ_i	§2.1	First Chern class
Ω	(9.2)	Substitution operator $q_i \mapsto \frac{i^{i-1}}{i!}$
$\Omega_{\infty}^{\text{br}}, \Omega_0^{\text{br}}$	§4.2	Variants of branch decompositions for trees
A_i	(7.5)	-
$A(t)$	(3.18)	Weight function for t
$B(t)$	(3.17)	Weight function for t
B_i	(7.5)	-
br	(3.2)	Fantechi-Pandharipande branch morphism
$C(t)$	(3.17)	Weight function for t
C	f(5.5)	Change of variables $x_i \rightsquigarrow y_i$
$D(t)$	(3.17)	Weight function for t
d	§3	Degree of a cover
E	b(7.5)	Even subseries operator

$\mathcal{E}_{0\infty}(\mathfrak{t})$	D3.10	Set of edges 0∞ -edges of \mathfrak{t}
$\mathcal{E}_{\infty\mathfrak{t}}(\mathfrak{t})$	D3.10	Set of edges $\infty\mathfrak{t}$ -edges of \mathfrak{t}
$\mathbb{F}^{g,\alpha}$	(3.5)	Faber-Hurwitz class
F_α^g	(3.10)	Faber-Hurwitz number, $\in \mathbb{Q}$; note $\mathbb{F}^{g,\alpha} = F_\alpha^g \mathbb{G}_{g,1}$
F_α^g	(3.12)	$(z; \mathbf{p})$: Generating series for Faber numbers F_α^g
F_m^g	(5.3)	(x_1, \dots, x_m) Symmetrized $F^g(z; \mathbf{p})$
f_j	T4.4	Generating series for $\mathcal{T}_{0,j}$
$f_{j,m}$	(9.1)	(x_1, \dots, x_m, u) : Symmetrized series for $\mathcal{T}_{0,j}$
$\mathbb{G}_{g,1}$	T3.5	A “natural generator” of $R_{2g-1}(\mathcal{M}_{g,1})$, C6.3
$\mathbb{G}_{g,d}$	T3.5	-
$\mathbb{G}_{g,d,\sim}$	T3.5	-
g	§2.1	Genus (of a curve)
g_j	T4.4	Generating series for $\mathcal{T}_{\infty,j}$
$g_{j,m}$	(9.1)	(x_1, \dots, x_m, u) symmetrized series for $\mathcal{T}_{\infty,j}$
H_α^0	(3.8)	Genus 0 (single) Hurwitz no.
$H_{\alpha,\beta}^0$	(3.23)	Genus 0 double Hurwitz no.
$\mathbb{H}_j^{g,\alpha}$	§3	Hurwitz class
\widehat{H}^0	(3.11)	$(z; \mathbf{p})$: Gen. series for genus 0 single Hurwitz nos. H_α^0
H^0	(4.1)	$(z, u; \mathbf{p}; \mathbf{q})$: Gen. series for genus 0 double Hurwitz nos. $H_{\alpha,\beta}^0$
$\widehat{\mathbf{H}}_m^0$	(5.3)	(x_1, \dots, x_m) : Symmetrized single Hurwitz series
\mathbf{H}_m^0	(9.1)	$(x_1, \dots, x_m, u; \mathbf{q})$: Symmetrized double Hurwitz series, genus 0
L	P3.1	Class in $\text{Sym}_{\alpha,\beta}^{r^g}(\mathbb{P}^1)$
$l(\alpha)$	I§3	Number of parts of α
$\overline{\mathcal{M}}_g$	§2.1	Moduli space of smooth curves
$\overline{\mathcal{M}}_{g,n}$	§2.1	Moduli space of stable b -pointed genus g curves
$\overline{\mathcal{M}}_{g,\alpha}(\mathbb{P}^1)$	§3	Moduli space of genus g relative stable maps to \mathbb{P}^1 relative to one pt. ∞
$\overline{\mathcal{M}}_{g,\alpha,\beta}(\mathbb{P}^1)$	§3.1.2	Moduli space of relative stable maps to \mathbb{P}^1 relative to two points 0 and ∞
$\overline{\mathcal{M}}_{g,\alpha,\beta}(\mathbb{P}^1)\sim$	§3.1.2	Moduli space of “rubber maps”
$\mathcal{M}_{g,\alpha}(\mathbb{P}^1)^{rt}$	§3.1.3	Moduli space of relative stable maps with rational tails
$\mathcal{M}_{g,\alpha,\beta}(\mathbb{P}^1)^{rt}$	§3.1.3	Similar to the above
$\mathcal{M}_{g,\alpha}(\mathbb{P}^1)\bullet$	§3.1.4	Possibly disconnected moduli space
$\mathbb{P}_m^g(\dots)$	(3.16)	$(\alpha_1, \dots, \alpha_m)$: Faber polynomial
\mathcal{P}	I§3	Set of all non-empty partitions
\mathbf{p}	(4.1)	(p_1, p_2, \dots)
p_i	f(3.11)	Indeterminate marking a part i in α
$Q(t)$	(9.4)	Generating series for indeterminates q_i
\mathbf{q}	(4.1)	(q_1, q_2, \dots)
q_i	f(4.1)	Indeterminate marking a part i in β

$R^*(\mathcal{M}_{g,n}^{rt})$	I§1	Tautological ring of $\mathcal{M}_{g,n}^{rt}$
r_α^g	(3.1)	“Expected” no. of branch pts. away from ∞
$r_{\alpha,\beta}^g$	(3.3)	“Expected” no. of branch pts. away from 0 and ∞
$r_{g,\alpha}^{\text{Fab}}$	(3.6)	Number of fixed branch pts. on \mathbb{P}^1
r_∞	(3.14)	-
$R_j(\mathcal{M})$	S2.2	$R^{\dim \mathcal{M}-j}(\mathcal{M})$ for open subset \mathcal{M} of $\overline{\mathcal{M}}_{g,n}$
S_i	fT5.3	Operator for terms of total degree i
$\text{sym}_{i_1, \dots, i_k}$	f(5.2)	Summation operator for symmetrization
$\mathcal{T}_{g,m}^j, \mathcal{T}_{0,j}$	D3.10	Sets of localization trees
$\mathcal{T}_{\infty,j}$		
T, T'	§5.3	Operators for maximum degree
$T \Xi_m$	§5	Fundamental transformation
\mathbb{T}^g	T4.4(b)	$(z, u; \mathbf{p})$: Tree sum related to generating series for $\mathbb{F}_{g,\alpha}^g$
\mathbb{T}_m^g	(5.3)	(x_1, \dots, x_m, u) : Symmetrized $\mathbb{T}^g(z, u; \mathbf{p})$
\mathfrak{t}	D3.10	Localization tree
$\mathcal{V}_0(\mathfrak{t})$	D3.10	Set of 0-vertices in \mathfrak{t}
$\mathcal{V}_\infty(\mathfrak{t})$	D3.10	Set of ∞ -vertices in \mathfrak{t}
$\mathcal{V}_t(\mathfrak{t})$	D3.10	Set of t -vertices in \mathfrak{t}
$\mathcal{U}_k(\mathcal{S})$	fL4.3	Unordered list
v, v_i	(9.3),(9.4)	Implicitly defined indets. for double Hurwitz series
v_i	§4.2	0-vertex labelled i
V_i	(9.8)	Transform of v_i
w, w_i	(5.5a)	Rooted tree series
\mathfrak{w}_γ	§4.2	Star of t -vertices rooted at an ∞ -vertex
x_i	(5.1)	Indeterminates for symmetrized series
\hat{Y}_m	(8.2)	-
$Y_i(u)$	(5.6)	-
y, y_i	(5.5b)	Implicitly defined indets. for symmetrized series

Acknowledgments. The third author has benefited from discussions with Renzo Cavalieri, Carel Faber, Y.-P. Lee, Rahul Pandharipande, Hsian-Hua Tseng, and especially Tom Graber, who developed many of the algebro-geometric foundations on which this paper is based. We also thank Ezra Getzler, Kefeng Liu, and Melissa Liu.

References

1. Bertram, A., Cavalieri, R., Todorov, G.: Evaluating tautological classes using only Hurwitz numbers. *Trans. Amer. Math. Soc.* 360(11), 6103–6111 (2008)
2. Bryan, J., Pandharipande, R.: The local Gromov-Witten theory of curves. *J. Amer. Math. Soc.* 21(1), 101–136 (2008)
3. Cavalieri, R.: Generating functions for Hurwitz-Hodge integrals. *Adv. Math.* 218(5), 1419–1429 (2008)

4. Deninger, C., Murre, J.: Motivic decomposition of abelian schemes and the Fourier transform. *J. Reine Angew. Math.* 422, 201–219 (1991)
5. Ekedahl, T., Lando, S., Shapiro, M., Vainshtein, A.: On Hurwitz numbers and Hodge integrals. *C. R. Acad. Sci. Paris Sér. I Math.* 328, 1175–1180 (1999)
6. Ekedahl, T., Lando, S., Shapiro, M., Vainshtein, A.: Hurwitz numbers and intersections on moduli spaces of curves. *Invent. Math.* 146(2), 297–327 (2001)
7. Faber, C.: A conjectural description of the tautological ring of the moduli space of curves. In: Faber, C., Looijenga, E. (eds.) *Moduli of Curves and Abelian Varieties*, pp. 109–129. Vieweg, Braunschweig (1999)
8. Faber, C.: Algorithms for computing intersection numbers on moduli spaces of curves, with an application to the class of the locus of Jacobians. In: Hulek, K., Catanese, F., Peters, C., Reid, M. (eds.) *New Trends in Algebraic Geometry*, pp. 93–109. London Math. Soc. Lecture Note Ser. 264. Cambridge Univ. Press, Cambridge (1999)
9. Faber, C.: MAPLE program for computing Hodge integrals. Personal communication. Available at <http://math.stanford.edu/~vakil/programs/>.
10. Faber, C.: Personal communication. (2006)
11. Faber, C., Pandharipande, R.: Relative maps and tautological classes. *J. Eur. Math. Soc.* 7(1) 13–49 (2005)
12. Fantechi, B., Pandharipande, R.: Stable maps and branch divisors. *Compositio Math.* 130(3), 345–364 (2002)
13. Gathmann, A.: Absolute and relative Gromov-Witten invariants of very ample hypersurfaces. *Duke Math. J.* 115(2), 171–203 (2002)
14. Getzler, E., Pandharipande, R.: Virasoro constraints and the Chern classes of the Hodge bundle. *Nuclear Phys. B* 530(3), 701–714 (1998)
15. Givental, A.: Semisimple Frobenius structures at higher genus. *Internat. Math. Res. Notices* 2001(23), 1265–1286 (2001)
16. Givental, A.: Gromov-Witten invariants and quantization of quadratic hamiltonians. *Mosc. Math. J.* 1(4), 551–568, 645 (2001)
17. Goulden, I.P., Jackson, D.M.: *Combinatorial Enumeration*. Wiley, New York (1983)
18. Goulden, I.P., Jackson, D.M.: Transitive factorisations into transpositions and holomorphic mappings on the sphere. *Proc. Amer. Math. Soc.* 125(1), 51–60 (1997)
19. Goulden, I.P., Jackson, D.M., Vainshtein, A.: The number of ramified coverings of the sphere by the torus and surfaces of higher genera. *Ann. Combin.* 4(1), 27–46 (2000)
20. Goulden, I.P., Jackson, D.M., Vakil, R.: The Gromov-Witten potential of a point, Hurwitz numbers, and Hodge integrals. *Proc. London Math. Soc.* (3) 83(3), 563–581 (2001)
21. Goulden, I.P., Jackson, D.M., Vakil, R.: Towards the geometry of double Hurwitz numbers. *Adv. Math.* 198(1), 43–92 (2005)
22. Goulden, I.P., Jackson, D.M., Vakil, R.: A short proof of the λ_g -Conjecture without Gromov-Witten theory: Hurwitz theory and the moduli of curves. *J. Reine Angew. Math.* 637, 175–191 (2009)
23. Graber, T., Pandharipande, R.: Localization of virtual classes. *Invent. Math.* 135(2), 487–518 (1999)
24. Graber, T., Vakil, R.: On the tautological ring of $\overline{\mathcal{M}}_{g,n}$. In: *Proceedings of the Seventh Gökova Geometry-Topology Conference 2000*, International Press, Somerville (2000)
25. Graber, T., Vakil, R.: Hodge integrals, Hurwitz numbers, and virtual localization. *Compositio Math.* 135(1), 25–36 (2003)
26. Graber, T., Vakil, R.: Relative virtual localization and vanishing of tautological classes on moduli spaces of curves. *Duke Math. J.* 130(1), 1–37 (2005)
27. Hori, K. et al.: *Gromov-Witten Theory and Mirror Symmetry*. Clay Math. Inst., Amer. Math. Soc. (2002)
28. Ionel, E.-N., Parker, T.H.: Relative Gromov-Witten invariants. *Ann. of Math.* (2) 157(1), 45–96 (2003)

29. Ionel, E.-N., Parker, T.H.: The symplectic sum formula for Gromov-Witten invariants. *Ann. of Math. (2)* 159(3), 935–1025 (2004)
30. Kazarian, M.E., Lando, S.K.: An algebro-geometric proof of Witten’s conjecture. *J. Amer. Math. Soc.* 20, 1079–1089 (2007)
31. Lee, Y.-P., Pandharipande, R.: Frobenius Manifolds, Gromov-Witten Theory, and Virasoro Constraints. to appear
32. Li, A.-M., Ruan, Y.: Symplectic surgery and Gromov-Witten invariants of Calabi-Yau 3-folds. *Invent. Math.* 145(1), 151–218 (2001)
33. Li, J.: Stable morphisms to singular schemes and relative stable morphisms. *J. Differential Geom.* 57(3), 509–578 (2001)
34. Li, J.: A degeneration formula of GW-invariants. *J. Differential Geom.* 60(2), 199–293 (2002)
35. Liu, K., Xu, H.: New properties of the intersection numbers on moduli spaces of curves. *Math. Res. Lett.* 14(6), 1041–1054 (2007)
36. Liu, K., Xu, H.: A proof of the Faber intersection number conjecture. *J. Differential Geom.* 83(2), 313–335 (2009)
37. Liu, C.-C.M., Liu, K., Zhou, J.: A proof of a conjecture of Mariño-Vafa on Hodge integrals. *J. Differential Geom.* 65(2), 289–340 (2003)
38. Looijenga, E.: On the tautological ring of \mathcal{M}_g . *Invent. Math.* 121(2), 411–419 (1995)
39. Okounkov, A., Pandharipande, R.: Gromov-Witten theory, Hurwitz numbers, and matrix models, I. In: Abramovich, D. et al. (eds.) *Algebraic Geometry—Seattle 2005*, pp. 325–414. Amer. Math. Soc., Providence (2009)
40. Pandharipande, R.: Hodge integrals and degenerate contributions. *Comm. Math. Phys.* 208(2), 489–506 (1999)
41. Pandharipande, R.: Three questions in Gromov-Witten theory. In: Li, D. (ed.) *Proceedings of the International Congress of Mathematicians, Vol. II (Beijing, 2002)*, pp. 503–512, Higher Ed. Press, Beijing (2002)
42. Shadrin, S., Shapiro, M., Vainshtein, A.: On double Hurwitz numbers in genus 0. In: *Proceedings of FPSAC’07. Tianjin, China (2007)*
43. Shadrin, S., Zvonkine, D.: Changes of variables in ELSV-type formulas. *Michigan Math. J.* 55(1), 209–228 (2007)
44. Vakil, R.: The moduli space of curves and its tautological ring. *Notices Amer. Math. Soc.* 50(6), 647–658 (2003)
45. Vakil, R.: The moduli space of curves and Gromov-Witten theory. In: Thaddeus, M. et al. (eds.) *Enumerative Invariants in Algebraic Geometry and String Theory*, pp. 143C198. Springer, Berlin (2008)