


Recall:




→ charts are tools to get a handle at the otherwise nameless abstract points of the manifold.

Problem:

How to define the abstract
"Tangent space, $T_p(M)$,"
to a diffable mfd at a point p ?

Intuition:



→ Proper definition should imply:

An n -dim mfd possesses for every point p an n -dim vector space of tangent vectors.

3 equivalent definitions of $T_p(M)$:

→ 1. "Algebraic" definition of $T_p(M)$:

Most powerful
b/c no need
for coordinates

Idea: $\square$ A tangent vector = directional derivative,
 $\square$ Derivatives definable through Leibniz rule:


$$(fg)' = f'g + fg'$$

2. "Physicist" definition of $T_p(M)$:

Idea: The elements of $T_p(M)$ are
to be vectors $\Rightarrow$ recognizable by how
their components change with charts.

3. "Geometric" definition of $T_p(M)$:

Idea: The elements of $T_p(M)$ are
to be actual tangent vectors
of one-dim. paths in the
manifold, that pass through p .



The 3 defs are equivalent, but:

We'll need all 3 occasionally!

→ we will do all 3:

1. Algebraic definition of $T_p(M)$

Idea: a) A tangent vector = directional derivative,

b) Derivatives definable through Leibniz rule:

$$(fg)' = f'g + fg'$$

Key example: $M = \mathbb{R}^n$

a) The tangent vectors ξ at a point p are identified with the directional 1st derivatives:

$$\xi = \sum_{i=1}^n \xi_i \frac{\partial}{\partial x_i} \Big|_{x=p}$$

b) Thus, tangent vectors at p should be those maps

$$\xi : f \rightarrow \xi(f) = \sum_{i=1}^n \xi_i \frac{\partial}{\partial x_i} f(x) \Big|_{x=p}$$

which obey the "Leibniz rule" at p :

$$\xi(fg) = \xi(f)g + f\xi(g) \Big|_{\text{at } p}$$

Q: How to express the local nature of $\xi \in T_p(M)$ properly?

A: ξ acts on function germs, not on functions.

Def: $\square$ Assume M, N are diffable mflds and $p \in M$.

$\square$ We say that two differentiable functions ϕ, ψ are germ-equivalent about p if in a neighborhood $U \subset M$ of p :

$$\phi(q) = \psi(q) \quad \forall q \in U$$

$\square$ Each such equivalence class of functions is called a germ at p .

$\square$ Then, the "germ" of ϕ at p , denoted $\bar{\phi}_p$, is the equivalence class of all functions ψ which are identical to ϕ in some neighborhood of p :

$$\psi \in \bar{\phi}_p \text{ if } \exists U_p \text{ (some open neighborhood of } p \text{ in } M) : \forall q \in U_p : \phi(q) = \psi(q)$$


↑
"there exists"

Notice: Assume $\phi: M \rightarrow N$ is diffable at $p \in M$.

Then all $\psi \in \bar{\phi}_p$ possess the same first derivative at p .

For example:

Consider germs of scalar functions f :



Note:

□ To specify a germ, it suffices to specify any arbitrary one of its functions.

□ The set of all germs at p is denoted $\mathcal{F}(p)$.

Note: □ One has for all $c \in \mathbb{R}$ and $f, g \in \mathcal{F}(p)$:

$$\overline{c \cdot f} = c \bar{f} \quad (a)$$

$$\overline{f \cdot g} = \bar{f} \bar{g} \quad (b)$$

$$\overline{f + g} = \bar{f} + \bar{g} \quad (c)$$

$\Rightarrow \mathcal{F}(p)$ obeys the axioms of an associative algebra.

Finally: Algebraic definition of $T_p(M)$

Recall idea: The elements of $T_p(M)$ are to be 1st derivatives $\Rightarrow$ definable by Leibniz rule.

Definition: The tangent space $T_p(M)$ is the set of "derivations" of $\mathcal{F}(p)$, i.e. the set of linear maps $\xi: \mathcal{F}(p) \rightarrow \mathbb{R}$ which obey:


$$\xi(\bar{f}_p \bar{g}_p) = \xi(\bar{f}_p) \cdot \bar{g}_p(p) + \bar{f}_p(p) \xi(\bar{g}_p)$$

$\begin{matrix} \parallel_{\xi(p)} & & \parallel_{f(p)} \\ \uparrow & \nearrow & \\ \text{remember this} & & (*) \end{matrix}$

Remark:

- ▢ this definition is abstract enough not only for arbitrary differentiable manifolds!
- ▢ this definition (as derivations of the algebra of functions) is also suitable for "Noncommutative Geometry": There, (Quantum Gravity) the algebra of functions $\mathcal{F}(p)$ is noncommutative.
- ▢ Note: Can't do Newton's derivatives then but algebraic def'n of derivation still works.

First example: a constant function, c , and its germ $\bar{c}$.



$c(x) := c$ and c is a constant: $c \in \mathbb{R}$

Then: $\xi(\bar{c}) = 0$ for all $\xi \in T_p(M)$

Proof: $\xi(\bar{c}) = c \xi(1) = c \xi(1 \cdot 1) \overset{\text{Leibniz rule}}{=} c(\xi(1) \cdot 1 + 1 \cdot \xi(1))$
 $= 2c \xi(1) \Rightarrow \xi(\bar{c}) = 0 \quad \checkmark$

Example: The case $M = \mathbb{R}^n$

If our definition for $T_p(M)$ is good, we expect that every $\xi \in T_p(M)$ is of the form:

$$\xi = \sum_{i=1}^n \xi^i \frac{\partial}{\partial x^i} \Big|_{x=p}$$

Proof:

□ We choose p to have coordinates $x = (0, 0, \dots)$.

□ Assume $\xi \in T_p(M)$ and $\bar{f} \in \mathcal{F}(p)$.

□ Notation: $h_{,i}(a', \dots, a^n) := \frac{\partial}{\partial a^i} h(a', \dots, a^n)$

Then:

(Note: these are not 3 numbers! These are 3 function germs, i.e., 3 equivalence classes of functions.)

$$\begin{aligned} \xi(\bar{f}(x)) &= \xi(\overline{f(0)} + \overline{f(x) - f(0)}) \\ &\stackrel{(a)}{=} \xi(\overline{f(0)} + \int_0^1 \frac{d}{dt} \bar{f}(tx', \dots, tx^n) dt) \end{aligned}$$

germ of a constant function
a constant function

$$\begin{aligned} &\stackrel{(b)}{=} \underbrace{\xi(\bar{f}(0))}_0 + \xi\left(\int_0^1 \sum_{i=1}^n \frac{\partial \bar{f}(tx', \dots, tx^n)}{\partial (tx^i)} \frac{d(tx^i)}{dt} dt\right) \\ &= \xi\left(\int_0^1 \sum_{i=1}^n \bar{f}_{,i}(tx', \dots, tx^n) \bar{x}^i dt\right) \end{aligned}$$

Linearity of $\xi \Rightarrow$

$$= \sum_{i=1}^n \xi \left(\int_0^1 \bar{f}_{,i}(tx', \dots, tx^n) dt \cdot \bar{x}^i \right)$$

Leibnitz rule $\Rightarrow$

$$= \sum_{i=1}^n \xi \left(\int_0^1 \bar{f}_{,i}(tx', \dots, tx^n) dt \right) \cdot \bar{x}^i \Big|_{x=p=0}^{\overset{=0}{}} + \sum_{i=1}^n \left(\int_0^1 \bar{f}_{,i}(tx', \dots, tx^n) dt \right) \Big|_{x=p=0} \cdot \xi(\bar{x}^i)$$

(Remember from ~~above~~ above)

$$\left(\int_0^1 c dt = c \Rightarrow \right)$$

$$= \sum_{i=1}^n \xi(\bar{x}^i) \int_0^1 \bar{f}_{,i}(0, \dots, 0) dt$$

$$= \sum_{i=1}^n \xi(\bar{x}^i) \frac{\partial}{\partial x^i} f(x', \dots, x^n) \Big|_{x=p=0}$$

$\Rightarrow$ Indeed, every $\xi \in T_p(M)$ is of the form

$$\xi = \sum_{i=1}^n \xi^i \frac{\partial}{\partial x^i} \Big|_{x=p} \quad (\text{I})$$

namely with

$$\xi^i = \xi(\bar{x}^i) \quad (\text{II})$$

□

Notice: Knowing how ξ acts on the coordinate functions $\bar{x}^i$ yields ξ^i (from II) and thus it means we know how ξ acts on all functions $\bar{f} \in \mathcal{F}(p)$, namely through (I).


But:

▢ This was the simple example:

$$M = \mathbb{R}^n$$

▢ How does our definition of $T_p(M)$ work for $M \neq \mathbb{R}^n$, concretely?

▢ Recall:




h gives abstract points a name, i.e. makes them concrete.

▢ Problem: How to make abstract $\xi \in T_p(M)$ concrete?

▢ Solution: Make use of charts in clever way!


Preparation: $T_p(M)$ and Diffeomorphisms.

Consider two diffable manifolds, M and N :



Note: If $N = \mathbb{R}^n$, then ϕ is a chart.

(that's the case we'll need but it's easy to keep a general N too)




Here: $\square$ $F(q)$ and $F(p)$ are algebras of function (germs).

$\square$ Given $\mathcal{C}$ we obtain a map $\mathcal{C}^*: F(q) \rightarrow F(p)$

$$\mathcal{C}^*: g \rightarrow f = \mathcal{C}^*(g) \text{ with } f(x) = g(\mathcal{C}(x)) \quad \forall x \in M$$

$$\text{i.e.: } f = \mathcal{C}^*(g) = g \circ \mathcal{C} \quad (+)$$



Here: $\square$ Given $\mathcal{C}^*: F(q) \rightarrow F(p)$ we obtain the "tangent map":

$$T_p \mathcal{C}: T_p(M) \rightarrow T_q(N)$$

$$T_p \mathcal{C}: \xi \rightarrow \eta$$

(When choosing $M = \mathbb{R}^n$, we obtain the desired concrete representation of $T_p(M)$ this way)

□ Namely:

$$\gamma = \xi \circ \varphi^*$$

i.e.:

$$\gamma(g) = \xi(\varphi^*(g))$$


□ From (+) $\Rightarrow$

$$\gamma(g) = \xi(g \circ \varphi)$$

The crucial special case:

- $N = \mathbb{R}^n$ (with $n = \dim(N)$)
- φ is invertible
- $(\Rightarrow \varphi^*$ is algebra isomorphism)
- $\Rightarrow T_p \varphi$ is vector space isomorphism

$\Rightarrow$ We do obtain a concrete handle on the abstract tangent vectors $\xi \in T_p(M)$, given a chart h :



Namely:

- Given a chart $\mathcal{C}$, every abstract point $p \in M$ has a concrete image $\mathcal{C}(p) \in \mathbb{R}^n$, and:
- Every abstract vector $\xi \in T_p(M)$ has a concrete image $\eta \in T_{\mathcal{C}(p)}(\mathbb{R}^n)$ namely:

$$\eta = T_p \mathcal{C}(\xi)$$

$\underbrace{\mathcal{C}(p)}_{x} = q$
- The image η is concrete because η is tangent vector to a point $q \in \mathbb{R}^n$, and it therefore must take the

form (we showed this):

$$\eta = \sum_{i=1}^n \eta^i \frac{\partial}{\partial x^i} \Big|_{x=q}$$

$\uparrow$ concrete numbers.

Conversely: (and very conveniently)

- Assuming a fixed $\mathcal{C}$, any choice of a $q = (x^1, \dots, x^n)$ denotes a $p \in M$ and any choice of a $(\eta^1, \dots, \eta^n)$ denotes a $\xi \in T_p(M)$.

$\uparrow$ some numbers

□ E.g. $\eta = \frac{\partial}{\partial x^i} \Big|_{x=p}$ is the image

of some abstract $\xi \in T_p(M)$, for fixed p .

Notation: $\xi = \frac{\partial}{\partial x^i} \Big|_{x=p}$

$\uparrow$ symbolic notation

Next:

If we hold p and $\xi \in T_p(M)$ fixed,

how do the numbers $(x^1, \dots, x^n)$

and $(\eta^1, \dots, \eta^n)$ change when we

change the chart? $\rightarrow$ Physicists' def of $T_p(M)$