

A review of machine learning applications in wildfire science and management

Piyush Jain^{*1,2}, Sean C P Coogan², Sriram Ganapathi Subramanian^{†3}, Mark Crowley^{‡3}, Steve Taylor⁴, and Mike D Flannigan²

¹Natural Resources Canada, Canadian Forest Service, Northern Forestry Centre, Edmonton, AB

²Canadian Partnership for Wildland Fire Science, University of Alberta, Renewable Resources, Edmonton, AB

³University of Waterloo, Electrical and Computer Engineering, Waterloo, ON

⁴Natural Resources Canada, Canadian Forest Service, Pacific Forestry Centre, Victoria, BC

July 27, 2020

Abstract

Artificial intelligence has been applied in wildfire science and management since the 1990s, with early applications including neural networks and expert systems. Since then the field has rapidly progressed congruently with the wide adoption of machine learning (ML) methods in the environmental sciences. Here, we present a scoping review of ML applications in wildfire science and management. Our overall objective is to improve awareness of ML methods among wildfire researchers and managers, as well as illustrate the diverse and challenging range of problems in wildfire science available to ML data scientists. To that end, we first present an overview of popular ML approaches used in wildfire science to date, and then review the use of ML in wildfire science as broadly categorized into six problem domains, including: 1) fuels characterization, fire detection, and mapping; 2) fire weather and climate change; 3) fire occurrence, susceptibility, and risk; 4) fire behavior prediction; 5) fire effects; and 6) fire management. Furthermore, we discuss the advantages and limitations of various ML approaches relating to data size, computational requirements, generalizability, and interpretability, as well as identify opportunities for future advances in the science and management of wildfires within a data science context. In total, we identified 300 relevant publications up to the end of 2019, where the most frequently used ML methods across problem domains included random forests, MaxEnt, artificial neural networks, decision trees, support vector machines, and genetic algorithms. As such, there exists opportunities to apply more current ML methods — including deep learning and agent based learning — in the wildfire sciences, especially in instances involving very large multivariate datasets. We must recognize, however, that despite the ability of ML methods to learn on their own, expertise in wildfire science is necessary to ensure realistic modelling of fire processes across multiple scales, while the complexity of some ML methods, such as deep learning, requires a dedicated and sophisticated knowledge of their application. Finally, we stress that the wildfire research and management communities play an active role in providing relevant, high quality, and freely available wildfire data for use by practitioners of ML methods.

Keywords: *machine learning, wildfire science, fire management, wildland fire, support vector machine, artificial neural network, decision trees, Bayesian networks, reinforcement learning, deep learning*

*piyush.jain@canada.ca

†ss@uwaterloo.ca

‡mcrowley@uwaterloo.ca

1 Introduction

Wildland fire is a widespread and critical element of the earth system [Bond and Keeley, 2005], and is a continuous global feature that occurs in every month of the year. Presently, global annual area burned is estimated to be approximately 420 Mha [Giglio et al., 2018], which is greater in area than the country of India. Globally, most of the area burned by wildfires occurs in grasslands and savannas. Humans are responsible for starting over 90% of wildland fires, and lightning is responsible for almost all of the remaining ignitions. Wildland fires can result in significant impacts to humans, either directly through loss of life and destruction to communities, or indirectly through smoke exposure. Moreover, as the climate warms we are seeing increasing impacts from wildland fire [Coogan et al., 2019]. Consequently, billions of dollars are spent every year on fire management activities aimed at mitigating or preventing wildfires' negative effects. Understanding and better predicting wildfires is therefore crucial in several important areas of wildfire management, including emergency response, ecosystem management, land-use planning, and climate adaptation to name a few.

Wildland fire itself is a complex process; its occurrence and behaviour are the product of several interrelated factors, including ignition source, fuel composition, weather, and topography. Furthermore, fire activity can be examined viewed across a vast range of scales, from ignition and combustion processes that occur at a scale of centimeters over a period of seconds, to fire spread and growth over minutes to days from meters to kilometers. At larger extents, measures of fire frequency may be measured over years to millennia at regional, continental, and planetary scales (see Simard [1991] for a classification of fire severity scales, and Taylor et al. [2013] for a review of numerical and statistical models that have been used to characterize and predict fire activity at a range of scales). For example, combustion and fire behavior are fundamentally physicochemical processes that can be usefully represented in mechanistic (i.e., physics-based) models at relatively fine scales [Coen, 2018]. However, such models are often limited both by the ability to resolve relevant physical processes, as well as the quality and availability of input data [Hoffman et al., 2016]. Moreover, with the limitations associated with currently available computing power it is not feasible to apply physical models to inform fire management and research across the larger and longer scales that are needed and in near real time. Thus, wildfire science and management rely heavily on the development of empirical and statistical models for meso, synoptic, strategic, and global scale processes [Simard, 1991], the utility of which are dependent upon their ability to represent the often complex and non-linear relationships between the variables of interest, as well as by the quality and availability of data.

While the complexities of wildland fire often present challenges for modelling, significant advances have been made in wildfire monitoring and observation primarily due to the increasing availability and capability of remote-sensing technologies. Several satellites (eg. NASA TERRA, AQUA and GOES), for instance, have onboard fire detection sensors (e.g., Advanced Very High Resolution Radiometer (AVHRR), Moderate Resolution Imaging Spectroradiometer (MODIS), Visible Infrared Imaging Radiometer Suite (VIIRS)), and these sensors along with those on other satellites (e.g., LANDSAT series) routinely monitor vegetation distributions and changes. Additionally, improvements in numerical weather prediction and climate models are simultaneously offering smaller spatial resolutions and longer lead forecast times [Bauer et al., 2015] which potentially offer improved predictability of extreme fire weather events. Such developments make a data-centric approach to wildfire modeling a natural evolution for many research problems given sufficient data. Consequently, there has been a growing interest in the use of Machine Learning (ML) methodologies in wildfire science and management in recent years.

Although no formal definition exists, we adopt the conventional interpretation of ML as the study of computer algorithms that can improve automatically through experience [Mitchell, 1997]. This approach is necessarily data-centric, with the performance of ML algorithms dependent on the quality and quantity of available data relevant to the task at hand. The field of ML has undergone an explosion of new algorithmic advances in recent years and is deeply connected to the broader field of Artificial Intelligence (AI). AI researchers aim to understand and synthesize intelligent agents which can act appropriately to their situation and objectives, adapt to changing environments, and learn from experience [Poole and Mackworth,

86 2010]. The motivations for using AI for forested ecosystem related research, including disturbances due to
 87 wildfire, insects, and disease, were discussed in an early paper [Schmoldt, 2001], while Olden et al. [2008]
 88 further argued for the use of ML methods to model complex problems in ecology. The use of ML models
 89 in the environmental sciences has seen a rapid uptake in the last decade, as is evidenced by recent reviews
 90 in the geosciences [Karpatne et al., 2017], forest ecology [Liu et al., 2018], extreme weather prediction
 91 [McGovern et al., 2017], flood forecasting [Mosavi et al., 2018], statistical downscaling [Vandal et al., 2018],
 92 remote sensing [Lary et al., 2016], and water resources [Shen, 2018, Sun and Scanlon, 2019]. Two recent
 93 perspectives have also made compelling arguments for the application of deep learning in earth system
 94 sciences [Reichstein et al., 2019] and for tackling climate change [Rolnick et al., 2019]. To date, however,
 95 no such paper has synthesized the diversity of ML approaches used in the various challenges facing wildland
 96 fire science.

97 In this paper, we review the current state of literature on ML applications in wildfire science and
 98 management. Our overall objective is to improve awareness of ML methods among fire researchers and
 99 managers, and illustrate the diverse and challenging problems in wildfire open to data scientists. This
 100 paper is organized as follows. In Section 2, we discuss commonly used ML methods, focusing on those
 101 most commonly encountered in wildfire science. In Section 3, we give an overview of the scoping review
 102 and literature search methodology employed in this paper. In this section we also highlight the results of
 103 our literature search and examine the uptake of ML methods in wildfire science since the 1990s. In Section
 104 4, we review the relevant literature within six broadly categorized wildfire modeling domains: (i) Fuels
 105 characterization, fire detection, and mapping; (ii) fire weather and climate change; (iii) fire probability
 106 and risk; (iv) fire behavior prediction; (v) fire effects; and (vi) fire management. In Section 5, we discuss
 107 our findings and identify further opportunities for the application of ML methods in wildfire science and
 108 management. Finally, in Section 6 we offer conclusions. Thus, this review will serve to guide and inform
 109 both researchers and practitioners in the wildfire community looking to use ML methods, as well as provide
 110 ML researchers the opportunity to identify possible applications in wildfire science and management.

111 2 Artificial Intelligence and Machine Learning

112 “**Definition: Machine Learning** - (Methods which) detect patterns in data, use the uncov-
 113 ered patterns to predict future data or other outcomes of interest”
 114 from *Machine Learning: A Probabilistic Perspective, 2012* [Murphy, 2012].

115 ML itself can be seen as a branch of AI or statistics, depending who you ask, that focuses on building
 116 predictive, descriptive, or actionable models for a given problem by using collected data, or incoming
 117 data, specific to that problem. ML methods learn directly from data and dispense with the need for
 118 a large number of expert rules or the need to model individual environmental variables with perfect
 119 accuracy. ML algorithms develop their own internal model of the underlying distributions when learning
 120 from data and thus need not be explicitly provided with physical properties of different parameters. Take
 121 for example, the task of modeling wildland fire spread, the relevant physical properties which include fuel
 122 composition, local weather and topography. The current state of the art in wildfire prediction includes
 123 physics-based simulators that fire fighters and strategic planners rely on to take many critical decisions
 124 regarding allocation of scarce firefighting resources in the event of a wildfire [Sullivan, 2007]. These physics-
 125 based simulators, however, have certain critical limitations; they normally render very low accuracies, have
 126 a prediction bias in regions where they are designed to be used, are often hard to design and implement due
 127 to the requirement of large number of expert rules. Furthermore, modelling many complex environmental
 128 variables is often difficult due to large resource requirements and complex or heterogeneous data formats.
 129 ML algorithms, however, learn their own mappings between parametric rules directly from data and do
 130 not require expert rules, which is particularly advantageous when the number of parameters are quite large
 131 and their physical properties quite complex, as in the case of wildland fire. Therefore, a ML approach to
 132 wildfire response may help to avoid many of the limitations of physics-based simulators.

A major goal of this review is to provide an overview of the various ML methods utilized in wildfire science and management. Importantly, we also provide a generalized framework for guiding wildfire scientists interested in applying ML methods to specific problem domains in wildland fire research. This conceptual framework, derived from the approach in [Murphy, 2012] and modified to show examples relevant to wildland fire and management is shown in Fig. 1. In general, ML methods can be identified as belonging to one of three types: supervised learning; unsupervised learning; or, agent based learning. We describe each of these below.

Supervised Learning - In supervised ML all problems can be seen as one of learning a parametrized function, often called a “model”, that maps inputs (i.e., predictor variables) to outputs (or “target variables”) both of which are known. The goal of supervised learning is to use an algorithm to learn the parameters of that function using available data. In fact, both linear and logistic regression can be seen as very simple forms of supervised learning. Most of the successful and popular ML methods fall into this category.

Unsupervised Learning - If the target variables are not available, then ML problems are typically much harder to solve. In unsupervised learning, the canonical tasks are dimensionality reduction and clustering, where relationships or patterns are extracted from the data without any guidance as to the “right” answer. Extracting embedded dimensions which minimize variance, or assigning datapoints to (labelled) classes which maximize some notion of natural proximity or other measures of similarity are examples of unsupervised ML tasks.

Agent Based Learning - Between supervised and unsupervised learning are a group of ML methods where learning happens by simulating behaviors and interactions of a single or a group of autonomous agents. These are general unsupervised methods which use incomplete information about the target variables, (i.e., information is available for some instances but not others), requiring generalizable models to be learned. A specific case in this space is Reinforcement Learning [Sutton and Barto, 1998], which is used to model decision making problems over time where critical parts of the environment can only be observed interactively through trial and error. This class of problems arises often in the real world and require efficient learning and careful definition of values (or preferences) and exploration strategies.

In the next section, we present a brief introduction to commonly used ML methods from the aforementioned learning paradigms. We note that this list is not meant to be exhaustive, and that some methods can accommodate both supervised and unsupervised learning tasks. It should be noted that the classification of a method as belonging to either ML or traditional statistics is often a question of taste. For the purpose of this review — and in the interests of economy — we have designated a number of methods as belonging to traditional statistics rather than ML. For a complete listing see tables 1 and 2.

2.1 Decision Trees

Decision Trees (DT) [Breiman, 2017] belong to the class of supervised learning algorithms and are another example of a universal function approximator, although in their basic form such universality is difficult to achieve. DTs can be used for both classification and regression problems. A decision tree is a set of if-then-else rules with multiple branches joined by decision nodes and terminated by leaf nodes. The decision node is where the tree splits into different branches, with each branch corresponding to the particular decision being taken by the algorithm whereas leaf nodes represent the model output. This could be a label for a classification problem or a continuous value in case of a regression problem. A large set of decision nodes is used in this way to build the DT. The objective of DTs are to accurately capture the relationships between input and outputs using the smallest possible tree that avoids overfitting. C4.5 [Quinlan, 1993] and Classification and Regression Trees (CART, [Breiman et al., 1984]) are examples of common single DT algorithms. Note that while the term CART is also used as an umbrella term for single tree methods, we use DT here to refer to all such methods. The majority of decision tree applications are ensemble decision tree (EDT) models that use multiple trees in parallel (ie. bootstrap aggregation or bagging) or sequentially (ie., boosting) to arrive at a final model. In this way, EDTs make use of many weak learners to form a strong learner while being robust to overfitting. EDTs are well described in many ML/AI textbooks and




Figure 1: A diagram showing the main machine learning types, types of data, and modeling tasks in relation to popular algorithms and potential applications in wildfire science and management. Note that the algorithms shown bolded are core ML methods whereas non-bolded algorithms are often not considered ML.

182 are widely available as implemented libraries.

183 2.1.1 Random Forests

184 A Random Forest (RF) [Breiman, 2001] is an ensemble model composed of a many individually trained
 185 DTs, and is the most popular implementation of a bagged decision tree. Each component DT in a RF
 186 model makes a classification decision where the class with the maximum number of votes is determined
 187 to be the final classification for the input data. RFs can also be used for regression where the final
 188 output is determined by averaging over the individual tree outputs. The underlying principle of the RF
 189 algorithm is that a random subset of features is selected at each node of each tree; the samples for training
 190 each component tree are selected using bagging, which resamples (with replacement) the original set of
 191 datapoints. The high performance of this algorithm is achieved by minimizing correlation between trees
 192 while reducing model variance so that a large number of different trees provides greater accuracy than
 193 individual trees. However, this improved performance comes at the cost of an increase in bias and loss of
 194 interpretability (although variable importance can still be inferred through permutation tests).

Machine Learning Methods

A3C	Asynchronous Advantage Actor-Critic
AdaBoost	Adaptive Boosting
ANFIS	Adaptive Neuro Fuzzy Inference System
ANN	Artificial Neural Networks
ADP	Approximate Dynamic Programming (a.k.a. reinforcement learning)
Bag	Bagged Decision Trees
BN	Bayesian Networks
BRT	Boosted Regression Trees (a.k.a. Gradient Boosted Machine)
BULC	Bayesian Updating of Land Cover
CART	Classification and Regression Tree
CNN	Convolutional Neural Network
DNN	Deep Neural Network
DQN	Deep Q-Network
DT	Decision Trees (incl. CART, J48, jRip)
EDT	Ensemble Decision Trees (incl. bagging and boosting)
ELM	Extreme Machine Learning (i.e., feedforward network)
GA	Genetic algorithms (a.k.a evolutionary algorithms)
GBM	Gradient Boosted Machine (a.k.a. Boosted Regression Trees, incl. XGBoost, AdaBoost, LogitBoost)
GMM	Gaussian Mixture Models
GP	Gaussian Processes
HCL	Hard Competitive Learning
HMM	Hidden Markov Models
ISODATA	Iterative Self-Organizing DATA algorithm
KNN	K Nearest Neighbor
KM	K-means Clustering
LB	LogitBoost (incl. AdaBoost)
LSTM	Long Short Term Memory
MaxEnt	Maximum Entropy
MCMC	Markov Chain Monte Carlo
MCTS	Monte Carlo Tree Search
MLP	Multilayer Perceptron
MDP	Markov Decision Process
NB	Naive Bayes
NFM	Neuro-Fuzzy models
PSO	Particle Swarm Optimization
RF	Random Forest
RL	Reinforcement Learning
RNN	Recurrent Neural Network
SGB	Stochastic Gradient Boosting
SOM	Self-organizing Maps
SVM	Support Vector Machines
t-SNE	T-distributed Stochastic Neighbor Embedding

Table 1: Table of acronyms and definitions for common machine learning algorithms referred to in text.

195 **2.1.2 Boosted Ensembles**

196 Boosting describes a strategy where one combines a set of weak learners — usually decision trees — to
197 make a strong learner using a sequential additive model. Each successive model improves on the previous

Non-machine learning methods	
DBSCAN	Density-based spatial clustering of applications with noise
GAM	Generalized Additive Model
GLM	Generalized Linear Model
KLR	Kernel Logistic Regression
LDA	Linear Discriminant Analysis
LR	Logistic Regression
MARS	Multivariate Adaptive Regression Splines
MLR	Multiple Linear Regression
PCA	Principal Component Analysis
SLR	Simple Linear regression

Table 2: Table of acronyms and definitions for common data analysis algorithms usually considered as foundational to, or outside of, machine learning itself.

198 by taking into account the model errors from the previous model, which can be done in more than one way.
 199 For example, the adaptive boosting algorithm, known as AdaBoost [Freund and Shapire, 1995], works by
 200 increasing the weight of observations that were previously misclassified. This can in principle reduce the
 201 classification error leading to a high level of precision [Hastie et al., 2009].

202 Another very popular implementation for ensemble boosted trees is Gradient Boosting Machine (GBMs),
 203 which makes use of the fact that each DT model represents a function that can be differentiated with re-
 204 spect to its parameters, i.e., how much a change in the parameters will change the output of the function.
 205 GBMs sequentially build an ensemble of multiple weak learners by following a simple gradient which points
 206 in the opposite direction to weakest results of the current combined model [Friedman, 2001].

207 The details for the GBM algorithm are as follows. Denoting the target output as Y , and given a
 208 tree-based ensemble model, represented as a function $T_i(X) \rightarrow Y$, after adding i weak learners already,
 209 the “perfect” function for the $(i + 1)$ th weak learner would be $h(x) = T_i(x) - Y$ which exactly corrects the
 210 previous model (i.e., $T_{(i+1)}(x) = T_i(x) + h(x) = Y$). In practice, we can only approach this perfect update
 211 by performing functional gradient descent where we use an approximation of the true residual (i.e., loss
 212 function) at each step. In our case this approximation is simply the sum of the residuals from each weak
 213 learner decision tree $L(Y, T(X)) = \sum_i Y - T_i(X)$. GBM explicitly uses the gradient $\nabla_{T_i} L(Y, T_i(X))$ of the
 214 loss function of each tree to fit a new tree and add it to the ensemble.

215 In a number of domains, and particularly in the context of ecological modeling GBM is often referred
 216 to as Boosted Regression Trees (BRTs) [Elith et al., 2008]. For consistency with the majority of literature
 217 reviewed in this paper we henceforth use the latter term. It should be noted that while deep neural networks
 218 (DNNs) and EDT methods are both universal function approximators, EDTs are more easily interpretable
 219 and faster to learn with less data than DNNs. However, there are fewer and fewer cases where trees-based
 220 methods can be shown to provide superior performance on any particular metric when DNNs are trained
 221 properly with enough data (see for example, Korotcov et al. [2017]).

222 2.2 Support Vector Machines

223 Another category of supervised learning includes Support Vector Machines (SVM) [Hearst et al., 1998] and
 224 related kernel-based methods. SVM is a classifier that determines the hyper-plane (decision boundary)
 225 in an n -dimensional space separating the boundary of each class, for data in n dimensions. SVM finds
 226 the optimal hyper-plane in such a way that the distance between the nearest point of each class to the
 227 decision boundary is maximized. If the data can be separated by a line then the hyper-plane is defined to
 228 be of the form $w^T x + b = 0$ where the w is the weight vector, x is the input vector and b is the bias. The
 229 distance of the hyper-plane to the closest data point d , called a support vector, is defined as the margin
 230 of separation. The objective is to find the optimal hyper-plane that minimizes the margin. If they are

not linearly separable, kernel SVM methods such as Radial Basis Functions (RBF) first apply a set of transformations to the data to a higher dimensional space where finding this hyperplane would be easier. SVMs have been widely used for both classification and regression problems, although recently developed deep learning algorithms have proved to be more efficient than SVMs given a large amount of training data. However, for problems with limited training samples, SVMs might give better performances than deep learning based classifiers.

2.3 Artificial Neural Networks and Deep Learning

The basic unit of an Artificial Neural Network (ANN) is a neuron (also called a perceptron or logistic unit). A neuron is inspired by the functioning of neurons in mammalian brains in that it can learn simple associations, but in reality it is much simpler than its biological counterpart. A neuron has a set of inputs which are combined linearly through multiplication with weights associated with the input. The final weighted sum forms the output signal which is then passed through a (generally) non-linear activation function. Examples of activation functions include sigmoid, tanh, and the Rectified Linear Unit (ReLU). This non-linearity is important for general learning since it creates an abrupt cutoff (or threshold) between positive and negative signals. The weights on each connection represent the function parameters which are fit using supervised learning by optimizing the threshold so that it reaches a maximally distinguishing value.

In practice, even simple ANNs, often called Multi-Layered Perceptrons (MLP), combine many neuron units in parallel, each processing the same input with independent weights. In addition, a second layer of hidden neuron units can be added to allow more degrees of freedom to fit general functions, see Figure 2(a). MLPs are capable of solving simple classification and regression problems. For instance, if the task is one of classification, then the output is the predicted class for the input data, whereas in the case of a regression task the output is the regressed value for the input data. Deep learning [LeCun et al., 2015] refers to using Deep Neural Networks (DNNs) which are ANNs with multiple hidden layers (nominally more than 3) and include Convolutional Neural Networks (CNNs) popularized in image analysis and Recurrent Neural Networks (RNNs) which can be used to model dynamic temporal phenomena. The architecture of DNNs can vary in connectivity between nodes, the number of layers employed, the types of activation functions used, and many other types of hyperparameters. Nodes within a single layer can be fully connected, or connected with some form of convolutional layer (e.g., CNNs), recurrent units (e.g., RNNs), or other sparse connectivity. The only requirement of all these connectivity structures and activation functions is that they are differentiable.

Regardless of the architecture, the most common process of training a ANN involves processing input data fed through the network layers and activation functions to produce an output. In the supervised setting, this output is then compared to the known true output (i.e., labelled training data) resulting in an error measurement (loss or cost function) used to evaluate model performance. The error for DNNs are commonly calculated as a cross entropy loss between the predicted output label and the true output label. Since every part of the network is mathematically differentiable we can compute a gradient for the entire network. This gradient is used to calculate the proportional change in each network weight needed to produce an infinitesimal increase in the likelihood of the network producing the same output for the most recent output. The gradient is then weighted by the computed error, and thereafter all the weights are updated in sequence using a backpropagation algorithm [Hecht-Nielsen, 1992].

ANNs can also be configured for unsupervised learning tasks. For example, self-organizing maps (SOMs) are a form of ANN adapted for dealing with spatial data and have therefore found widespread use in the atmospheric sciences [Skific and Francis, 2012]. A SOM is a form of unsupervised learning that consists of a two-dimensional array of nodes as the input layer, representing say, a gridded atmospheric variable at a single time. The algorithm clusters similar atmospheric patterns together and results in a dimensionality reduction of the input data. More recently, unsupervised learning methods from deep learning, such as autoencoder networks, are starting to replace SOMs in the environmental sciences [Shen, 2018].




Figure 2: Logistic regression can be seen as basic building block for neural networks, with no hidden layer and a sigmoid activation function. Classic shallow neural networks (also known as Multi-Layer Perceptrons) have at least one hidden layer and can have a variety of activation functions. Deep neural networks essentially have a much larger number of hidden layers as well as use additional regularization and optimization methods to enhance training.

279 **2.4 Bayesian methods**

280 **2.4.1 Bayesian Networks**

281 Bayesian networks (Bayes net, belief network; BN) are a popular tool in many applied domains because
 282 they provide an intuitive graphical language for specifying the probabilistic relationships between variables
 283 as well as tools for calculating the resulting probabilities [Pearl, 1988]. The basis of BNs is Bayes' theorem,
 284 which relates the conditional and marginal probabilities of random variables. BNs can be treated as a ML
 285 task if one is trying to automatically fit the parameters of the model from data, or even more challenging,
 286 to learn the best graphical structure that should be used to represent a dataset. BNs have close ties to
 287 causal reasoning, but it is important to remember that the relationships encoded in a BN are inherently
 288 correlational rather than causal. BNs are acyclic graphs, consisting of nodes and arrows (or arcs), defining
 289 a probability distribution over variables \mathcal{U} . The set of parents of a node (variable) X , denoted π_X , are all
 290 nodes with directed arcs going into X . BNs provide compact representation of conditional distributions
 291 since $p(X_i|X_1, \dots, X_{i-1}) = p(X_i|\pi_{X_i})$ where X_1, \dots, X_{i-1} are arranged to be all of the ancestors of X_i
 292 other than its direct parents. Each node X is associated with a conditional probability table over X and
 293 its parents defining $p(X|\pi_X)$. If a node has no parents, a prior distribution is specified for $p(X)$. The joint
 294 probability distribution of the network is then specified by the chain rule $P(U) = \prod_{X \in \mathcal{U}} p(X|\pi_X)$.

295 **2.4.2 Naïve Bayes**

296 A special case of a BN is the Naïve Bayes (NB) classifier, which assumes conditional independence between
 297 input features, which allows the likelihood function to be constructed by a simple multiplication of the
 298 conditional probability of each input variable conditional on the output. Therefore, while NB is fast

299 and straightforward to implement, prediction accuracy can be low for problems where the assumption of
300 conditional independence does not hold.

301 2.4.3 Maximum Entropy

302 Maximum Entropy (MaxEnt), originally introduced by [Phillips et al. \[2006\]](#), is a presence only framework
303 that fits a spatial probability distribution by maximising entropy, consistent with existing knowledge.
304 MaxEnt can be considered a Bayesian method since it is compatible with an application of Bayes Theorem
305 as existing knowledge is equivalent to specifying a prior distribution. MaxEnt has found widespread use
306 in landscape ecology species distribution modeling [[Elith, Phillips, Hastie, Dudík, Chee, and Yates, 2011](#)],
307 where prior knowledge consists of occurrence observations for the species of interest.

308 2.5 Reward based methods

309 2.5.1 Genetic Algorithms

310 Genetic algorithms (GA) are heuristic algorithms inspired by Darwin’s theory of evolution (natural selec-
311 tion) and belong to a more general class of evolutionary algorithms [[Mitchell, 1996](#)]. GAs are often used to
312 generate solutions to search and optimization problems by using biologically motivated operators such as
313 mutation, crossover, and selection. In general, GAs involve several steps. The first step involves creating
314 an initial population of potential solutions, with each solution encoded as a chromosome. Second a fitness
315 function appropriate to the problem is defined, which returns a fitness score determining how likely an
316 individual is to be chosen for reproduction. The third step requires the selection of pairs of individuals,
317 denoted as parents. In the fourth step, a new population of finite individuals are created by generating
318 two new offspring from each set of parents using crossover, whereby a new chromosome is created by some
319 random selection process from each parents chromosomes. In the final step called mutation, a small sample
320 of the new population is chosen and a small perturbation is made to the parameters to maintain diversity.
321 The entire process is repeated many times until the desired results are satisfactory (based on the fitness
322 function), or some measure of convergence is reached.

323 2.5.2 Reinforcement Learning

324 Reinforcement learning (RL) represents a very different learning paradigm to supervised or unsupervised
325 learning. In RL, an agent (or actor) interacts with its environment and learns a desired behavior (set of
326 actions) in order to maximize some reward. RL is a solution to a Markov Decision Process (MDP) where
327 the transition probabilities are not explicitly known but need to be learned. This type of learning is well
328 suited to problems of automated decision making, such as required for automated control (e.g., robotics)
329 or for system optimization (e.g., management policies). Various RL algorithms include Monte Carlo Tree
330 Search (MTCS), Q-Learning, and Actor-Critic algorithms. For an introduction to RL see [Sutton and Barto](#)
331 [\[2018\]](#).

332 2.6 Clustering methods

333 Clustering is the process of splitting a set of points into groups where each point in a group is more similar to
334 its own group than any other group. There are different ways in which clustering can be done, for example,
335 the K-means (KM) clustering algorithm [[MacQueen et al., 1967](#)], based on a centroid model, is perhaps
336 the most well-known clustering algorithm. In K-means, the notion of similarity is based on closeness to
337 the centroid of each cluster. K-means is an iterative process in which the centroid of a group and points
338 belonging to a group are updated at each step. The K-means algorithm consists of five steps: (i) specify
339 the number of clusters; (ii) each data point is randomly assigned to a cluster; (iii) the centroids of each
340 cluster is calculated; (iv) the points are reassigned to the nearest centroids, and (v) cluster centroids are
341 recomputed. Steps iv and v repeat until no further changes are possible. Although KM is the most widely

342 used clustering algorithm, several other clustering algorithms exist including, for example, agglomerative
343 Hierarchical Clustering (HC), Gaussian Mixture Models (GMMs) and Iterative Self-Organizing DATA
344 (ISODATA).

345 2.7 Other methods

346 2.7.1 K-Nearest Neighbor

347 The K-Nearest Neighbors (KNN) algorithm is a simple but very effective supervised classification algorithm
348 which is based on the intuitive premise that similar data points are in close proximity according to some
349 metric [Altman, 1992]. Specifically, a KNN calculates the similarity of data points to each other using the
350 Euclidean distance between the K nearest data points. The optimal value of K can be found experimentally
351 over a range values using the classification error. KNN is widely used in applications where a search query
352 is performed such that results should be similar to another pre-existing entity. Examples of this include
353 finding similar images to a specified image and recommender systems. Another popular application of
354 KNN is outlier (or anomaly) detection, whereby the points (in a multidimensional space) farthest away
355 from their nearest neighbours may be classified as outliers.

356 2.7.2 Neuro-Fuzzy models

357 Fuzzy logic is an approach for encoding expert human knowledge into a system by defining logical rules
358 about how different classes overlap and interact without being constrained to “all-or-nothing” notions of
359 set inclusion or probability of occurrence. Although early implementations of fuzzy logic systems depended
360 on setting rules manually, and therefore are not considered machine learning, using fuzzy rules as inputs
361 or extracting them from ML methods are often described as “neuro-fuzzy” methods. For example, the
362 Adaptive Neuro-Fuzzy Inference System (ANFIS) [Jang, 1993] fuses fuzzy logical rules with an ANN
363 approach, while trying to maintain the benefits of both. ANFIS is a universal function approximator
364 like ANNs. However, since this algorithm originated in the 1990s, it precedes the recent deep learning
365 revolution so is not necessarily appropriate for very large data problems with complex patterns arising in
366 high-dimensional spaces. Alternatively, human acquired fuzzy rules can be integrated into ANNs learning;
367 however, it is not guaranteed that the resulting trained neural network will still be interpretable. It
368 should be noted that fuzzy rules and fuzzy logic are not a major direction of research within the core ML
369 community.

370 3 Literature search and scoping review

371 The combination of ML and wildfire science and management comprises a diverse range of topics in a rela-
372 tively nascent field of multidisciplinary research. Thus, we employed a scoping review methodology [Arksey
373 and O’Malley, 2005, Levac et al., 2010] for this paper. The goal of a scoping review is to characterize the
374 existing literature in a particular field of study, particularly when a topic has yet to be extensively reviewed
375 and the related concepts are complex and heterogeneous [Pham, Rajić, Greig, Sargeant, Papadopoulos,
376 and Mcewen, 2014]. Furthermore, scoping reviews can be particularly useful for summarizing and dissem-
377 inating research findings, and for identifying research gaps in the published literature. A critical review of
378 methodological advances and limitations and comparison with other methods is left for future work. We
379 performed a literature search using the Google Scholar and Scopus databases and the key words “wild-
380 fire” or “wildland fire” or “forest fire” or “bushfire” in combination with “machine learning” or “random
381 forest” or “decision trees” or “regression trees” or “support vector machine” or “maximum entropy” or
382 “neural network” or “deep learning” or “reinforcement learning”. We also used the Fire Research Institute
383 online database (<http://fireresearchinstitute.org>) using the following search terms: “Artificial In-
384 telligence”; “Machine Learning”; “Random Forests”; “Expert Systems”; and “Support Vector Machines”.




Figure 3: Number of publications by year for 300 publications on topic of ML and wildfire science and management as identified in this review.




Figure 4: Number of ML applications by category and by year for 300 publications on topic of ML and wildfire science and management as identified in this review.

385 Furthermore, we obtained papers from references cited within papers we had obtained using literature
 386 databases.

387 After performing our literature search, we identified a total of 300 publications relevant to the topic of
 388 ML applications in wildfire science and management (see supplementary material for a full bibliography).
 389 Furthermore, a search of the Scopus database revealed a dramatic increase in the number of wildfire and

390 ML articles published in recent years (see Fig. 3). After identifying publications for review, we further
 391 applied the following criteria to exclude non-relevant or unsuitable publications, including: (i) conference
 392 submissions where a journal publication describing the same work was available; (ii) conference posters;
 393 (iii) articles in which the methodology and results were not adequately described to conduct an assessment
 394 of the study; (iv) articles not available to as either by open access or by subscription; and (v) studies that
 395 did not present new methodologies or results.

396 4 Wildfire applications

397 In summary, we found a total of 300 journal papers or conference proceedings on the topic of ML applica-
 398 tions in wildfire science and management. We found the problem domains with the highest application of
 399 ML methods was *Fire Occurrence, Susceptibility and Risk* (127 papers) followed by *Fuels Characterization,*
 400 *Fire Detection And Mapping* (66 papers), *Fire Behaviour Prediction* (43 papers), *Fire Effects* (35 papers),
 401 *Fire Weather and Climate Change* (20 papers), and *Fire Management* (16 papers). Within Fire Occur-
 402 rence, Susceptibility and Risk, the subdomains with the most papers were *Fire Susceptibility Mapping* (71
 403 papers) and *Landscape Controls on Fire* (101 papers). Refer to table 3 and the supplementary material
 404 for a break-down of each problem subdomain and ML methods used, as well as study areas considered.

405 4.1 Fuels Characterization, Fire Detection, and Mapping

406 4.1.1 Fuels characterization

407 Fires ignite in a few fuel particles; subsequent heat transfer between particles through conduction, radiation
 408 and convection, and the resulting fire behavior (fuel consumption, spread rate, intensity) is influenced by
 409 properties of the live and dead vegetative fuels, including moisture content, biomass, and vertical and
 410 horizontal distribution. Fuel properties are a required input in all fire behavior models, whether it be
 411 a simple categorical vegetation type, as in the Canadian FBP System, or as physical quantities in a 3
 412 dimensional space (eg. see FIRETEC model). Research to predict fuel properties has been carried out
 413 at two different scales 1) regression applications to predict quantities such as the crown biomass of single
 414 trees from more easily measured variables such as height and diameter, and 2) classification applications to
 415 map fuel type descriptors or fuel quantities over a landscape from visual interpretation of air photographs
 416 or by interpretation of the spectral properties of remote sensing imagery. However, relatively few studies
 417 have employed ML to wildfire fuel prediction, leaving the potential for substantially more research in this
 418 area.

419 In an early study, [Riaño et al. \[2005\]](#) used an ANN to predict and map the equivalent water thickness
 420 and dry matter content of wet and dry leaf samples from 49 species of broad leaf plants using reflectance and
 421 transmittance values in the Ispra region of Italy. [Pierce et al. \[2012\]](#) used RF to classify important canopy
 422 fuel variables (e.g. canopy cover, canopy height, canopy base height, and canopy bulk density) related to
 423 wildland fire in Lassen Volcanic National Park, California, using field measurements, topographic data,
 424 and NDVI to produce forest canopy fuel maps. Likewise, [Viegas et al. \[2014\]](#) used RF with Landfire
 425 and biophysical variables to perform fuel classification and mapping in Eastern Oregon. The authors
 426 of the aforementioned study achieved relatively high overall modelling accuracy, for example, 97% for
 427 forest height, 86% for forest cover, and 84% for existing vegetation group (i.e. fuel type). [López-Serrano
 428 et al. \[2016\]](#) compared the performance of three common ML methods (i. SVM; ii. KNN; and iii. RF) and
 429 *multiple linear regression* in estimating above ground biomass in the Sierra Madre Occidental, Mexico. The
 430 authors reported the advantages and limitations of each method, concluding that that the *non-parametric*
 431 ML methods had an advantage over multiple linear regression for biomass estimation. [García et al. \[2011\]](#)
 432 used SVM to classify LiDAR and multispectral data to map fuel types in Spain. [Chirici et al. \[2013\]](#)
 433 compared the use of CART, RF, and Stochastic Gradient Boosting SGB, an ensemble tree method that
 434 uses both boosting and bagging, for mapping forest fuel types in Italy, and found that SGB had the highest
 435 overall accuracy.

Section	Domain	NFM	SVM	KM	GA	BN	BRT	ANN	DT	RF	KNN	MAXENT	DL	NB	Other
1.1	Fuels characterization	-	2	-	-	-	1	1	1	4	1	-	-	-	-
1.2	Fire detection	2	3	1	1	1	-	12	-	-	-	-	18	-	3
1.3	Fire perimeter and severity mapping	1	12	1	2	-	1	6	1	4	2	1	-	-	6
2.1	Fire weather prediction	-	-	1	-	-	-	-	-	1	-	-	-	-	3
2.2	Lightning prediction	-	-	-	-	-	-	-	1	2	-	-	-	-	-
2.3	Climate change	-	1	-	-	-	6	2	2	5	-	7	-	-	-
3.1	Fire occurrence prediction	-	3	-	-	1	-	7	1	5	1	2	-	1	4
3.2	Landscape-scale Burned area prediction	-	1	1	1	-	-	1	1	2	-	1	1	-	1
3.3	Fire Susceptibility Mapping	2	12	1	3	2	8	16	9	26	-	27	1	2	3
3.4	Landscape controls on fire	2	10	1	3	2	19	11	15	40	1	30	1	1	2
4.1	Fire Spread and Growth	-	-	-	13	2	-	4	-	1	1	-	3	-	2
4.2	Burned area and fire severity prediction	-	7	-	1	1	3	10	7	6	3	-	2	1	5
5.1	Soil erosion and deposits	-	-	1	-	-	-	1	1	-	-	1	-	-	-
5.2	Smoke and particulate levels	-	2	-	-	-	3	3	-	5	2	-	-	-	2
5.3	Post-fire regeneration and ecology	-	1	-	1	1	6	1	2	10	-	2	-	1	-
5.4	Socioeconomic effects	-	-	-	-	1	-	-	-	-	-	-	-	-	-
6.1	Planning and policy	-	-	-	1	1	-	-	-	2	-	-	-	-	2
6.2	Fuel treatment	-	-	-	1	1	-	-	-	-	-	-	-	-	1
6.3	Wildfire preparedness and response	-	-	-	1	2	1	1	-	-	-	1	1	-	1
6.4	Social factors	-	-	-	-	1	-	-	-	-	-	-	-	-	-

Table 3: Summary of application of ML methods applied to different problem domains in wildfire science and management. A table of acronyms for the ML methods are given in 1. Note that in some cases a paper may use more than one ML method and/or appear in multiple problem domains.

436 4.1.2 Fire detection

437 Detecting wildfires as soon as possible after they have ignited, and therefore while they are still relatively
438 small, is critical to facilitating a quick and effective response. Traditionally, fires have mainly been detected
439 by human observers, by distinguishing smoke in the field of view directly from a fire tower, or from a video
440 feed from a tower, aircraft, or from the ground. All of these methods can be limited by spatial or temporal
441 coverage, human error, the presence of smoke from other fires and by hours of daylight. Automated
442 detection of heat signatures or smoke in infra-red or optical images can extend the spatial and temporal
443 coverage of detection, the detection efficiency in smoky conditions, and remove bias associated with human
444 observation. The analytical task is a classification problem that is quite well suited to ML methods.

445 For example, [Arrue et al. \[2000\]](#) used ANNs for infrared (IR) image processing (in combination with
446 visual imagery, meteorological and geographic data used in a decision function using fuzzy logic), to identify
447 true wildfires. Several researchers have similarly employed ANNs for fire detection [[Al-Rawi et al., 2001](#),
448 [Angayarkkani and Radhakrishnan, 2010](#), [Fernandes et al., 2004a,b](#), [Li et al., 2015](#), [Soliman et al., 2010](#),
449 [Utkin et al., 2002](#), [Sayad et al., 2019](#)]. In addition, [Liu et al. \[2015\]](#) used ANNs on wireless sensor networks
450 to build a fire detection system, where multi-criteria detection was used on multiple attributes (e.g. flame,
451 heat, light, and radiation) to detect and raise alarms. Other ML methods used in fire detection systems
452 include SVM to automatically detect wildfires from videoframes [[Zhao et al., 2011](#)], GA for multi-objective
453 optimization of a LiDAR-based fire detection system [[Cordoba et al., 2004](#)], BN in a vision-based early fire
454 detection system [[Ko et al., 2010](#)], ANFIS [[Angayarkkani and Radhakrishnan, 2011](#), [Wang et al., 2011](#)],
455 and KM [[Srinivasa et al., 2008](#)].

456 CNNs (ie. deep learning), which are able to extract features and patterns from spatial images and
457 are finding widespread use in object detection tasks, have recently been applied to the problem of fire
458 detection. Several of these applications trained the models on terrestrial based images of fire and/or smoke
459 [[Zhang et al., 2016, 2018a,b](#), [Yuan et al., 2018](#), [Akhloufi et al., 2018](#), [Barmpoutis et al., 2019](#), [Jakubowski
460 et al., 2019](#), [João Sousa et al., 2019](#), [Li et al., 2018b, 2019](#), [Muhammad et al., 2018](#), [Wang et al., 2019](#)].
461 Of particular note, [Zhang et al. \[2018b\]](#) found CNNs outperformed a SVM-based method and [Barmpoutis
462 et al. \[2019\]](#) found a Faster region-based CNN outperformed another CNN based on YOLO (“you only look
463 once”). [Yuan et al. \[2018\]](#) used CNN combined with optical flow to include time-dependent information.
464 [Li et al. \[2018b\]](#) similarly used a 3D CNN to incorporate both spatial and temporal information and so
465 were able to treat smoke detection as a segmentation problem for video images. Another approach by [Cao
466 et al. \[2019\]](#) used convolutional layers as part of a Long Short Term Memory (LSTM) Neural network for
467 smoke detection from a sequence of images (ie. video feed). They found the LSTM method achieved 97.8%
468 accuracy, a 4.4% improvement over a single image-based deep learning method.

469 Perhaps of greater utility for fire management were fire/smoke detection models trained on either
470 unmanned aerial vehicle (UAV) images [[Zhao et al., 2018](#), [Alexandrov et al., 2019](#)] or satellite imagery
471 including GOES-16 [[Phan and Nguyen, 2019](#)] and MODIS [[Ba et al., 2019](#)]. [Zhao et al. \[2018\]](#) compared
472 SVM, ANN and 3 CNN models and found their 15-layer CNN performed best with an accuracy of 98%. By
473 comparison, the SVM based method, which was unable to extract spatial features, only had an accuracy of
474 43%. [Alexandrov et al. \[2019\]](#) found YOLO was both faster and more accurate than a region-based CNN
475 method in contrast to [Barmpoutis et al. \[2019\]](#).

476 4.1.3 Fire perimeter and severity mapping

477 Fire maps have two management applications: 1) Accurate maps of the location of the active fire perimeter
478 are important for daily planning of suppression activities and/or evacuations, including modeling fire
479 growth 2) Maps of the final burn perimeter and fire severity are important for assessing and predicting the
480 economic and ecological impacts of wildland fire and for recovery planning. Historically, fire perimeters were
481 sketch-mapped from the air, from a ground or aerial GPS or other traverse, or by air-photo interpretation.
482 Developing methods for mapping fire perimeters and burn severity from remote sensing imagery has been
483 an area of active research since the advent of remote sensing in the 1970s, and is mainly concerned with

484 classifying active fire areas from inactive or non burned areas, burned from unburned areas (for extinguished
485 fires), or fire severity measures such as the Normalized Burn Ratio [Lutes et al., 2006].

486 In early studies using ML methods for fire mapping Al-Rawi et al. [2001] and Al-Rawi et al. [2002] used
487 ANNs (specifically, the supervised ART-II neural network) for burned scar mapping and fire detection. Pu
488 and Gong [2004] compared Logistic Regression (LR) with ANN for burned scar mapping using Landsat
489 images; both methods achieved high accuracy ($> 97\%$). Interestingly, however, the authors found that
490 LR was more efficient for their relatively limited data set. The authors in Zammit et al. [2006] performed
491 burned area mapping for two large fires that occurred in France using satellite images and three ML
492 algorithms, including SVM, K-nearest neighbour, and the K-means algorithm; overall SVM had the best
493 performance. Likewise, E. Dragozi, I. Z. Gitas, D.G. Stavarakoudis [2011] compared the use of SVM against
494 a nearest neighbour method for burned area mapping in Greece and found better performance with SVM.
495 In fact, a number of studies [Alonso-Benito et al., 2008, Cao et al., 2009, Petropoulos et al., 2010, 2011, Zhao
496 et al., 2015, Pereira et al., 2017, Branham et al., 2017, Hamilton et al., 2017] have successfully used SVM
497 for burned scar mapping using satellite data. Mitrakis et al. [2012] performed burned area mapping in the
498 Mediterranean region using a variety of ML algorithms, including a fuzzy neuron classifier (FNC), ANN,
499 SVM, and AdaBoost, and found that, while all methods displayed similar accuracy, the FNC performed
500 slightly better. Dragozi et al. [2014] applied SVM and a feature selection method (based on fuzzy logic)
501 to IKONOS imagery for burned area mapping in Greece. Another approach to burned area mapping in
502 the Mediterranean used an ANN and MODIS hotspot data [Gómez and Pilar Martín, 2011]. Pereira et al.
503 [2017] used a one class SVM, which requires only positive training data (i.e. burned pixels), for burned
504 scar mapping, which may offer a more sample efficient approach than general SVMs – the one class SVM
505 approach may be useful in cases where good wildfire training datasets are difficult to obtain. In Mithal
506 et al. [2018], the authors developed a three-stage framework for burned area mapping using MODIS data
507 and ANNs. Crowley et al. [2019] used Bayesian Updating of Landcover (BULC) to merge burned-area
508 classifications from three remote sensing sources (Landsat-8, Sentinel-2 and MODIS). Celik [2010] used
509 GA for change detection in satellite images, while Sunar and Özkan [2001] used the interactive Iterative
510 Self-Organizing DATA algorithm (ISODATA) and ANN to map burned areas.

511 In addition to burned area mapping, ML methods have been used for burn severity mapping, including
512 GA [Brumby et al., 2001], MaxEnt [Quintano et al., 2019], bagged decision trees [Sá et al., 2003], and others.
513 For instance, Hultquist et al. [2014] used three popular ML approaches (Gaussian Process Regression (GPR)
514 [Rasmussen and Williams, 2006], RF, and SVM) for burn severity assessment in the Big Sur ecoregion,
515 California. RF gave the best overall performance and had lower sensitivity to different combinations of
516 variables. All ML methods, however, performed better than conventional multiple regression techniques.
517 Likewise, Hultquist et al. [2014] compared the use of GPR, RF, and SVM for burn severity assessment, and
518 found that RF displayed the best performance. Another recent paper by Collins et al. [2018] investigated
519 the applicability of RF for fire severity mapping, and discussed the advantages and limitations of RF for
520 different fire and land conditions.

521 One recent paper by Langford et al. [2019] used a 5-layer deep neural network (DNN) for mapping fires
522 in Interior Alaska with a number of MODIS derived variables (eg. NDVI and surface reflectance). They
523 found that a validation-loss (VL) weight selection strategy for the unbalanced data set (i.e., the no-fire
524 class appeared much more frequently than fire) allowed them to achieve better accuracy compared with a
525 XGBoost method. However, without the VL approach, XGBoost outperformed the DNN, highlighting the
526 need for methods to deal with unbalanced datasets in fire mapping.

527 4.2 Fire Weather and Climate Change

528 4.2.1 Fire weather prediction

529 Fire weather is a critical factor in determining whether a fire will start, how fast it will spread, and where
530 it will spread. Fire weather observations are commonly obtained from surface weather station networks
531 operated by meteorological services or fire management agencies. Weather observations may be interpolated

532 from these point locations to a grid over the domain of interest, which may include diverse topographical
533 conditions; the interpolation task is a regression problem. Weather observations may subsequently be
534 used in the calculation of meteorologically based fire danger indices, such as the Canadian Fire Weather
535 Index (FWI) System [Van Wagner, 1987]. Future fire weather conditions and danger indices are commonly
536 forecast using the output from numerical weather prediction (NWP) models (e.g., The European Forest
537 Fire Information System [San-Miguel-Ayanz et al., 2012]). However, errors in the calculation of fire danger
538 indices that have a memory (such as the moisture indices of the FWI System) can accumulate in such
539 projections. It is noteworthy that surface fire danger measures may be correlated with large scale weather
540 and climatic patterns.

541 To date there has been relatively few papers that address fire weather and danger prediction using ma-
542 chine learning. The first effort [Crimmins, 2006] used self-organizing maps (SOMs) to explore the synoptic
543 climatology of extreme fire weather in the southwest USA. He found three key patterns representing south-
544 westerly flow and large geopotential height gradients that were associated with over 80% of the extreme
545 fire weather days as determined by a fire weather index. Nauslar et al. [2019] used SOMs to determine the
546 timing of the North American Monsoon that plays a major role on the length of the active fire season in
547 the southwest USA. Lagerquist et al. [2017] also used SOMs to predict extreme fire weather in northern
548 Alberta, Canada. Extreme fire weather was defined by using extreme values of the Fine Fuel Moisture
549 Code (FFMC), Initial Spread Index (ISI) and the Fire Weather Index (FWI), all components of the Cana-
550 dian Fire Weather Index (FWI) System [Van Wagner, 1987]. Good performance was achieved with the
551 FFMC and the ISI and this approach has the potential to be used in near real time, allowing input into
552 fire management decision systems. Other efforts have used a combination of conventional and machine
553 learning approaches to interpolate meteorological fire danger in Australia [Sanabria et al., 2013].

554 4.2.2 Lightning prediction

555 Lightning is second most common cause of wildfires (behind human causes); thus predicting the location
556 and timing of future storms/strikes is of great importance to predicting fire occurrence. Electronic lightning
557 detection systems have been deployed in many parts of the world for several decades and have accrued rich
558 strike location/time datasets. Lightning prediction models have employed these data to derive regression
559 relationships with atmospheric conditions and stability indices that can be forecast with NWP. Ensemble
560 forecasts of lightning using RF is a viable modelling approach for Alberta, Canada [Blouin et al., 2016].
561 Bates et al. [2017] used two machine learning methods (CART and RF) and three statistical methods to
562 classify wet and dry thunderstorms (lightning associated with dry thunderstorms are more likely to start
563 fires) in Australia.

564 4.2.3 Climate Change

565 Transfer modeling, whereby a model produced for one study region and/or distribution of environmental
566 conditions is applied to other cases [Phillips et al., 2006], is a common approach in climate change science.
567 Model transferability should be considered when using ML methods to estimated projected quantities due
568 to climate change or other environmental changes. With regards to climate change, transfer modeling is
569 essentially an extrapolation task. Previous studies in the context of species distribution modeling have
570 indicated ML approaches may be suitable for transfer modeling under future climate scenarios. For exam-
571 ple, Heikkinen et al. [2012] indicated MaxEnt and generalized boosting methods (GBM) have the better
572 transferability than either ANN and RF, and that the relatively poor transferability of RF may be due to
573 overfitting.

574 There are several publications on wildfires and climate change that use ML approaches. Amatulli
575 et al. [2013] found that Multivariate Adaptive Regression Splines (MARS) were better predictors of future
576 monthly area burned for 5 European countries as compared to Multiple Linear Regression and RF. [Parks
577 et al., 2016] projected fire severity for future time periods in Western USA using BRT. Young et al. [2017]
578 similarly used BRT to project future fire intervals in Alaska and found up to a fourfold increase in (30

year) fire occurrence probability by 2100. Several authors used MaxEnt to project future fire probability globally [Moritz et al., 2012], for Mediterranean ecosystems [Batllori et al., 2013], in Southwest China [Li et al., 2017], the pacific northwestern USA [Davis et al., 2017], and for south central USA [Stroh et al., 2018]. An alternative approach for projecting future potential burn probability was employed by Stralberg et al. [2018] who used RF to determine future vegetation distributions as inputs to ensemble Burn-P3 simulations. Another interesting paper of note was by Boulanger et al. [2018] who built a consensus model with 2 different predictor datasets and 5 different regression methods (generalised linear models, RF, BRT, CART and MARS) to make projections of future area burned in Canada. The consensus model can be used to quantify uncertainty in future area burned estimates. The authors noted that model uncertainty for future periods ($> 200\%$) can be higher than that of different climate models under different carbon forcing scenarios. This highlights the need for further work in the application of ML methods for projecting future fire danger under climate change.

4.3 Fire Occurrence, Susceptibility and Risk

Papers in this domain include prediction of fire occurrence and area burned (at a landscape or seasonal scales), mapping of fire susceptibility (or similar definitions of risk) and analysis of landscape or environmental controls on fire.

4.3.1 Fire occurrence prediction

Predictions of the number and location of fire starts in the upcoming day(s) are important to preparedness planning — that is, the acquisition of resources, including the relocation of mobile resources and readiness for expected fire activity. The origins of fire occurrence prediction (FOP) models go back almost 100 years [Nadeem et al., 2020]. FOP models typically use regression methods to relate the response variable (fire reports or hotspots) to weather, lightning, and other covariates for a geographic unit, or as a spatial probability. The seminal work of Brillinger and others in developing the spatio-temporal FOP framework is reviewed in Taylor et al. [2013] The most commonly used ML method in studies predicting fire occurrence were ANNs. As early as 1996, Vega-Garcia et al. [1996] used an ANN for human-caused wildfire prediction in Alberta, Canada, correctly predicting 85% of no-fire observations and 78% of fire observations. Not long after, Alonso-Betanzos et al. [2002] and Alonso-Betanzos et al. [2003] used ANN to predict a daily fire occurrence risk index using temperature, humidity, rainfall, and fire history, as part of a larger system for real-time wildfire management system in the Galicia region of Spain. Vasilakos et al. [2007] used separate ANNs for three different indices representing fire weather (Fire Weather Index; FWI), hazard (Fire Hazard Index; FHI), and risk (Fire Risk Index) to create a composite fire ignition index (FII) for estimating the probability of wildfire occurrence on the Greek island of Lesvos. Sakr et al. [2010] used meteorological variables in a SVM to create a daily fire risk index corresponding to the number of fires that could potentially occur on a particular day. Sakr et al. [2011] then compared the use of SVM and ANN for fire occurrence prediction based only on relative humidity and cumulative precipitation up to the specific day. While Sakr et al. [2011] reported low errors for the number of fires predicted by both the SVM and ANN models, ANN models outperformed SVM; however, the SVM performed better on binary classification of fire/no fire. It is important to note, however, that ANNs encompass a wide range of possible network architectures. In an Australian study, Dutta et al. [2013] compared the use of ten different types of ANN models for estimating monthly fire occurrence from climate data, and found that an Elman RNN performed the best.

After 2012, RF became the more popular method for predicting fire occurrence among the papers reviewed here. Stojanova et al. [2012] evaluated several machine learning methods for predicting fire outbreaks using geographical, remote sensed, and meteorological data in Slovenia, including single classifier methods (i.e., KNN, Naive Bayes, DT (using the J48 and jRIP algorithms), LR, SVM, and BN), and ensemble methods (AdaBoost, DT with bagging, and RF). The ensemble methods DT with bagging and RF displayed the best predictive performance with bagging having higher precision and RF having better

626 recall. Vecín-Arias et al. [2016] found that RF performed slightly better than LR for predicting lightning
627 fire occurrence in the Iberian Peninsula, based on topography, vegetation, meteorology, and lightning
628 characteristics. Similarly, Cao et al. [2017] found that a cost-sensitive RF analysis outperformed GLM
629 and ANN models for predicting wildfire ignition susceptibility. In recent non-comparative studies, Yu
630 et al. [2017] used RF to predict fire risk ratings in Cambodia using publicly available remote sensed
631 products, while Van Beusekom et al. [2018] used RF to predict fire occurrence in Puerto Rico and found
632 precipitation was found to be the most important predictor. The maximum entropy (MaxEnt) method
633 has also been used for fire occurrence prediction [De Angelis et al., 2015, Chen et al., 2015]. For example,
634 De Angelis et al. [2015] used MaxEnt to evaluate different meteorological variables and fire-indices (e.g.
635 the Canadian Fire Weather Index, FWI) for daily fire risk forecasting in the mountainous Canton Ticino
636 region of Switzerland. The authors of that study found that combinations of such variables increased
637 predictive power for identifying daily meteorological conditions for wildfires. Dutta et al. [2016] use a two-
638 stage machine learning approach (ensemble of unsupervised deep belief neural networks with conventional
639 supervised ensemble machine learning) to predict bush-fire hot spot incidence on a weekly time-scale. In
640 the first unsupervised deep learning phase, Dutta et al. [2016] used Deep Belief Networks (DBNet; an
641 ensemble deep learning method) to generate simple features from environmental and climatic surfaces.
642 In the second supervised ensemble classification stage, features extracted from the first stage were fed
643 as training inputs to ten ML classifiers (i.e., conventional supervised Binary Tree, Linear Discriminant
644 Analyser, Naïve Bayes, KNN, Bagging Tree, AdaBoost, Gentle Boosting Tree, Random Under-Sampling
645 Boosting Tree, Subspace Discriminant, and Subspace KNN) to establish the best classifier for bush fire
646 hotspot estimation. The authors found that bagging and the conventional KNN classifier were the two
647 best classifiers with 94.5% and 91.8% accuracy, respectively.

648 4.3.2 Landscape scale burned area prediction

649 The use of ML methods in studies of burned area prediction have only occurred relatively recently compared
650 to other wildfire domains, yet such studies have incorporated a variety of ML methods. For example, Cheng
651 and Wang [2008] used an RNN to forecast annual average area burned in Canada, while Archibald et al.
652 [2009] used RF to evaluate the relative importance of human and climatic drivers of burnt area in Southern
653 Africa. Arnold et al. [2014] used Hard Competitive Learning (HCL) to identify clusters of unique pre-fire
654 antecedent climate conditions in the interior western US which they then used to construct fire danger
655 models based on MaxEnt.

656 Mayr et al. [2018] evaluated five common statistical and ML methods for predicting burned area and
657 fire occurrence in Namibia, including GLM, Multivariate Adaptive Regression Splines (MARS), Regression
658 Trees from Recursive Partitioning (RPART), RF, and SVMs for Regression (SVR). The RF model
659 performed best for predicting burned area and fire occurrence; however, adjusted R^2 values were slightly
660 higher for RPART and SVR in both cases. Likewise, de Bem et al. [2018] compared the use of LR and
661 ANN for modelling burned area in Brazil. Both LR and ANN showed similar performance; however, the
662 ANN had better accuracy values when identifying non-burned areas, but displayed lower accuracy when
663 classifying burned areas.

664 4.3.3 Fire Susceptibility Mapping

665 A considerable number of references (71) used various ML algorithms to map wildfire susceptibility, cor-
666 responding to either the spatial probability or density of fire occurrence (or other measures of fire risk
667 such as burn severity) although other terms such as fire vulnerability and risk have also been used. The
668 general approach was to build a spatial fire susceptibility model using either remote sensed or agency
669 reported fire data with some combination of landscape, climate, structural and anthropogenic variables as
670 explanatory variables. In general, the various modeling approaches used either a presence only framework
671 (e.g., MaxEnt) or a presence/absence framework (e.g., BRT or RF).

Early attempts at fire susceptibility mapping used CART [Amatulli et al., 2006, Amatulli and Camia, 2007, Lozano et al., 2008]. Amatulli and Camia [2007] compared fire density maps in central Italy using CART and multivariate adaptive regression splines (MARS) and found while CART was more accurate that MARS led to smoother density model. More recent work has used ensemble based classifiers, such as RF and BRT, or ANNs (see table S.3.3 in supplementary material for a full list) Several of these papers also compared ML and non-ML methods for fire susceptibility mapping and in general found superior performance from the ML methods. Specifically, Adab [2017] mapped fire hazard in the Northeast of Iran, and found ANN performed better than binary logistic regression (BLR) with an AUC of 87% compared with 81% for BLR. Bisquert et al. [2012] found ANN outperformed logistic regression for mapping fire risk in the North-west of Spain. Goldarag et al. [2016] also compared ANN and linear regression for fire susceptibility mapping in Northern Iran and found ANN had much better accuracy (93.49%) than linear regression (65.76%). Guo et al. [2016b] and Guo et al. [2016a] compared RF and logistic regression for fire susceptibility mapping in China and found RF led to better performance. Oliveira et al. [2012] compared RF and LR for fire density mapping in Mediterranean Europe and found RF outperformed linear regression. De Vasconcelos et al. [2001] found ANN had better classification accuracy than logistic regression for ignition probability maps in parts of Portugal.

Referring to table 3 and section S.3.3 of the supplementary material a frequently used ML method for fire susceptibility mapping was Maximum Entropy (MaxEnt) which is extensively used in landscape ecology for species distribution modeling [Elith et al., 2011]. In particular, Vilar et al. [2016] found MaxEnt performed better than GLM for fire susceptibility mapping in central Spain with respect to sensitivity (i.e., true positive rate) and commission error (i.e., false positive rate), even though the AUC was lower. Of further note, Duane et al. [2015] partitioned their fire data into topography-driven, wind-driven and convection-driven fires in Catalonia and mapped the fire susceptibility for each fire type.

Other ML methods used for regional fire susceptibility mapping include Bayesian networks [Bashari et al., 2016, Dlamini, 2011] and novel hybrid methods such as Neuro-Fuzzy systems [Jaafari et al., 2019, Tien Bui et al., 2017]. Bashari et al. [2016] noted that Bayesian networks may be useful because it allows probabilities to be updated when new observations become available. SVM was also used by a number of authors as a benchmark for other ML methods [Ghorbanzadeh et al., 2019b, Gigović et al., 2019, Hong et al., 2018, Jaafari, 2019, Ngoc Thach et al., 2018, Rodrigues and De la Riva, 2014, Sachdeva et al., 2018, Tehrany et al., 2018, Tien Bui et al., 2017, van Breugel et al., 2016, Zhang et al., 2019] but as we discuss below, it did not perform as well as other methods to which it was being compared.

There were two applications of ML for mapping global fire susceptibility including Moritz et al. [2012] who used MaxEnt and Luo et al. [2013] who used RF. Both of these papers found that at a global scale, precipitation was one of the most important predictors of fire risk.

The majority of papers considered thus far used the entire study period (typically 4 or more years) to map fire susceptibility, therefore neglecting the temporal aspect of fire risk. However, a few authors have considered various temporal factors to map fire susceptibility. Martín et al. [2019] included seasonality and holidays as explanatory variables for fire probability in northeast Spain. Vacchiano et al. [2018] predicted fire susceptibility separately for the winter and summer seasons. Several papers produced maps of fire susceptibility in the Eastern US by month of year [Peters et al., 2013, Peters and Iverson, 2017]. Parisien et al. [2014] examined differences in annual fire susceptibility maps and a 31 year climatology for the USA, highlighting the role of climate variability as a driver of fire occurrence. In particular, they found FWI90 (the 90th percentile of the Canadian Fire Weather Index) was the dominant factor for annual fire risk but not for climatological fire risk. Cao et al. [2017] considered a 10 day resolution (corresponding to the available fire data) for fire risk mapping, which makes their approach similar to fire occurrence prediction.

In addition to fire susceptibility mapping, a few papers focused on other aspects of fire risk including mapping probability of burn severity classes [Holden et al., 2009, Parks et al., 2018, Tracy et al., 2018]. Parks et al. [2018] additionally considered the role of fuel treatments on fire probability which has obvious implications for fire management. Additionally Ghorbanzadeh et al. [2019a] combined fire susceptibility maps with vulnerability and infrastructure indicators to produce a fire hazard map.

722 A number of papers directly compared three or more ML (and sometimes non-ML) methods for fire
 723 susceptibility mapping. Here we highlight some of these papers, which elucidate the performance and
 724 advantages/disadvantages of various ML methods. [Cao et al. \[2017\]](#) found a cost-sensitive RF model
 725 outperformed a standard RF model, ANN as well as probit and logistic regression. [Ghorbanzadeh et al.](#)
 726 [\[2019b\]](#) compared ANN, SVM and RF and found the best performance with RF. [Gigović et al. \[2019\]](#)
 727 compared SVM and RF for fire susceptibility mapping in combination with Bayesian averaging to generate
 728 ensemble models. They found the ensemble model led to marginal improvement (AUC = 0.848) over SVM
 729 (AUC=0.834) and RF (AUC=0.844). For mapping both wildfire ignitions and potential natural vegetation
 730 in Ethiopia [van Breugel et al. \[2016\]](#) also considered ensemble models consisting of a weighted combination
 731 of ML methods (RF, SVM, BRT, MaxEnt, ANN, CART) and non-ML methods (GLM and MARS) and
 732 concluded the ensemble member performed best over a number of metrics. However, in this paper RF
 733 showed the best overall performance of all methods including the ensemble model.

734 [Jaafari et al. \[2018\]](#) compared 5 decision tree based classifiers for wildfire susceptibility mapping in Iran.
 735 Here, the Alternating Decision tree (ADT) classifier achieved the highest performance (accuracy 94.3%) in
 736 both training and validation sets. [Ngoc Thach et al. \[2018\]](#) compared SVM, RF and a Multilayer Perceptron
 737 (MLP) neural network for forest fire danger mapping in the region of Tjuan chau in Vietnam. They found
 738 the performance of all models were comparable although MLP had the highest AUC values. Interestingly
 739 [Pourtaghi et al. \[2016\]](#) found that a generalized additive model (GAM) outperformed RF and BRT for fire
 740 susceptibility mapping in the Golestan province in Iran. This was one of the few examples we found where
 741 a non-ML method outperformed ML methods. [Rodrigues and De la Riva \[2014\]](#) compared RF, BRT, SVM
 742 and logistic regression for fire susceptibility mapping and found RF led to the highest accuracy as well as
 743 the most parsimonious model. [Tehrany et al. \[2018\]](#) compared a LogitBoost ensemble-based decision tree
 744 (LEDT) algorithm with SVM, RF and Kernel logistic regression (KLR) for fire susceptibility mapping in
 745 Lao Cai region of Vietnam and found the best performance with LEDT, closely followed by RF. Finally,
 746 of particular note, [Zhang et al. \[2019\]](#) compared CNN, RF, SVM, ANN and KLR for fire susceptibility
 747 mapping in the Yunnan Province of China. This was the only application of deep learning we could find
 748 for fire susceptibility mapping. The authors found that CNN outperformed the other algorithms with
 749 overall accuracy of 87.92% compared with RF (84.36%), SVM (80.04%), MLP (78.47%), KLR (81.23%).
 750 They noted that the benefit of CNN is that it incorporates spatial correlations so that it can learn spatial
 751 features. However, the downside is that deep learning models are not as easily interpreted as other ML
 752 methods (such as RF and BRT).

753 4.3.4 Landscape controls on fire

754 Many of the ML methods used in fire susceptibility mapping have also been used to examine landscape
 755 controls – ie. the relative importance of weather, vegetation, topography, structural and anthropogenic
 756 variables – on fire activity, which may facilitate hypothesis formation and testing or model building. From
 757 table 3 the most commonly used methods in this section were MaxEnt, RF, BRT and ANN. These methods
 758 all allow for the determination of variable importance (i.e. the relative influence of predictor variables in a
 759 given model of a response variable). A commonly used method to ascertain variable importance is through
 760 the use of partial dependence plots [[Hastie et al., 2009](#)]. This method works by averaging over models
 761 that exclude the predictor variable of interest, with the resulting reduction in AUC (or other performance
 762 metrics) representing the marginal effect of the variable on the response. Partial dependence plots have the
 763 advantage of being able to be applied to a wide range of ML methods. A related method for determining
 764 variable importance, often used for RFs, is a permutation test which involves random permutation of each
 765 predictor variable [[Strobl et al., 2007](#)]. Another model-dependent approach used for ANN is the use of
 766 partial derivatives (of the activation functions of hidden and output nodes) as outlined by [Vasilakos et al.](#)
 767 [\[2009\]](#). It should be noted that while many other methods for model interpretation and variable dependence
 768 exist, a discussion of these methods is outside the scope of this paper.

769 In general, the drivers of fire occurrence or area burned varied greatly by the study area considered
 770 (including the size of area) and the methods used. Consistent with other work on “top down” and “bottom

up” drivers of fire activity, at large scales climate variables were often determined to be the main drivers of fire activity whereas at smaller scales anthropogenic or structural factors exerted a larger influence. Here we discuss some of the papers that highlight the diversity of results for different study areas and spatial scales (global, country, ecoregion, urban) but refer the reader to section S.3.4 of the supplementary material for a full listing of papers in this section. Note that many of the papers listed under section S.3.4 also belong to the fire susceptibility mapping section and have already been discussed there.

Aldersley et al. [2011] considered drivers of monthly area burned at global and regional scales using both regression trees and RF. They found climate factors (high temperature, moderate precipitation, and dry spells) were the most important drivers at the global scale, although at the regional scale the models exhibited higher variability due to the influence of anthropogenic factors. At a continental scale Mansuy et al. [2019] used MaxEnt to show that climate variables were the dominant controls (over landscape and human factors) on area burned for most ecoregions for both protected areas and outside these areas, although anthropogenic factors exerted a stronger influence in some regions such as the Tropical Wet Forests ecoregion. [Masrur et al., 2018] used RF to investigate controls on circumpolar arctic fire and found June surface temperature anomalies were the most important variable for determining the likelihood of wildfire occurrence on an annual scale. Chingono and Mbohwa [2015] used MaxEnt to model fire occurrences in Southern Africa where most fires are human-caused and found vegetation (i.e., dry mass productivity and NDVI) were the main drivers of biomass burning. Curt et al. [2015] used BRT to examine drivers of fire in New Caledonia. Interestingly, they found that human factors (such as distance to villages, cities or roads) were dominant influences for predicting fire ignitions whereas vegetation and weather factors were most important for area burned. Curt et al. [2016] modeled fire probabilities by different fire ignition causes (lightning, intentional, accidental, negligence professional and negligence personal) in Southeastern France. They found socioeconomic factors (eg. housing and road density) were the dominant factors for ignitions and area burned for human-caused fires. Fernandes et al. [2016] used BRT to examine large fires in Portugal and found high pyrodiversity (ie. spatial structure due to fire recurrence) and low landscape fuel connectivity were important drivers of area burned. Curt et al. [2016] modeled fire probabilities by different fire ignition causes (lightning, intentional, accidental, negligence professional and negligence personal) in Southeastern France. They found socioeconomic factors (eg. housing and road density) were the dominant factors for ignitions and area burned for human-caused fires. Leys et al. [2017] used RF to find the drivers that determine sedimentary charcoal counts in order to reconstruct grassfire history in the Great Plains, USA. Not surprisingly, they found fire regime characteristics (eg. area burned and fire frequency) were the most important variables and concluded that charcoal records can therefore be used to reconstruct fire histories. Li et al. [2009] used ANNs to show that wildfire probability was strongly influenced by population density in Japan, with a peak determined by the interplay of positive and negative effects of human presence. This relationship, however, becomes more complex when weather parameters and forest cover percentage are added to the model. Liu et al. [2013] used BRT to study factors influencing fire size in the Great Xingan Mountains in Northeastern China. Their method included a “moving window” resampling technique that allowed them to look at the relative influence of variables at different spatial scales. They showed that the most dominant factors influencing fire size were fuel and topography for small fires, but fire weather became the dominant factor for larger fires. For regions of high population density, anthropogenic or structural factors are often dominant for fire susceptibility. For example Molina et al. [2019] used MaxEnt to show distance to roads, settlements or powerlines were the dominant factors for fire occurrence probability in the Andalusia region in southern Spain. MaxEnt has also been used for estimating spatial fire probability under different scenarios such as future projections of housing development and private land conservation [Syphard et al., 2016]. One study in China using RF found mean spring temperature was the most important variable for fire occurrence whereas forest stock was most important for area burned [Ying et al., 2018].

Some authors examined controls on fire severity using high resolution data for a single large fire. For example, several authors used RF to examine controls on burn severity for the 2013 Rim fire in the Sierra Nevada [Lydersen et al., 2014, Kane et al., 2015, Lydersen et al., 2017]. At smaller spatial scales fire

821 weather was the most important variable for fire severity, whereas fuel treatments were most important
 822 at larger spatial scales [Lydersen et al., 2017]. A similar study by Harris and Taylor [2017] showed that
 823 previous fire severity was an important factor influencing fire severity for the Rim fire. For the 2005 Riba
 824 de Saelices fire, Viedma et al. [2015] looked at factors contributing to burn severity using a BRT model
 825 and found burning conditions (including fire weather variables) were more important compared than stand
 826 structure and topography. For burn severity these papers all used the Relativized differenced Normalized
 827 Burn Ratio (RdNBR) metric, derived from Landsat satellite images, which allowed spatial modeling at
 828 high resolutions (eg. 30m by 30m). In addition to the more commonly used ML methods one paper by
 829 Wu et al. [2015] used KNN to identify spatially homogeneous fire environment zones by clustering climate,
 830 vegetation, topography, and human activity related variables. They then used CART to examine variable
 831 importance for each of three fire environment zones in south-eastern China. For landscape controls on fire
 832 there were few studies comparing multiple ML methods. One such study by Nelson et al. [2017] compared
 833 CART, BRT and RF for classifying different fire size classes in British Columbia, Canada. For both central
 834 and periphery regions they found the best performing model was BRT followed by CART and RF. For
 835 example, in the central region BRT achieved a classification accuracy of 88% compared with 82.9% and
 836 49.6% for the CART and RF models respectively. It is not clear from the study why RF performed poorly,
 837 although it was noted that variable importance differs appreciably between the three models.

838 4.4 Fire Behavior Prediction

839 In general, fire behavior includes physical processes and characteristics at a variety of scales including
 840 combustion rate, flaming, smouldering residence time fuel consumption, flame height, and flame depth.
 841 However, the papers in this section deal mainly with larger scale processes and characteristics such as the
 842 prediction of fire spread rates, fire growth, burned area, and fire severity, conditional on the occurrence
 843 (ignition) of one, or more, wildfires. Here, our emphasis is on prognostic applications, in contrast to the
 844 *Fuels Characterization, Fire Detection and Mapping* problem domain, in which we focused on diagnostic
 845 applications.

846 4.4.1 Fire spread and growth

847 Predicting the spread of a wildland fire is an important task for fire management agencies, particularly to
 848 aid in the deployment of suppression resources or to anticipate evacuations one or more days in advance.
 849 Thus, a large number of models have been developed using different approaches. In a series of reviews
 850 Sullivan [2009a,b,c] described fire spread models he classified as being of physical or quasi-physical nature,
 851 or empirical or quasi-empirical nature, as well as mathematical analogues and simulation models. Many
 852 fire growth simulation models convert one dimensional empirical or quasi-empirical spread rate models to
 853 two dimensions and then propagate a fire perimeter across a modelled landscape.

854 A wide range of ML methods have been applied to predict fire growth. For example, Markuzon and
 855 Kolitz [2009] tested several classifiers (RF, BNs, and KNN) to estimate if a fire would become large either
 856 one or two days following its observation; they found each of the tested methods performed similarly with
 857 RF correctly classifying large fires at a rate over 75%, albeit with a number of false positives. Vakalis
 858 et al. [2004] used a ANN in combination with a fuzzy logic model to estimate the rate of spread in the
 859 mountainous region of Attica in Greece. A number of papers used genetic algorithms (GAs) to optimize
 860 input parameters to a physics or empirically based fire simulator in order to improve fire spread predictions
 861 [Abdalhaq et al., 2005, Rodriguez et al., 2008, Rodriguez et al., 2009, Artés et al., 2014, 2016, Carrillo
 862 et al., 2016, Denham et al., 2012, Cencerrado et al., 2012, 2013, 2014, Artés et al., 2017, Denham and
 863 Laneri, 2018]. For example, Cencerrado et al. [2014] developed a framework based on GAs to shorten the
 864 time needed to run deterministic fire spread simulations. They tested the framework using the FARSITE
 865 [Finney, 2004] fire spread simulator with different input scenarios sampled from distributions of vegetation
 866 models, wind speed/direction, and dead/live fuel moisture content. The algorithm used a fitness function
 867 which discarded the most time-intensive simulations, but did not lead to an appreciable decrease in the

868 accuracy of the simulations. Such an approach is potentially useful for fire management where it is desirable
 869 to predict fire behavior as far in advance as possible so that the information can be enacted upon. This
 870 approach may greatly reduce overall simulation time by reducing the input parameter space as also noted
 871 by Artés et al. [2016] and Denham et al. [2012], or through parallelization of simulation runs for stochastic
 872 approaches [Artés et al., 2017, Denham and Laneri, 2018]. A different goal was considered by Ascoli et al.
 873 [2015] who used a GA to optimize fuel models in Southern Europe by calibrating the model with respect
 874 to rate of spread observations.

875 Kozik et al. [2013] presented a fire spread model that used a novel ANN implementation that incorpo-
 876 rated a Kalman filter for data assimilation that could potentially be run in real-time, the resulting model
 877 more closely resembling that of complex cellular automata than a traditional ANN. The same authors later
 878 implemented this model and simulated fire growth under various scenarios with different wind speeds and
 879 directions, or both, although a direct comparison with real fire data was not possible [Kozik et al., 2014].

880 Zheng et al. [2017] simulated fire spread by integrating a cellular automata (CA) model with an Extreme
 881 Learning Machine (ELM; a type of feedforward ANN). Transition rules for the CA were determined by
 882 the ELM trained with data from historical fires, as well as vegetation, topographic, and meteorological
 883 data. Likewise, Chetehouna et al. [2015] used ANNs to predict fire behavior, including rate of spread,
 884 and flame height and angle. In contrast, Subramanian and Crowley [2017] formulated the problem of fire
 885 spread prediction as a Markov Decision Process, where they proposed solutions based on both a classic
 886 reinforcement learning algorithm and a deep reinforcement learning algorithm – the authors found the
 887 deep learning approach improved on the traditional approach when tested on two large fires in Alberta,
 888 Canada. The authors further developed this work to compare five widely used reinforcement learning
 889 algorithms [Subramanian and Crowley, 2018], and found that the Asynchronous Advantage Actor-Critic
 890 (A3C) and Monte Carlo Tree Search (MCTS) algorithms achieved the best accuracy. Meanwhile, Khakzad
 891 [2019] developed a fire spread model to predict the risk of fire spread in Wildland-Industrial Interfaces,
 892 using Dynamic Bayesian Networks (DBN) in combination with a deterministic fire spread model. The
 893 Canadian Fire Behavior Prediction (FBP) system, which uses meteorological and fuel conditions data as
 894 inputs, determined the fire spread probabilities from one node to another in the aforementioned DBN.

895 More recently Hodges and Lattimer [2019] trained a (deep learning) CNN to predict fire spread using
 896 environmental variables (topography, weather and fuel related variables). Outputs of the CNN were spatial
 897 grids corresponding to the probability the burn map reached a pixel and the probability the burn map
 898 did not reach a pixel. Their method achieved a mean precision of 89% and mean sensitivity of 80% with
 899 reference 6 hourly burn maps computed using the physics-based FARSITE simulator. Radke et al. [2019]
 900 also used a similar approach to predict daily fire spread for the 2016 Beaver Creek fire in Colorado.

901 4.4.2 Burned area and fire severity prediction

902 There are a number of papers that focus on using ML approaches to directly predict the final area burned
 903 from a wildfire. Cortez and Morais [2007] compared multiple regression and four different ML methods
 904 (DT, RF, ANN, and SVM) to predict area burned using fire and weather (i.e., temperature, precipitation,
 905 relative humidity and wind speed) data from the Montesinho natural park in northeastern Portugal, and
 906 found that SVM displayed the best performance. A number of publications subsequently used the data
 907 from Cortez and Morais [2007] to predict area burned using various ML methods, including ANN [Safi and
 908 Bouroumi, 2013, Storer and Green, 2016], genetic algorithms [Castelli et al., 2015], both ANN and SVM
 909 [Al-Janabi et al., 2018], and decision trees [Alberg, 2015, Li et al., 2018a]. Notably, Castelli et al. [2015]
 910 found that a GA variant outperformed other ML methods including SVM. Xie and Shi [2014] used a similar
 911 set of input variables with SVM to predict burned area in for Guangzhou City in China. In addition to
 912 these studies, Toujani et al. [2018] used hidden Markov models (HMM) to predict burned area in the north-
 913 west of Tunisia, where the spatiotemporal factors used as inputs to the model were initially clustered using
 914 self-organizing maps (SOMs). Liang et al. [2019] compared back-propagation neural networks, recurrent
 915 neural networks (RNN) and Long Short Term Memory (LSTM) neural networks to predict wildfire scale,
 916 a quantity related to area burned and fire duration, in Alberta Canada. They found the highest accuracy

917 (90.9%) was achieved with LSTM.

918 Most recently, [Xie and Peng \[2019\]](#) compared a number of machine learning methods for estimating area
919 burned (regression) and binary classification of fire sizes (> 5 Ha) in Montesinho natural park, Portugal.
920 For the regression task, they found a tuned RF algorithm performed better than standard RF, tuned
921 and standard gradient boosted machines, tuned and standard generalized linear models (GLMs) and deep
922 learning. For the classification problem they found extreme gradient boosting and deep learning had a
923 higher accuracy than CART, RF, SVM, ANN, and logistic regression.

924 By attempting to predict membership of burned area size classes, a number of papers were able to
925 recast the problem of burned area prediction as a classification problem. For example, [Yu et al. \[2011\]](#)
926 used a combination of SOMs and back-propagation ANNs to classify forest fires into size categories based
927 on meteorological variables. This approach gave [Yu et al. \[2011\]](#) better accuracy (90%) when compared
928 with a rules-based method (82%). [Özbayoğlu and Bozer \[2012\]](#) estimated burned area size classes using
929 geographical and meteorological data using three different machine learning methods: i) Multilayer
930 Perceptron (MLP); ii) Radial Basis Function Networks (RBFN); and iii) SVM. Overall, the best perform-
931 ing method was MLP, which achieved a 65% success rate, using humidity and windspeed as predictors.
932 [Zwirgmaier et al. \[2013\]](#) used a BN to predict area burned classes using historical fire data, fire weather
933 data, fire behaviour indices, land cover, and topographic data. [Shidik and Mustofa \[2014\]](#) used a hybrid
934 model (Fuzzy C-Means and Back-Propagation ANN) to estimate fire size classes using data from [Cortez
935 and Morais \[2007\]](#), where the hybrid model performed best with an accuracy of 97.50% when compared
936 with Naive Bayes (55.5%), DT (86.5%), RF (73.1%), KNN (85.5%) and SVM (90.3%). [Mitsopoulos and
937 Mallinis \[2017\]](#) compared BRT, RF and Logistic Regression to predict 3 burned area classes for fires in
938 Greece. They found RF led to the best performance of the three tested methods and that fire suppression
939 and weather were the two most important explanatory variables. [Coffield et al. \[2019\]](#) compared CART,
940 RF, ANN, KNN and gradient boosting to predict 3 burned area classes at time of ignition in Alaska.
941 They found a parsimonious model using CART with Vapor Pressure Deficit (VPD) provided the best
942 performance of the models and variables considered.

943 We found only one study that used ML to predict fire behavior related to fire severity, which is important
944 in the context of fire ecology, suggesting that there are opportunities to apply ML in this domain of wildfire
945 science. In that paper, [Zald and Dunn \[2018\]](#) used RF to determine that the most important predictor of
946 fire severity was daily fire weather, followed by stand age and ownership, with less predictability given by
947 topographic features.

948 4.5 Fire Effects

949 Fire Effects prediction studies have largely used regression based approaches to relate costs, losses, or other
950 impacts (e.g., soils, post-fire ecology, wildlife, socioeconomic factors) to physical measures of fire severity
951 and exposure. Importantly, this category also includes wildfire smoke and particulate modelling (but not
952 smoke detection which was previously discussed in the fire detection section).

953 4.5.1 Soil Erosion and Deposits

954 [Mallinis et al. \[2009\]](#) modelled potential post-fire soil erosion risk following a large intensive wildfire in the
955 Mediterranean area using CART and k-means algorithms. In that paper, before wildfire, 55% of the study
956 area was classified as having severe or heavy erosion potential, compared to 90% post-fire, with an overall
957 classification accuracy of 86%. Meanwhile, [Buckland et al. \[2019\]](#) used ANNs to examine the relationships
958 between sand deposition in semi-arid grasslands and wildfire occurrence, land use, and climatic conditions.
959 The authors then predicted soil erosion levels in the future given climate change assumptions.

960 4.5.2 Smoke and Particulate Levels

961 Smoke emitted from wildfires can seriously lower air quality with adverse effects on the health of both
 962 human and non-human animals, as well as other impacts. Thus, it is not surprising that ML methods
 963 have been used to understand the dynamics of smoke from wildland fire. For example, Yao et al. [2018b]
 964 used RF to predict the minimum height of forest fire smoke using data from the CALIPSO satellite. More
 965 commonly, ML methods have also been used to estimate population exposure to fine particulate matter
 966 (e.g., PM_{2.5}: atmospheric particulate matter with diameter less than 2.5 μ m), which can be useful for
 967 epidemiological studies and for informing public health actions. One such study by Yao et al. [2018a]
 968 also used RF to estimate hourly concentrations of PM_{2.5} in British Columbia, Canada. Zou et al. [2019]
 969 compared RF, BRT and MLR to estimate regional PM_{2.5} concentrations in the Pacific Northwest and
 970 found RF performed much better than the other algorithms. In another very broad study covering several
 971 datasets and ML methods, Reid et al. [2015] estimated spatial distributions of PM_{2.5} concentrations
 972 during the 2008 northern California wildfires. The authors of the aforementioned study used 29 predictor
 973 variables and compared 11 different statistical models, including RF, BRT, SVM, and KNN. Overall, the
 974 BRT and RF models displayed the best performance. Emissions other than particulate matter have also
 975 been modelled using ML, as Lozhkin et al. [2016] used an ANN to predict carbon monoxide concentrations
 976 emitted from a peat fire in Siberia, Russia. In another study, the authors used ten different statistical and
 977 ML methods and 21 covariates (including weather, geography, land-use, and atmospheric chemistry) to
 978 predict ozone exposures before and after wildfire events [Watson et al., 2019]. Here, gradient boosting gave
 979 the best results with respect to both root mean square error and R^2 values, followed by RF and SVM. In a
 980 different application related to smoke, Fuentes et al. [2019] used ANNs to detect smoke in several different
 981 grape varieties used for wine making.

982 4.5.3 Post-fire regeneration, succession, and ecology

983 The study of post-fire regeneration is an important aspect of understanding forest and ecosystem responses
 984 and resilience to wildfire disturbances, with important ecological and economic consequences. RF, for
 985 example, has been a popular ML method for understanding the important variables driving post-fire
 986 regeneration [João et al., 2018, Vijayakumar et al., 2016]. Burn severity (a measure of above and below
 987 ground biomass loss due to fire) is an important metric for understanding the impacts of wildfire on
 988 vegetation and post-fire regeneration, soils, and potential successional shifts in forest composition, and as
 989 such, has been included in many ML studies in this section, including [Barrett et al., 2011, Cai et al., 2013,
 990 Cardil et al., 2019, Chapin et al., 2014, Divya and Vijayalakshmi, 2016, Fairman et al., 2017, Han et al.,
 991 2015, Johnstone et al., 2010, Liu and Yang, 2014, Martín-Alcón and Coll, 2016, Sherrill and Romme, 2012,
 992 Thompson and Spies, 2010]. For instance, Cardil et al. [2019] used BRT to demonstrate that remotely-
 993 sensed data (i.e., Relative Differenced Normalized Burn Ratio index; RdNBR) can provide an acceptable
 994 assessment of fire-induced impacts (i.e., burn severity) on forest vegetation, while [Fairman et al., 2017]
 995 used RF to identify the variables most important in explaining plot-level mortality and regeneration of
 996 *Eucalyptus pauciflora* in Victoria, Australia, affected by high-severity wildfires and subsequent re-burns.
 997 Debouk et al. [2013] assessed post-fire vegetation regeneration status using field measurements, a canopy
 998 height model, and Lidar (i.e., 3D laser scanning) data with a simple ANN. Post-fire regeneration also has
 999 important implications for the successional trajectories of forested areas, and a few studies have examined
 1000 this using ML approaches [Barrett et al., 2011, Cai et al., 2013, Johnstone et al., 2010]. For example,
 1001 Barrett et al. [2011] used RF to model fire severity, from which they made an assessment of the area
 1002 susceptible to a shift from coniferous to deciduous forest cover in the Alaskan boreal forest, while Cai et al.
 1003 [2013] used BRT to assess the influence of environmental variables and burn severity on the composition
 1004 and density of post-fire tree recruitment, and thus the trajectory of succession, in northeastern China. In
 1005 other studies not directly related to post-fire regeneration, Hermosilla et al. [2015] used RF to attribute
 1006 annual forest change to one of four categories, including wildfire, in Saskatchewan, Canada, while [Jung
 1007 et al., 2013] used GA and RF to estimate the basal area of post-fire residual spruce (*Picea obovate*) and fir

1008 (*Abies sibirica*) stands in central Siberia using remotely sensed data. Magadzire et al. [2019] used MaxEnt
1009 to demonstrate that fire return interval and species life history traits affected the distribution of plant
1010 species in South Africa. ML has also been used to examine fire effects on the hydrological cycle, as Poon
1011 et al. [2018] used SVM to estimate both pre- and post-wildfire evapotranspiration using remotely sensed
1012 variables.

1013 Considering the potential impacts of wildfires on wildlife, it is perhaps surprising that relatively few of
1014 such studies have adopted ML approaches. However, ML methods have been used to predict the impacts
1015 of wildfire and other drivers on species distributions and arthropod communities. Hradsky et al. [2017], for
1016 example, used non-parametric BNs to describe and quantify the drivers of faunal distributions in wildfire-
1017 affected landscapes in southeastern Australia. Similarly, Reside et al. [2012] used MaxEnt to model bird
1018 species distributions in response to fire regime shifts in northern Australia, which is an important aspect
1019 of conservation planning in the region. ML has also been used to look at the effects of wildfire on fauna at
1020 the community level, as Luo et al. [2017] used DTs, Association Rule Mining, and AdaBoost to examine
1021 the effects of fire disturbance on spider communities in Cangshan Mountain, China.

1022 4.5.4 Socioeconomic effects

1023 ML methods have been little used to model socio-economic impacts of fire to date. We found one study
1024 in which BNs were used to predict the economic impacts of wildfires in Greece from 2006-2010 due to
1025 housing losses [Papakosta et al., 2017]. The authors did this by first defining a causal relationship between
1026 the participating variables, and then using BNs to estimate housing damages. It is worth noting that the
1027 problem of detecting these causal relationships from data is a difficult task and remains an active area of
1028 research in artificial intelligence.

1029 4.6 Fire management

1030 The goal of contemporary fire management is to have the appropriate amount of fire on the landscape, which
1031 may be accomplished through the management of vegetation including prescribed burning, the management
1032 of human activities (prevention), and fire suppression. Fire management is a form of risk management that
1033 seeks to maximize fire benefits and minimize costs and losses [Finney, 2005]. Fire management decisions
1034 have a wide range of scales, including long-term strategic decisions about the acquisition and location of
1035 resources or the application of vegetation management in large regions, medium-term tactical decisions
1036 about the acquisition of additional resources, relocation, or release of resources during the fire season, and
1037 short-term real time operational decisions about the deployment and utilization of resources on individual
1038 incidents. Fire preparedness and response is a supply chain with a hierarchical dependence. Taylor [2020]
1039 describes 20 common decision types in fire management and maps the spatial-temporal dimensions of their
1040 decision spaces.

1041 Fire management models can be predictive, such as the probability of initial attack success, or pre-
1042 scriptive such as to maximize/minimize an objective function (e.g., optimal helicopter routing to minimize
1043 travel time in crew deployment). While advances have been made in the domain of wildfire management
1044 using ML techniques, there have been relatively few studies in this area compared to other wildfire problem
1045 domains. Thus, there appears to be great potential for ML to be applied to wildfire management problems,
1046 which may lead to novel and innovative approaches in the future.

1047 4.6.1 Planning and policy

1048 An important area of fire management is planning and policy, where various ML methods have been
1049 applied to address pertinent challenges. For example, Bao et al. [2015] used GA, which are useful for
1050 solving multi-objective optimization problems, to optimize watchtower locations for forest fire monitoring.
1051 Bradley et al. [2016] used RF to investigate the relationship between the protected status of forest in the
1052 western US and burn severity. Likewise, Ruffault and Mouillot [2015] also used BRTs to assess the impact

of fire policy introduced in the 1980s on fire activity in southern France and the relationships between fire and weather, and [Penman et al. \[2011\]](#) used BNs to build a framework to simultaneously assess the relative merits of multiple management strategies in Wollemi National Park, NSW, Australia. [McGregor et al. \[2016\]](#) used Markov decision processes (MDP) and model free Monte Carlo method to create fast running simulations (based on the FARSITE simulator) to create interactive visualizations of forest futures over 100 years based on alternate high-level suppression policies. [McGregor et al. \[2017\]](#) demonstrated ways in which a variety of ML and optimization methods can be used to create an interactive approximate simulation tool for fire managers. The authors of the aforementioned study utilized a modified version of the FARSITE fire-spread simulator, which was augmented to run thousands of simulation trajectories while also including new models of lightning strike occurrences, fire duration, and a forest vegetation simulator. [McGregor et al. \[2017\]](#) also clearly show how decision trees can be used to analyze a hierarchy of decision thresholds for deciding whether to suppress a fire or not; their hierarchy splits on fuel levels, then intensity estimations, and finally weather predictors to arrive at a generalizable policy.

4.6.2 Fuel treatment

ML methods have also been used to model the effects of fuel treatments in order to mitigate wildfire risk. For example, [Penman et al. \[2014\]](#) used a BN to examine the relative risk reduction of using prescribed burns on the landscape versus within the 500m interface zone adjacent to houses in the Sydney basin, Australia. [Lauer et al. \[2017\]](#) used approximate dynamic programming (also known as reinforcement learning) to determine the optimal timing and location of fuel treatments and timber harvest for a fire-threatened landscape in Oregon, USA, with the objective of maximizing wealth through timber management. Similarly, [Arca et al. \[2015\]](#) used GA for multi-objective optimization of fuel treatments.

4.6.3 Wildfire preparedness and response

Wildfire preparedness and response issues have also been examined using ML techniques. [Costafreda-Aumedes et al. \[2015\]](#) used ANNs to model the relationships between daily fire load, fire duration, fire type, fire size, and response time, as well as personnel and terrestrial/aerial units deployed for individual wildfires in Spain. Most of the models in [Costafreda-Aumedes et al. \[2015\]](#) highlighted the positive correlation of burned area and fire duration with the number of resources assigned to each fire, and some highlighted the negative influence of daily fire load. In another study, [Penman et al. \[2015\]](#) used Bayesian Networks to assess the relative influence of preventative and suppression management strategies on the probability of house loss in the Sydney basin, Australia. [O'Connor et al. \[2017\]](#) used BRT to develop a predictive model of fire control locations in the Northern Rocky Mountains, USA, based on the likelihood of final fire perimeters, while [Homchaudhuri et al. \[2010\]](#) used GAs to optimize fireline generation. [Rodrigues et al. \[2019\]](#) modelled the probability that wildfire will escape initial attack using a RF model trained with fire location, detection time, arrival time, weather, fuel types, and available resources data. Important variables in [Rodrigues et al. \[2019\]](#) included fire weather and simultaneity of events. [Julian and Kochenderfer \[2018a\]](#) used two different RL algorithms to develop a system for autonomous control of one or more aircraft in order to monitor active wildfires.

4.6.4 Social factors

Recently, the use of ML in fire management has grown to encompass more novel aspects of fire management, even including the investigation of criminal motives related to arson, as [Delgado et al. \[2018\]](#) used BNs to characterize wildfire arsonists in Spain thereby identifying five motivational archetypes (i.e., slight negligence; gross negligence; impulsive; profit; and revenge).

5 Discussion

ML methods have seen a spectacular evolution in development, accuracy, computational efficiency, and application in many fields since the 1990s. It is therefore not surprising that ML has been helpful in providing new insights into several critical sustainability and social challenges in the 21st century [Gomes, 2009, Sullivan et al., 2014, Butler, 2017]. The recent uptake and success of ML methods has been driven in large part by ongoing advances in computational power and technology. For example, the recent use of bandwidth optimized Graphics Processing Units (GPUs) takes advantage of parallel processing for simultaneous execution of computationally expensive tasks, which has facilitated a wider use of computationally demanding but more accurate methods like DNNs. The advantages of powerful but efficient ML methods are therefore widely anticipated as being useful in wildfire science and management.

However, despite some early papers suggesting that data driven techniques would be useful in forest fire management [Latham, 1987, Kourtz, 1990, 1993], our review has shown that there was relatively slow adoption of ML-based research in wildfire science up to the 2000s compared with other fields, followed by a sharp increase in publication rate in the last decade. In the early 2000s, data mining techniques were quite popular and classic ML methods such as DTs, RF, and bagging and boosting techniques began to appear in the wildfire science literature (e.g., Stojanova et al. [2006]). In fact, some researchers started using simple feed forward ANNs for small scale applications as early as the mid 1990s and early 2000s (e.g., McCormick et al. [1999], Al-Rawi et al. [2002]). In the last three decades, almost all major ML methods have been used in some way in wildfire applications, although some more computationally demanding methods, such as SOMs and cellular automata, have only been actively experimented with in the last decade [Toujani et al., 2018, Zheng et al., 2017]. Furthermore, the recent development of DL algorithms, with a particular focus on extracting spatial features from images, has led to a sharp rise in the application of DL for wildfire applications in the last decade. It is evident, however, from our review that while an increasing number of ML methodologies have been used across a variety of fire research domains over the past 30 years, this research is unevenly distributed among ML algorithms, research domains and tasks, and has had limited application in fire management.

Many fire science and management questions can be framed within a fire risk context. Xi et al. [2019] discussed the advantages of adopting a risk framework with regard to statistical modeling of wildfires. There the risk components of “hazard”, “vulnerability” and “exposure” are replaced respectively by fire probability, fire behavior and fire effects. Most fire management activities can be framed as risk controls to mitigate these components of risk. Traditionally, methods used in wildfire fire science to address these various questions have included physical modeling (e.g., Sullivan [2009a,b,c]), statistical methods (e.g., Taylor et al. [2013], Xi et al. [2019]), simulation modeling (e.g., Keane et al. [2004]), and operations research methods (Martell [2015], Minas et al. [2012]).

In simple terms, any analytical study begins with one or more of four questions: “what happened?”; “why did it happen?”; “what will happen?”; or “what to do?” Corresponding data driven approaches to address these questions are respectively called descriptive, diagnostic, predictive, and prescriptive analytics. The type of analytical approach adopted then circumscribes the types of methodological approaches (e.g., regression, classification, clustering, dimensionality reduction, decision making) and sets of possible algorithms appropriate to the analysis.

In our review, we found that studies incorporating ML methods in wildland fire science were predominantly associated with descriptive or diagnostic analytics, reflecting the large body of work on fire detection and mapping using classification methods, and on fire susceptibility mapping and landscape controls on fire using regression approaches. In many cases, the ML methods identified in our review are an alternative to statistical methods used for clustering and regression. While the aforementioned tasks are undoubtedly very important for understanding wildland fire, we found much less work associated with predictive or prescriptive analytics, such as fire occurrence prediction (predictive), fire behaviour prediction (predictive), and fire management (prescriptive). This may be because: a) particular domain knowledge is required to frame fire management problems; b) fire management data are often not publicly available, need a lot of

work to transform into an easily analyzable form, or do not exist at the scale of the problem; and c) some fire management problems are not suited or can't be fully addressed by ML approaches. We note that much of the work on fire risk in the fire susceptibility and mapping domain used historical fire and environmental data to map fire susceptibility; therefore, while that work aims to inform future fire risk, it cannot be considered to be predictive analytics, except, for example, in cases where it was used in combination with climate change projections. It appears then that, in general, wildfire science research is currently more closely aligned with descriptive and diagnostic analytics, whereas wildfire management goals are aligned with predictive and prescriptive analytics. This fundamental difference identifies new opportunities for research in fire management, which we discuss later in this paper.

In the remainder of the paper, we examine some considerations for the use of ML methods, including: data considerations, model selection and accuracy, implementation challenges, interpretation, opportunities, and implications for fire management.

5.1 Data considerations

ML is a data-centric modeling paradigm concerned with finding patterns in data. Importantly, data scientists need to determine, often in collaboration with fire managers or domain experts, whether there are suitable and sufficient data for a given modeling task. Some of the criteria for suitable data include whether: a) the predictands and covariates are or can be wrangled into the same temporal and spatial scale; b) the observations are a representative sample of the full range of conditions that may occur in application of a model to future observations; and c) whether the data are at spatiotemporal scale appropriate to the fire science or management question. The first of these criteria can be relaxed in some ML models such as ANNs and DNNs, where inputs and outputs can be at different spatial or temporal scales for appropriately designed network architectures, although data normalization may still be required. The second criterion also addresses the important question of whether enough data exists for training a given algorithm for a given problem. In general, this question depends on the nature of the problem, complexity of the underlying model, data uncertainty and many other factors (see Roh et al. [2018] for a further discussion of data requirements for ML). In any case, many complex problems require a substantive data wrangling effort, to acquire, perform quality assurance, and fuse data into sampling units at the appropriate spatiotemporal scale. An example of this in daily fire occurrence prediction, where observations of a variety of features (e.g., continuous measures such as fire arrival time and location, or lightning strike times and locations) are discretized into three-dimensional (e.g., longitude, latitude, and day) cells called voxels. Another important consideration for the collection and use of data in machine learning is selection bias. A form of spatial selection bias called preferential sampling occurs when sampling occurs preferentially in locations where one expects a certain response [Diggle et al., 2010]. For example, preferential sampling may occur in air monitoring, because sensors may be placed in locations where poor air quality is expected [Shaddick and Zidek, 2014]. In general, preferential sampling or other selection biases may be avoided altogether by selecting an appropriate sampling strategy at the experimental design phase, or, where this is not possible, to take it into account in model evaluation [Zadrozny, 2004].

For the problem domain fire detection and mapping, most applications of ML used some form of imagery (e.g., remote sensed satellite images or terrestrial photographs). In particular, many papers used satellite data (e.g., Landsat, MODIS) to determine vegetation differences before and after a fire and so were able to map area burned. For fire detection, many applications considered either remote sensed data for hotspot or smoke detection, or photographs of wildfires (used as inputs to an image classification problem). For fire weather and climate change, the three main sources of data were either weather station observations, climate reanalyses (modelled data that include historical observations), or GCMs for future climate projections. Reanalyses and GCMs are typically highly dimensional large gridded spatiotemporal datasets which require careful feature selection and/or dimensional reduction for ML applications. Fire occurrence prediction, susceptibility, and risk applications used a large number of different environmental variables as predictors, but almost all used fire locations and associated temporal information as predictands. Fire data itself is usually collated from fire management agencies in the form of georeferenced points or perimeter

1193 data, along with reported dates, ignition cause, and other related variables. Care should be taken using
1194 such data because changes in reporting standards or accuracy may lead to data inhomogeneity. As well as
1195 fire locations and perimeters, fire severity is an attribute of much interest to fire scientists. Fire severity is
1196 often determined from remotely sensed data and represented using variables such as the Differenced Nor-
1197 malized Burn Ratio (dNBR) and variants, or through field sampling. However, remote sensed estimates of
1198 burn severity should be considered as proxies as they have low skill in some ecosystems. Other fire ecology
1199 research historically relies on in situ field, sampling although many of the ML applications attempt to
1200 resolve features of interest using remote sensed data. Smoke data can also be derived from remote sensed
1201 imagery or from air quality sensors (e.g., PM_{2.5}, atmospheric particulate matter less than 2.5 μm).

1202 Continued advances in remote sensing, as well as the quality and availability of remote sensed data prod-
1203 ucts, in weather and climate modeling have led to increased availability of large spatiotemporal datasets,
1204 which presents both an opportunity and challenge for the application of ML methods in wildfire research
1205 and management. The era of “big data” has seen the development of cloud computing platforms to provide
1206 the computing and data storage facilities to deal with these large datasets. For example, in our review we
1207 found two papers [Crowley et al., 2019, Quintero et al., 2019] that used Google Earth Engine which inte-
1208 grates geospatial datasets with a coding environment [Gorelick et al., 2017]. In any case, data processing
1209 and management plays an important role in the use of large geospatial datasets.

1210 5.2 Model selection and accuracy

1211 Given a wildfire science question or management problem and available relevant data, a critical question to
1212 ask is what is the most appropriate modeling tool to address the problem? Is it a standard statistical model
1213 (e.g., linear regression or LR), a physical model (e.g., FIRETEC or other fire simulator), a ML model, or a
1214 combination of approaches? Moreover, which specific algorithm will yield the most accurate classification
1215 or regression. Given the heterogeneity of research questions, study areas, and datasets considered in the
1216 papers reviewed here, it is not possible to comprehensively answer these questions with respect to ML
1217 approaches. Even in the case where multiple studies used the same dataset [Cortez and Morais, 2007,
1218 Safi and Bouroumi, 2013, Storer and Green, 2016, Castelli et al., 2015, ALJanabi et al., 2018, Alberg,
1219 2015, Li et al., 2018a, Castelli et al., 2015] the different research questions considered meant a direct
1220 comparison of ML methods was not possible between research studies. However, a number of individual
1221 studies did make comparisons between multiple ML methods, or between ML and statistical methods for
1222 a given wildfire modeling problem and dataset. Here we highlight some of their findings to provide some
1223 guidance with respect to model selection. In our review (see section 4 and the supplementary material), we
1224 found 29 papers comparing ML and statistical methods, where in the majority of these cases ML methods
1225 were found to be more accurate than traditional statistical methods (e.g., GLMs), or displayed similar
1226 performance [Pu and Gong, 2004, Bates et al., 2017, de Bem et al., 2018]. In only one study on climate
1227 change by Amatulli et al. [2013], MARS was found to be superior to RF for their analytical task. A sizable
1228 number of the comparative studies (14) involved classification problems that used LR as a benchmark
1229 method against ANN or ensemble tree methods. For studies comparing multiple ML methods, there was
1230 considerable variation in the choice of most accurate method; however, in general ensemble methods tended
1231 to outperform single classifier methods (e.g., Stojanova et al. [2012], Dutta et al. [2016], Mayr et al. [2018],
1232 Nelson et al. [2017], Reid et al. [2015], Watson et al. [2019]), except in one case where the most accurate
1233 model (CART) was also the most parsimonious [Coffield et al., 2019]. A few more recent papers also
1234 highlighted the advantages of DL over other methods. In particular, for fire detection, Zhang et al. [2018b]
1235 compared CNNs with SVM and found that CNNs were more accurate, while Zhao et al. [2018] similarly
1236 found CNNs superior to SVMs and ANNs. For fire susceptibility mapping, Zhang et al. [2019] found CNNs
1237 were more accurate than RF, SVMs, and ANNs. For time series forecasting problems, Liang et al. [2019]
1238 found LSTMs outperformed ANNs. Finally, Cao et al. [2019] found that using an LSTM combined with a
1239 CNN led to better fire detection performance from video compared with CNNs alone.

1240 In any case, more rigorous inter-model comparisons are needed to reveal in which conditions, and in
1241 what sense particular methods are more accurate, as well as to establish procedures for evaluating accuracy.

ML methods are also prone to overfitting, so it is important to evaluate models with robust test datasets using appropriate cross-validation strategies. For example, the naïve application of cross-validation to data that have spatial or spatio-temporal dependencies may lead to overly optimistic evaluations [Roberts et al., 2017]. In general, one also desires to minimise errors associated with either under-specification or over-specification of the model, a problem known as the bias-variance trade-off [Geman et al., 1992]. However, several recent advances have been made to reduce overfitting in ML models, for instance, regularization techniques in DNNs [Kukačka et al., 2017]. Moreover, when interpreting comparisons between ML and statistical methods, we should be cognizant that just as some ML methods require expert knowledge, the accuracy of statistical methods can also vary with the skill of the practitioner. Thompson and Calkin [2011] also emphasize the need for identifying sources of uncertainty in modeling so that they can better managed.

5.3 Implementation Challenges

Beyond data and model selection, two important considerations for model specification are feature selection and spatial autocorrelation. Knowledge of the problem domain is extremely important in identifying a set of candidate features. However, while many ML methods are not limited by the number of features, more variables do not necessarily make for a more accurate, interpretable, or easily implemented model [Schoenberg, 2016, Breiman, 2001] and can lead to overfitting and increased computational time. Two different ML methods to enable selection of a reduced and more optimal set of features include GAs and PSO. Sachdeva et al. [2018] used a GA to select input features for BRT and found this method gave the best accuracy compared with ANN, RF, SVM, SVM with PSO (PSO-SVM), DTs, logistic regression, and NB. Hong et al. [2018] employed a similar approach for fire susceptibility mapping and found this led to improvements for both SVM and RF compared with their non-optimized counterparts. Tracy et al. [2018] used a novel random subset feature selection algorithm for feature selection, which they found led to higher AUC values and lower model complexity. Jaafari et al. [2019] used a NFM combined with the imperialist competitive algorithm (a variant of GA) for feature selection which led to very high model accuracy (0.99) in their study. Tien Bui et al. [2017] used PSO to choose inputs to a NFN and found this improved results. [Zhang et al., 2019] also considered the information gain ratio for feature selection. As noted in Moritz et al. [2012] and Mayr et al. [2018], one should also take spatial autocorrelation into account when modeling fire probabilities spatially. In general, the presence of spatial autocorrelation violates the assumption of independence for parametric models, which can degrade model performance. One approach to deal with autocorrelation requires subsampling to remove any spatial autocorrelation Moritz et al. [2012]. It is also often necessary to subsample from non-fire locations due to class imbalance between ignitions and non-ignitions (e.g., Cao et al. [2017], Zhang et al. [2019]). Song et al. [2017] considered spatial econometric models and found a spatial autocorrelation model worked better than RF, although Kim et al. [2019] note that RF may be robust to spatial autocorrelation with large samples. In contrast to many ML methods, a strength of CNNs is its ability to exploit spatial correlation in the data to enable the extraction of spatial features.

5.4 Interpretation

A major obstacle for the adoption of ML methods to fire modeling tasks is the perceived lack of interpretability or explainability of such methods, which are often considered to be “black box” models. Users (in this case fire fighters and managers) need to trust ML model predictions, and so have the confidence and justification to apply these models, particularly in cases where proposed solutions are considered novel. Model interpretability should therefore be an important aspect of model development if models are to be selected and deployed in fire management operations. Model interpretability varies significantly across the different types of ML. For example, conventional thinking is that tree-based methods are more interpretable than neural network methods. This is because a single decision tree classifier can be rendered as a flow chart corresponding to if-then-else statements, whereas an ANN represents a nonlinear function

1289 approximated through a series of nonlinear activations. However, because they combine multiple trees in
1290 an optimized way, ensemble tree classifiers are less interpretable than single tree classifiers. On the other
1291 hand, BNs are one example of an ML technique where good explanations for results can be inferred due
1292 to their graphical representation; however, full Bayesian learning on large-scale data is very computationally
1293 expensive which may have limited early applications; however, as computational power has increased
1294 we have seen an increase in the popularity of BNs in wildfire science and management applications (e.g.,
1295 [Penman et al. \[2015\]](#), [Papakosta et al. \[2017\]](#)).

1296 DL-based architectures are widely considered to be among the least interpretable ML models, despite
1297 the fact that they can achieve very accurate function approximation [[Chakraborty et al., 2017](#)]. In fact, this
1298 is demonstrative of the well-known trade-off between prediction accuracy and interpretability (see [Kuhn
1299 and Johnson \[2013\]](#) for an in-depth discussion). The ML community, however, recognizes the problem
1300 of interpretability and work is underway to develop methods that allow for greater interpretability of ML
1301 methods, including methods for DL (see for example, [McGovern et al. \[2019\]](#)) or model-agnostic approaches
1302 [[Ribeiro et al., 2016](#)]. [Runge et al. \[2019\]](#) further argue that casual inference methods should be used in
1303 conjunction with predictive models to improve our understanding of physical systems. Finally, it is worth
1304 noting that assessing variable importance (see Sec. 4.3.4) for a given model can play a role in model
1305 interpretation.

1306 5.5 Opportunities

1307 Our review highlights a number of potential opportunities in wildfire science and management for ML
1308 applications where ML has not yet been applied or is under-utilized. Here we examine ML advances in
1309 other areas of environmental science that have analogous problems in wildland fire science and which may
1310 be useful for identifying further ML applications. For instance, [Li et al. \[2011\]](#) compared ML algorithms for
1311 spatial interpolation and found that a RF model combined with geostatistical methods yielded good results;
1312 a similar method could be used to improve interpolation of fire weather observations from weather stations,
1313 and so enhance fire danger monitoring. [Rasp and Lerch \[2018\]](#) showed that ANNs could improve weather
1314 forecasts by post-processing ensemble forecasts, an approach which could similarly be applied to improve
1315 short-term forecasts of fire weather. [Belayneh et al. \[2014\]](#) used ANNs and SVMs combined with wavelet
1316 transforms for long term drought forecasting in Ethiopia; such methods could also be useful for forecasting
1317 drought in the context of fire danger potential. In the context of numerical weather prediction, [Cohen et al.
1318 \[2019\]](#) found better predictability using ML methods than dynamical models for subseasonal to seasonal
1319 weather forecasting, suggesting similar applications for long-term fire weather forecasting. [McGovern et al.
1320 \[2017\]](#) discussed how AI techniques can be leveraged to improve decision making around high-impact
1321 weather. More recently, [Reichstein et al. \[2019\]](#) have further argued for the use of DL in the environmental
1322 sciences, citing its potential to extract spatiotemporal features from large geospatial datasets. [Kussul et al.
1323 \[2017\]](#) used CNNs to classify land cover and crop types and found that CNNs improved the results over
1324 standard ANN models; a similar approach could be used for fuels classification, which is an important input
1325 to fire behaviour prediction models. [Shi et al. \[2016\]](#) also used CNNs to detect clouds in remote sensed
1326 imagery and were able to differentiate between thin and thick cloud. A similar approach could be used
1327 for smoke detection, which is important for fire detection, as well as in determining the presence of false
1328 negatives in hotspot data (due to smoke or cloud obscuration). Finally, recent proposals have called for
1329 hybrid models that combine process-based models and ML methods [[Reichstein et al., 2019](#)]. For example,
1330 ML models may replace user-specified parameterizations in numerical weather prediction models [[Brenowitz
1331 and Bretherton, 2018](#)]. Other recent approaches use ML methods to determine the solutions to nonlinear
1332 partial differential equations [[Raissi and Karniadakis \[2018\]](#), [Raissi et al. \[2019\]](#)]. Such methods could find
1333 future applications in improving fire behaviour prediction models based on computationally expensive
1334 physics-based fire simulators, in coupled fire-atmosphere models, or in smoke dispersion modeling. In any
1335 case, the applications of ML that we have outlined are meant for illustrative purposes and are not meant
1336 to represent an exhaustive list of all possible applications.

5.6 Implications for fire management

We believe ML has been under-utilized in fire management, particularly with respect to problems belonging to either predictive or prescriptive analytics. Fire management comprises a set of risk control measures, which are often cast in the framework of the emergency response phases: prevention; mitigation; preparedness; response; recovery; and review [Tymstra et al., 2019]. In terms of financial expenditure, by far the largest percentage spent in the response phase [Stocks and Martell, 2016]. In practice, fire management is largely determined by the need to manage resources in response to active or expected wildfires, typically for lead times of days to weeks, or to manage vegetative fuels. This suggests the opportunity for increased research in areas of fire weather prediction, fire occurrence prediction, and fire behaviour prediction, as well as optimizing fire operations and fuel treatments. The identification of these areas, as well as the fact that wildfire is both a spatial and temporal process, further reiterate the need for ML applications for time series forecasting.

From this review, there were few papers that used time series ML methods for forecasting problems, suggesting an opportunity for further work in this area. In particular, recurrent neural networks (RNNs) were used for fire behavior prediction [Cheng and Wang, 2008, Kozik et al., 2013, 2014] and fire occurrence prediction [Dutta et al., 2013]. The most common variant of RNNs are Long Short Term Memory (LSTM) networks [Hochreiter and Schmidhuber, 1997], which have been used for burned area prediction [Liang et al., 2019] and fire detection [Cao et al., 2019]. Because these methods implicitly model dynamical processes, they should lead to improve forecasting models compared with standard ANNs. For example Gensler et al. [2017] have used LSTMs to forecast solar power and Kim et al. [2017] used CNNs combined with LSTM for forecasting precipitation. We anticipate that these methods could also be employed for fire weather, fire occurrence, and fire behaviour prediction.

We note that there are a number of operational research and management science methods used in fire management research including queuing, optimization, and simulation of complex system dynamics (e.g., Martell [2015]) where ML algorithms don't seem to provide an obvious alternative. For example, planning models to simulate the interactions between fire management resource configurations and fire dynamics reviewed by [Mavsar et al., 2013]. From our review, a few papers used agent-based learning methods for fire management. In particular, reinforcement learning was used for optimizing fuel treatments [Lauer et al., 2017] or for autonomous control of aircraft for fire monitoring [Julian and Kochenderfer, 2018a]. GAs were used for generating optimal firelines for active fires [Homchaudhuri et al., 2010] and for reducing the time for fire simulation [Cencerrado et al., 2014]. However, more work is needed to identify where ML methods could contribute to tactical, operational, or strategic fire management decision making.

An important challenge for the fire research and management communities is enabling the transition of potentially useful ML models to fire management operations. Although we identified several papers that emphasized their ML models could be deployed in fire management operations [Artés et al., 2016, Alonso-Betanzos et al., 2002, Iliadis, 2005, Stojanova et al., 2012, Davis et al., 1989, 1986, Liu et al., 2015], it can be difficult to assess whether and how a study has been adopted by, or influenced, fire management agencies. This challenge is often exacerbated by a lack of resources and/or funding, as well as the different priorities and institutional cultures of researchers and fire managers. One possible solution to this problem would be the formation of working groups dedicated to enabling this transition, preferably at the research proposal phase. In general, enabling operational ML methods will require tighter integration and greater collaboration between the research and management communities, particularly with regards to project design, data compilation and variable selection, implementation, and interpretation. However, it is worth noting that this is not a problem unique to ML, it is a long-standing and common issue in many areas of fire research and other applied science disciplines, where continuous effort is required to maintain communications and relationships between researchers and practitioners.

Finally, we would like to stress that we believe the wildfire research and management communities should play an active role in providing relevant, high quality, and freely available wildfire data for use by practitioners of ML methods. For example, burned area and fire weather data made available by Cortez and Morais [2007] was subsequently used by a number of authors in their work. It is imperative that the

1387 quality of data collected by management agencies be as robust as possible, as the results of any modelling
 1388 process are dependent upon the data used for analysis. It is worth considering how new data on, for
 1389 example, hourly fire growth or the daily use of fire management resources, could be used in ML methods
 1390 to yield better predictions or management recommendations — using new tools to answer new questions
 1391 may require better or more complete data. Conversely, we must recognize that despite ML models being
 1392 able to learn on their own, expertise in wildfire science is necessary to ensure realistic modelling of wildfire
 1393 processes, while the complexity of some ML methods (e.g., DL) requires a dedicated and sophisticated
 1394 knowledge of their application (we note that many of the most popular ML methods used in this study are
 1395 fairly easy to implement, such as RF, MaxEnt, and DTs). The observation that no single ML algorithm is
 1396 superior for all classes of problem, an idea encapsulated by the “no free lunch” theorem [Wolpert, 1996],
 1397 further reinforces the need for domain-specific knowledge. Thus, the proper implementation of ML in
 1398 wildfire science is a challenging endeavor, often requiring multidisciplinary teams and/or interdisciplinary
 1399 specialists to effectively produce meaningful results.

1400 5.7 A word of caution

1401 ML holds tremendous potential for a number of wildfire science and management problem domains. As
 1402 indicated in this review, much work has already been undertaken in a number of areas, although further
 1403 work is clearly needed for fire management specific problems. Despite this potential, ML should not be
 1404 considered a panacea for all fire research areas. ML is best suited to problems where there is sufficient high-
 1405 quality data, and this is not always the case. For example, for problems related to fire management policy,
 1406 data is needed at large spatiotemporal scales (i.e., ecosystem/administrative spatial units at timescales of
 1407 decades or even centuries), and such data may simply not yet exist in current inventories. At the other
 1408 extreme, data is needed at very fine spatiotemporal scales for fire spread and behavior modeling, including
 1409 high resolution fuel maps and surface weather variables which are often not available at the required scale
 1410 and are difficult to acquire even in an experimental context. Another limitation of ML may occur when
 1411 one attempts make predictions where no analog exists in the observed data, such as may be the case with
 1412 climate change prediction.

1413 6 Conclusions

1414 Our review shows that the application of ML methods in wildfire science and management has been steadily
 1415 increasing since their first use in the 1990s, across core problem domains using a wide range ML methods.
 1416 The bulk of work undertaken thus far has used traditional methods such as RF, BRT, MaxEnt, SVM
 1417 and ANNs, partly due to the ease of application and partly due to their simple interpretability in many
 1418 cases. However, problem domains associated with predictive (e.g., predicted fire behavior) or prescriptive
 1419 analytics (e.g. optimizing fire management decisions) have seen much less work with ML methods. We
 1420 therefore suggest opportunities exist for both the wildfire community and ML practitioners to apply ML
 1421 methods in these areas. Moreover, the increasing availability of large spatio-temporal datasets, from climate
 1422 models or remote sensing for example, may be amenable to the use of deep learning methods, which can
 1423 efficiently extract spatial or temporal features from data. Another major opportunity is the application of
 1424 agent based learning to fire management operations, although many other opportunities exist. However,
 1425 we must recognize that despite ML models being able to learn on their own, expertise in wildfire science
 1426 is necessary to ensure realistic modelling of wildfire processes across multiple scales, while the complexity
 1427 of some ML methods (e.g. DL) requires a dedicated and sophisticated knowledge of their application.
 1428 Furthermore, a major obstacle for the adoption of ML methods to fire modeling tasks is the perceived
 1429 lack of interpretability of such methods, which are often considered to be black box models. The ML
 1430 community, however, recognizes this problem and work is underway to develop methods that allow for
 1431 greater interpretability of ML methods (see for example, [McGovern et al., 2019]). Data driven approaches
 1432 are by definition data dependent — if the fire management community wants to more fully exploit powerful

1433 ML methods, we need to consider data as a valuable resource and examine what further information on
 1434 fire events or operations are needed to apply ML approaches to management problems. Thus, wildland
 1435 fire science is a diverse multi-faceted discipline that requires a multi-pronged approach, a challenge made
 1436 greater by the need to mitigate and adapt to current and future fire regimes.

1437 Acknowledgments

1438 The motivation for this paper arose from the “Not the New Normal” BC AI Wildfire Symposium held in
 1439 Vancouver, BC, on 12 October 2018. The authors would also like to thank Intact Insurance, the Canadian
 1440 Partnership for Wildland Fire Science, the NSERC Discovery Grants program and the Microsoft AI for
 1441 Social Good program for their support.

1442 References

- 1443 Baker Abdalhaq, Ana Cortés, Tomàs Margalef, and Emilio Luque. Enhancing wildland fire prediction on
 1444 cluster systems applying evolutionary optimization techniques. *Future Generation Computer Systems*,
 1445 21(1):61–67, 2005. ISSN 0167739X. doi: 10.1016/j.future.2004.09.013.
- 1446 Hamed Adab. Landfire hazard assessment in the Caspian Hyrcanian forest ecoregion with the long-term
 1447 MODIS active fire data. *Natural Hazards*, 87(3):1807–1825, jul 2017. ISSN 0921-030X. doi: 10.1007/
 1448 s11069-017-2850-2. URL <http://link.springer.com/10.1007/s11069-017-2850-2>.
- 1449 Hamed Adab, Azadeh Atabati, Sandra Oliveira, and Ahmad Moghaddam Gheshlagh. Assessing fire hazard
 1450 potential and its main drivers in Mazandaran province, Iran: a data-driven approach. *Environmental*
 1451 *Monitoring and Assessment*, 190(11):670, nov 2018. ISSN 0167-6369. doi: 10.1007/s10661-018-7052-1.
 1452 URL <http://link.springer.com/10.1007/s10661-018-7052-1>.
- 1453 Moulay A. Akhloufi, Roger Booto Tokime, and Hassan Ellassady. Wildland fires detection and
 1454 segmentation using deep learning. In Mohammad S. Alam, editor, *Pattern Recognition and Track-*
 1455 *ing XXIX*, page 11. SPIE, apr 2018. ISBN 9781510618091. doi: 10.1117/12.2304936. URL
 1456 [https://www.spiedigitallibrary.org/conference-proceedings-of-spie/10649/2304936/
 1457 Wildland-fires-detection-and-segmentation-using-deep-learning/10.1117/12.2304936.
 1458 full](https://www.spiedigitallibrary.org/conference-proceedings-of-spie/10649/2304936/Wildland-fires-detection-and-segmentation-using-deep-learning/10.1117/12.2304936.full).
- 1459 K. R. Al-Rawi, J. L. Casanova, and A. Calle. Burned area mapping system and fire detection system, based
 1460 on neural networks and NOAA-AVHRR imagery. *International Journal of Remote Sensing*, 22(10):2015–
 1461 2032, jan 2001. ISSN 0143-1161. doi: 10.1080/01431160117531. URL [https://www.tandfonline.com/
 1462 doi/full/10.1080/01431160117531](https://www.tandfonline.com/doi/full/10.1080/01431160117531).
- 1463 K R Al-Rawi, J L Casanova, A Romo, and E M Louakfaoui. Integrated fire evolution monitoring system
 1464 (IFEMS) for monitoring spatial-temporal behaviour of multiple fire phenomena. *International Journal*
 1465 *of Remote Sensing*, 23(10):1967–1983, 2002. ISSN 01431161. doi: 10.1080/01431160110069809. URL
 1466 <http://www.tandfonline.com/action/journalInformation?journalCode=tres20>.
- 1467 Dima Alberg. An Interval Tree Approach to Predict Forest Fires using Meteorological Data. *International*
 1468 *Journal of Computer Applications*, 132(4):17–22, 2015. doi: 10.5120/ijca2015907398.
- 1469 Andrew Aldersley, Steven J. Murray, and Sarah E. Cornell. Global and regional analysis of climate and
 1470 human drivers of wildfire. *Science of the Total Environment*, 409(18):3472–3481, 2011. ISSN 00489697.
 1471 doi: 10.1016/j.scitotenv.2011.05.032. URL <http://dx.doi.org/10.1016/j.scitotenv.2011.05.032>.

- 1472 Dmitriy Alexandrov, Elizaveta Pertseva, Ivan Berman, Igor Pantiukhin, and Aleksandr Kapitonov. Anal-
 1473 ysis of machine learning methods for wildfire security monitoring with an unmanned aerial vehicles. In
 1474 *Conference of Open Innovation Association, FRUCT*, volume 2019-April, pages 3–9. IEEE Computer
 1475 Society, may 2019. ISBN 9789526865386. doi: 10.23919/FRUCT.2019.8711917.
- 1476 Samaher Al-Janabi, Ibrahim Al-Shourbaji, and Mahdi A. Salman. Assessing the suitability of soft comput-
 1477 ing approaches for forest fires prediction. *Applied Computing and Informatics*, 14(2):214–224, jul 2018.
 1478 ISSN 2210-8327. doi: 10.1016/J.ACI.2017.09.006. URL [https://www.sciencedirect.com/science/
 1479 article/pii/S2210832717301539](https://www.sciencedirect.com/science/article/pii/S2210832717301539).
- 1480 A. Alonso-Benito, P. A. Hernandez-Leal, A. Gonzalez-Calvo, M. Arbelo, and A. Barreto. Analysis of
 1481 Different Methods for Burnt Area Estimation using Remote Sensing and Ground Truth Data. In *IGARSS
 1482 2008 - 2008 IEEE International Geoscience and Remote Sensing Symposium*, pages III – 828–III – 831.
 1483 IEEE, 2008. ISBN 978-1-4244-2807-6. doi: 10.1109/IGARSS.2008.4779477. URL [http://ieeexplore.
 1484 ieee.org/document/4779477/](http://ieeexplore.ieee.org/document/4779477/).
- 1485 Amparo Alonso-Betanzos, Oscar Fontenla-Romero, Bertha Guijarro-BerdiñasBerdi, Elena Hernández-
 1486 Pereira, Juan Canda, Eulogio Jimenez, José Luis Legido, Susana MuñizMu, Cristina Paz-Andrade,
 1487 and María Inmaculada Paz-Andrade. A Neural Network Approach for Forestal Fire Risk Estimation.
 1488 Technical report, 2002.
- 1489 Amparo Alonso-Betanzos, Oscar Fontenla-Romero, Bertha Guijarro-Berdiñas, Elena Hernández-Pereira,
 1490 María Inmaculada Paz Andrade, Eulogio Jiménez, Jose Luis Legido Soto, and Tarsy Carballas. An
 1491 intelligent system for forest fire risk prediction and fire fighting management in Galicia. *Expert Systems
 1492 with Applications*, 25(4):545–554, 2003. ISSN 09574174. doi: 10.1016/S0957-4174(03)00095-2.
- 1493 N. S. Altman. An introduction to kernel and nearest-neighbor nonparametric regression. *American Statis-
 1494 tician*, 46(3):175–185, 1992. ISSN 15372731. doi: 10.1080/00031305.1992.10475879.
- 1495 Giuseppe Amatulli and Andrea Camia. Exploring the relationships of fire occurrence variables by means
 1496 of CART and MARS models. In *Proceedings of the 4th International Wildland Fire Conference*, pages
 1497 1–11, 2007.
- 1498 Giuseppe Amatulli, Maria João Rodrigues, Marco Trombetti, and Raffaella Lovreglio. Assessing long-
 1499 term fire risk at local scale by means of decision tree technique. *Journal of Geophysical Research:
 1500 Biogeosciences*, 111(G4), dec 2006. ISSN 01480227. doi: 10.1029/2005JG000133. URL [http://doi.
 1501 wiley.com/10.1029/2005JG000133](http://doi.wiley.com/10.1029/2005JG000133).
- 1502 Giuseppe Amatulli, Andrea Camia, and Jesús San-Miguel-Ayanz. Estimating future burned areas un-
 1503 der changing climate in the EU-Mediterranean countries. *Science of The Total Environment*, 450-
 1504 451:209–222, apr 2013. ISSN 0048-9697. doi: 10.1016/J.SCITOTENV.2013.02.014. URL <https://www.sciencedirect.com/science/article/pii/S0048969713001770>.
- 1506 K. Angayarkkani and N. Radhakrishnan. An Intelligent System For Effective Forest Fire Detection Using
 1507 Spatial Data. feb 2010. URL <http://arxiv.org/abs/1002.2199>.
- 1508 K. Angayarkkani and N. Radhakrishnan. An effective technique to detect forest fire region through AN-
 1509 FIS with spatial data. In *ICECT 2011 - 2011 3rd International Conference on Electronics Computer
 1510 Technology*, volume 3, pages 24–30, 2011. ISBN 9781424486779. doi: 10.1109/ICECTECH.2011.5941794.
- 1511 Bachisio Arca, Tiziano Ghisu, and Giuseppe A. Trunfio. GPU-accelerated multi-objective optimization
 1512 of fuel treatments for mitigating wildfire hazard. *Journal of Computational Science*, 11:258–268, 2015.
 1513 ISSN 18777503. doi: 10.1016/j.jocs.2015.08.009.

- 1514 Sally Archibald, David P. Roy, Brian W. van Wilgen, and Robert J. Scholes. What limits fire? An
 1515 examination of drivers of burnt area in Southern Africa. *Global Change Biology*, 15(3):613–630, 2009.
 1516 ISSN 13541013. doi: 10.1111/j.1365-2486.2008.01754.x.
- 1517 Juan P. Argañaraz, Gregorio Gavier Pizarro, Marcelo Zak, Marcos A. Landi, and Laura M Bellis. Hu-
 1518 man and biophysical drivers of fires in Semiarid Chaco mountains of Central Argentina. *Science of*
 1519 *The Total Environment*, 520:1–12, jul 2015. ISSN 0048-9697. doi: 10.1016/J.SCITOTENV.2015.
 1520 02.081. URL [https://www-sciencedirect-com.login.ezproxy.library.ualberta.ca/science/
 1521 article/pii/S0048969715002338](https://www-sciencedirect-com.login.ezproxy.library.ualberta.ca/science/article/pii/S0048969715002338).
- 1522 Hilary Arksey and Lisa O’Malley. Scoping studies: towards a methodological framework.
 1523 *International Journal of Social Research Methodology*, 8(1):19–32, feb 2005. ISSN 1364-
 1524 5579. doi: 10.1080/1364557032000119616. URL [http://www.tandfonline.com/doi/abs/10.1080/
 1525 1364557032000119616](http://www.tandfonline.com/doi/abs/10.1080/1364557032000119616).
- 1526 James D. Arnold, Simon C. Brewer, and Philip E. Dennison. Modeling Climate-Fire Connections within
 1527 the Great Basin and Upper Colorado River Basin, Western United States. *Fire Ecology*, 10(2):64–75,
 1528 aug 2014. ISSN 19339747. doi: 10.4996/fireecology.1002064. URL [http://fireecologyjournal.org/
 1529 journal/abstract/?abstract=220](http://fireecologyjournal.org/journal/abstract/?abstract=220).
- 1530 A. Arpacı, B. Malowerschnig, O. Sass, and H. Vacik. Using multi variate data mining techniques for
 1531 estimating fire susceptibility of Tyrolean forests. *Applied Geography*, 53:258–270, sep 2014. ISSN
 1532 0143-6228. doi: 10.1016/J.APGEOG.2014.05.015. URL [https://www.sciencedirect.com/science/
 1533 article/abs/pii/S0143622814001106](https://www.sciencedirect.com/science/article/abs/pii/S0143622814001106).
- 1534 B.C. Arrue, A. Ollero, and J.R. Matinez de Dios. An intelligent system for false alarm reduction in
 1535 infrared forest-fire detection. *IEEE Intelligent Systems*, 15(3):64–73, may 2000. ISSN 1094-7167. doi:
 1536 10.1109/5254.846287. URL <http://ieeexplore.ieee.org/document/846287/>.
- 1537 Tomàs Artés, Andrés Cencerrado, Ana Cortés, and Tomàs Margalef. Core Allocation Policies on Multicore
 1538 Platforms to Accelerate Forest Fire Spread Predictions. *PPAM 2013: Parallel Processing and Applied*
 1539 *Mathematics*, pages 151–160, 2014. ISSN 16113349. doi: 10.1007/978-3-642-55195-6. URL [https:
 1540 //link.springer.com/chapter/10.1007/978-3-642-55195-6_{_}14{#}enumeration](https://link.springer.com/chapter/10.1007/978-3-642-55195-6_{_}14{#}enumeration).
- 1541 Tomàs Artés, Andrés Cencerrado, Ana Cortés, and Tomàs Margalef. Real-time genetic spatial opti-
 1542 mization to improve forest fire spread forecasting in high-performance computing environments. *In-*
 1543 *ternational Journal of Geographical Information Science*, 30(3):594–611, 2016. ISSN 13623087. doi:
 1544 10.1080/13658816.2015.1085052. URL <http://dx.doi.org/10.1080/13658816.2015.1085052>.
- 1545 Tomàs Artés, Andrés Cencerrado, Ana Cortés, and Tomàs Margalef. Time aware genetic algorithm for
 1546 forest fire propagation prediction: exploiting multi-core platforms. *Concurrency Computation*, 29(9):
 1547 1–18, 2017. ISSN 15320634. doi: 10.1002/cpe.3837.
- 1548 Davide Ascoli, Giorgio Vacchiano, Renzo Motta, and Giovanni Bovio. Building Rothermel fire behaviour
 1549 fuel models by genetic algorithm optimisation. *International Journal of Wildland Fire*, 24(3):317–328,
 1550 2015. ISSN 10498001. doi: 10.1071/WF14097.
- 1551 Rui Ba, Chen Chen, Jing Yuan, Weiguo Song, and Siuming Lo. SmokeNet: Satellite Smoke Scene Detection
 1552 Using Convolutional Neural Network with Spatial and Channel-Wise Attention. *Remote Sensing*, 11(14):
 1553 1702, jul 2019. ISSN 2072-4292. doi: 10.3390/rs11141702. URL [https://www.mdpi.com/2072-4292/
 1554 11/14/1702](https://www.mdpi.com/2072-4292/11/14/1702).
- 1555 Shitai Bao, Ningchuan Xiao, Zehui Lai, Heyuan Zhang, and Changjoo Kim. Optimizing watchtower
 1556 locations for forest fire monitoring using location models. *Fire Safety Journal*, 71(December 2013):100–
 1557 109, 2015. ISSN 03797112. doi: 10.1016/j.firesaf.2014.11.016. URL [http://dx.doi.org/10.1016/j.
 1558 firesaf.2014.11.016](http://dx.doi.org/10.1016/j.firesaf.2014.11.016).

- 1559 Avi Bar Massada, Alexandra D. Syphard, Susan I. Stewart, and Volker C. Radeloff. Wildfire ignition-
 1560 distribution modelling: a comparative study in the Huron?Manistee National Forest, Michigan, USA.
 1561 *International Journal of Wildland Fire*, 22(2):174, apr 2013. ISSN 1049-8001. doi: 10.1071/WF11178.
 1562 URL <http://www.publish.csiro.au/?paper=WF11178>.
- 1563 Panagiotis Barmpoutis, Kosmas Dimitropoulos, Kyriaki Kaza, and Nikos Grammalidis. Fire Detection from
 1564 Images Using Faster R-CNN and Multidimensional Texture Analysis. In *ICASSP, IEEE International
 1565 Conference on Acoustics, Speech and Signal Processing - Proceedings*, volume 2019-May, pages 8301–
 1566 8305. Institute of Electrical and Electronics Engineers Inc., may 2019. ISBN 9781479981311. doi:
 1567 10.1109/ICASSP.2019.8682647.
- 1568 K. Barrett, A. D. McGuire, E. E. Hoy, and E. S. Kasischke. Potential shifts in dominant forest cover in
 1569 interior Alaska driven by variations in fire severity. *Ecological Applications*, 21(7):2380–2396, 2011. ISSN
 1570 10510761. doi: 10.1890/10-0896.1.
- 1571 Hossein Bashari, Ali Asghar Naghipour, Seyed Jamaledin Khajeddin, Hamed Sangoony, and Pejman Tah-
 1572 masebi. Risk of fire occurrence in arid and semi-arid ecosystems of Iran: an investigation using Bayesian
 1573 belief networks. *Environmental Monitoring and Assessment*, 188(9):531, sep 2016. ISSN 0167-6369. doi:
 1574 10.1007/s10661-016-5532-8. URL <http://link.springer.com/10.1007/s10661-016-5532-8>.
- 1575 Bryson C. Bates, Andrew J. Dowdy, Richard E. Chandler, Bryson C. Bates, Andrew J. Dowdy, and
 1576 Richard E. Chandler. Classification of Australian Thunderstorms Using Multivariate Analyses of Large-
 1577 Scale Atmospheric Variables. *Journal of Applied Meteorology and Climatology*, 56(7):1921–1937, jul
 1578 2017. ISSN 1558-8424. doi: 10.1175/JAMC-D-16-0271.1. URL [http://journals.ametsoc.org/doi/
 1579 10.1175/JAMC-D-16-0271.1](http://journals.ametsoc.org/doi/10.1175/JAMC-D-16-0271.1).
- 1580 Enric Batllori, Marc-André Parisien, Meg A. Krawchuk, and Max A. Moritz. Climate change-induced
 1581 shifts in fire for Mediterranean ecosystems. *Global Ecology and Biogeography*, 22(10):1118–1129, oct
 1582 2013. ISSN 1466822X. doi: 10.1111/geb.12065. URL <http://doi.wiley.com/10.1111/geb.12065>.
- 1583 Peter Bauer, Alan Thorpe, and Gilbert Brunet. The quiet revolution of numerical weather prediction.
 1584 *Nature*, 525(7567):47–55, sep 2015. ISSN 0028-0836. doi: 10.1038/nature14956. URL [http://www.
 1585 nature.com/articles/nature14956](http://www.nature.com/articles/nature14956).
- 1586 A. Belayneh, J. Adamowski, B. Khalil, and B. Ozga-Zielinski. Long-term SPI drought forecasting in
 1587 the Awash River Basin in Ethiopia using wavelet neural networks and wavelet support vector regression
 1588 models. *Journal of Hydrology*, 508:418–429, jan 2014. ISSN 00221694. doi: 10.1016/j.jhydrol.2013.10.052.
- 1589 Mar Bisquert, Eduardo Caselles, Juan Manuel Sánchez, and Vicente Caselles. Application of artificial
 1590 neural networks and logistic regression to the prediction of forest fire danger in Galicia using MODIS data.
 1591 *International Journal of Wildland Fire*, 21(8):1025, dec 2012. ISSN 1049-8001. doi: 10.1071/WF11105.
 1592 URL <http://www.publish.csiro.au/?paper=WF11105>.
- 1593 Karen D. Blouin, Mike D. Flannigan, Xianli Wang, and Bohdan Kochtubajda. Ensemble lightning predic-
 1594 tion models for the province of Alberta, Canada. *International Journal of Wildland Fire*, 25(4):421–432,
 1595 apr 2016. ISSN 1049-8001. doi: 10.1071/WF15111. URL [http://www.publish.csiro.au/?paper=
 1596 WF15111](http://www.publish.csiro.au/?paper=WF15111).
- 1597 William J. Bond and Jon E. Keeley. Fire as a global 'herbivore': The ecology and evolution of flammable
 1598 ecosystems. *Trends in Ecology and Evolution*, 20(7):387–394, jul 2005. ISSN 01695347. doi: 10.1016/j.
 1599 tree.2005.04.025.
- 1600 Yan Boulanger, Marc-André Parisien, and Xianli Wang. Model-specification uncertainty in future area
 1601 burned by wildfires in Canada. *International Journal of Wildland Fire*, 27(3):164, apr 2018. ISSN
 1602 1049-8001. doi: 10.1071/WF17123. URL <http://www.publish.csiro.au/?paper=WF17123>.

- 1603 Curtis M. Bradley, Chad T. Hanson, and Dominick A. DellaSala. Does increased forest protection corre-
 1604 spond to higher fire severity in frequent-fire forests of the western United States? *Ecosphere*, 7(10):1–13,
 1605 2016. ISSN 21508925. doi: 10.1002/ecs2.1492.
- 1606 Jonathan Branham, Nicholas Hamilton, Dale Hamilton, and Barry Myers. Evaluation of
 1607 Image Spatial Resolution for Machine Learning Mapping of Wildland Fire Effects, jul
 1608 2017. URL [https://scholarworks.boisestate.edu/icur/2017/Poster{ }Session/26https://](https://scholarworks.boisestate.edu/icur/2017/Poster{ }Session/26https://link.springer.com/chapter/10.1007/978-3-030-01054-6{ }29)
 1609 link.springer.com/chapter/10.1007/978-3-030-01054-6{ }29.
- 1610 Leo Breiman. Statistical modeling: The two cultures. *Statistical Science*, 16(3):199–215, 2001. ISSN
 1611 08834237. doi: 10.1214/ss/1009213726.
- 1612 Leo Breiman. *Classification and regression trees*. Routledge, 2017.
- 1613 Leo Breiman, Jerome Friedman, Richard A Olshen, and Charles J Stone. Classification and regression
 1614 trees Chapman & Hall. *New York*, 1984.
- 1615 N. D. Brenowitz and C. S. Bretherton. Prognostic Validation of a Neural Network Unified Physics
 1616 Parameterization. *Geophysical Research Letters*, 45(12):6289–6298, jun 2018. ISSN 00948276. doi:
 1617 10.1029/2018GL078510. URL <http://doi.wiley.com/10.1029/2018GL078510>.
- 1618 Steven P. Brumby, Neal R. Harvey, Jeffrey J. Bloch, James P. Theiler, Simon J. Perkins, Aaron C. Young,
 1619 and John J. Szymanski. Evolving forest fire burn severity classification algorithms for multispectral
 1620 imagery. *Algorithms for Multispectral, Hyperspectral, and Ultraspectral Imagery VII*, 4381(August 2001):
 1621 236–245, 2001. doi: 10.1117/12.437013.
- 1622 C. E. Buckland, R. M. Bailey, and D. S. G. Thomas. Using artificial neural networks to predict
 1623 future dryland responses to human and climate disturbances. *Scientific Reports*, 9(1):3855, dec
 1624 2019. ISSN 2045-2322. doi: 10.1038/s41598-019-40429-5. URL [http://www.nature.com/articles/](http://www.nature.com/articles/s41598-019-40429-5)
 1625 [s41598-019-40429-5](http://www.nature.com/articles/s41598-019-40429-5).
- 1626 Declan Butler. AI summit aims to help world’s poorest, jun 2017. ISSN 14764687.
- 1627 Wenhua Cai, Jian Yang, Zhihua Liu, Yuanman Hu, and Peter J. Weisberg. Post-fire tree recruitment of a
 1628 boreal larch forest in Northeast China. *Forest Ecology and Management*, 307:20–29, 2013. ISSN 03781127.
 1629 doi: 10.1016/j.foreco.2013.06.056. URL <http://dx.doi.org/10.1016/j.foreco.2013.06.056>.
- 1630 X. Cao, J. Chen, B. Matsushita, H. Imura, and L. Wang. An automatic method for burn scar mapping
 1631 using support vector machines. *International Journal of Remote Sensing*, 30(3):577–594, feb 2009.
 1632 ISSN 0143-1161. doi: 10.1080/01431160802220219. URL [https://www.tandfonline.com/doi/full/](https://www.tandfonline.com/doi/full/10.1080/01431160802220219)
 1633 [10.1080/01431160802220219](https://www.tandfonline.com/doi/full/10.1080/01431160802220219).
- 1634 Yichao Cao, Feng Yang, Qingfei Tang, and Xiaobo Lu. An Attention Enhanced Bidirectional LSTM for
 1635 Early Forest Fire Smoke Recognition. *IEEE Access*, pages 1–1, oct 2019. doi: 10.1109/access.2019.
 1636 2946712.
- 1637 Yinxue Cao, Ming Wang, and Kai Liu. Wildfire Susceptibility Assessment in Southern China: A Compar-
 1638 ison of Multiple Methods. *International Journal of Disaster Risk Science*, 8(2):164–181, jun 2017. ISSN
 1639 21926395. doi: 10.1007/s13753-017-0129-6.
- 1640 Adrián Cardil, Blas Mola-Yudego, Ángela Blázquez-Casado, and José Ramón González-Olabarria. Fire
 1641 and burn severity assessment: Calibration of Relative Differenced Normalized Burn Ratio (RdNBR)
 1642 with field data. *Journal of Environmental Management*, 235(January):342–349, 2019. ISSN 10958630.
 1643 doi: 10.1016/j.jenvman.2019.01.077. URL <https://doi.org/10.1016/j.jenvman.2019.01.077>.

- 1644 Carlos Carrillo, Tomàs Artés, Ana Cortés, and Tomàs Margalef. Error function impact in dynamic data-
 1645 driven framework applied to forest fire spread prediction. *Procedia Computer Science*, 80:418–427, 2016.
 1646 ISSN 18770509. doi: 10.1016/j.procs.2016.05.342. URL [http://dx.doi.org/10.1016/j.procs.2016.](http://dx.doi.org/10.1016/j.procs.2016.05.342)
 1647 [05.342](http://dx.doi.org/10.1016/j.procs.2016.05.342).
- 1648 Mauro Castelli, Leonardo Vanneschi, and Aleš Popovič. PREDICTING BURNED AREAS OF FOR-
 1649 EST FIRES: AN ARTIFICIAL INTELLIGENCE APPROACH. *Fire Ecology*, 11(1):106–118, apr 2015.
 1650 ISSN 19339747. doi: 10.4996/fireecology.1101106. URL [http://fireecologyjournal.org/journal/](http://fireecologyjournal.org/journal/abstract/?abstract=236)
 1651 [abstract/?abstract=236](http://fireecologyjournal.org/journal/abstract/?abstract=236).
- 1652 Turgay Celik. Change detection in satellite images using a genetic algorithm approach. *IEEE Geoscience*
 1653 *and Remote Sensing Letters*, 7(2):386–390, 2010. ISSN 1545598X. doi: 10.1109/LGRS.2009.2037024.
- 1654 Andrés Cencerrado, Ana Cortés, and Tomàs Margalef. Genetic algorithm characterization for the quality
 1655 assessment of forest fire spread prediction. *Procedia Computer Science*, 9:312–320, 2012. ISSN 18770509.
 1656 doi: 10.1016/j.procs.2012.04.033.
- 1657 Andrés Cencerrado, Ana Cortés, and Tomàs Margalef. Applying probability theory for the quality as-
 1658 sessment of a wildfire spread prediction framework based on genetic algorithms. *The Scientific World*
 1659 *Journal*, 2013, 2013. ISSN 1537744X. doi: 10.1155/2013/728414.
- 1660 Andrés Cencerrado, Ana Cortés, and Tomàs Margalef. Response time assessment in forest fire spread
 1661 simulation: An integrated methodology for efficient exploitation of available prediction time. *Environ-*
 1662 *mental Modelling & Software*, 54:153–164, apr 2014. ISSN 1364-8152. doi: 10.1016/J.ENVSOFT.2014.
 1663 01.008. URL [https://www.sciencedirect-com.login.ezproxy.library.ualberta.ca/science/](https://www.sciencedirect-com.login.ezproxy.library.ualberta.ca/science/article/pii/S1364815214000176)
 1664 [article/pii/S1364815214000176](https://www.sciencedirect-com.login.ezproxy.library.ualberta.ca/science/article/pii/S1364815214000176).
- 1665 Supriyo Chakraborty, Richard Tomsett, Ramya Raghavendra, Daniel Harborne, Moustafa Alzantot,
 1666 Federico Cerutti, Mani Srivastava, Alun Preece, Simon Julier, Raghuvver M Rao, and Others. In-
 1667 terpretability of deep learning models: a survey of results. In *2017 IEEE SmartWorld, Ubiq-*
 1668 *uitous Intelligence & Computing, Advanced & Trusted Computed, Scalable Computing & Commu-*
 1669 *nications, Cloud & Big Data Computing, Internet of People and Smart City Innovation (Smart-*
 1670 *World/SCALCOM/UIC/ATC/CBDCOM/IOP/SCI)*, pages 1–6. IEEE, 2017.
- 1671 Fs Chapin, T.N. Hollingsworth, and Re Hewitt. Fire effects on seedling establishment success across
 1672 treeline: implications for future tree migration and flammability in a changing climate. 2014. URL
 1673 <http://digitalcommons.unl.edu/jfspresearch/82/>.
- 1674 Feng Chen, Yongsheng Du, Shukui Niu, and Jinlong Zhao. Modeling Forest Lightning Fire Occurrence
 1675 in the Daxinganling Mountains of Northeastern China with MAXENT. *Forests*, 6(12):1422–1438, apr
 1676 2015. ISSN 1999-4907. doi: 10.3390/f6051422. URL <http://www.mdpi.com/1999-4907/6/5/1422>.
- 1677 Tao Cheng and Jiaqiu Wang. Integrated Spatio-temporal Data Mining for Forest Fire Prediction. *Trans-*
 1678 *actions in GIS*, 12(5):591–611, sep 2008. ISSN 13611682. doi: 10.1111/j.1467-9671.2008.01117.x. URL
 1679 <http://doi.wiley.com/10.1111/j.1467-9671.2008.01117.x>.
- 1680 Khaled Chetehouna, Eddy El Tabach, Loubna Bouazaoui, and Nicolas Gascoin. Predicting the flame
 1681 characteristics and rate of spread in fires propagating in a bed of Pinus pinaster using Artificial Neu-
 1682 ral Networks. *Process Safety and Environmental Protection*, 98:50–56, nov 2015. ISSN 0957-5820.
 1683 doi: 10.1016/J.PSEP.2015.06.010. URL [https://www.sciencedirect.com/science/article/pii/](https://www.sciencedirect.com/science/article/pii/S0957582015001111)
 1684 [S0957582015001111](https://www.sciencedirect.com/science/article/pii/S0957582015001111).
- 1685 Tatenda T Chingono and C Mbohwa. Fire Hazard Modelling in Southern Africa. In *Proceedings of the*
 1686 *World Congress on Engineering and Computer Science*, San Francisco, 2015. URL [http://www.iaeng.](http://www.iaeng.org/publication/WCECS2015/WCECS2015_{_}pp514-519.pdf)
 1687 [org/publication/WCECS2015/WCECS2015_{_}pp514-519.pdf](http://www.iaeng.org/publication/WCECS2015/WCECS2015_{_}pp514-519.pdf).

- 1688 G. Chirici, R. Scotti, A. Montagni, A. Barbati, R. Cartisano, G. Lopez, M. Marchetti, R. E. Mcroberts,
 1689 H. Olsson, and P. Corona. Stochastic gradient boosting classification trees for forest fuel types mapping
 1690 through airborne laser scanning and IRS LISS-III imagery. *International Journal of Applied Earth*
 1691 *Observation and Geoinformation*, 25(1):87–97, 2013. ISSN 15698432. doi: 10.1016/j.jag.2013.04.006.
- 1692 Emilio Chuvieco, F. Javier Salas, Luis Carvacho, and Francisco Rodríguez-Silva. Integrated fire risk
 1693 mapping. In *Remote Sensing of Large Wildfires*, pages 61–100. Springer Berlin Heidelberg, Berlin,
 1694 Heidelberg, 1999. doi: 10.1007/978-3-642-60164-4_5. URL [http://link.springer.com/10.1007/](http://link.springer.com/10.1007/978-3-642-60164-4_5)
 1695 [978-3-642-60164-4_5](http://link.springer.com/10.1007/978-3-642-60164-4_5).
- 1696 Hamish Clarke, Rebecca Gibson, Brett Cirulis, Ross A Bradstock, and Trent D Penman. Developing
 1697 and testing models of the drivers of anthropogenic and lightning-caused wildfire ignitions in south-
 1698 eastern Australia. *Journal of Environmental Management*, 235:34–41, apr 2019. ISSN 0301-4797.
 1699 doi: 10.1016/J.JENVMAN.2019.01.055. URL [https://www-sciencedirect-com.login.ezproxy.](https://www-sciencedirect-com.login.ezproxy.library.ualberta.ca/science/article/pii/S0301479719300568)
 1700 [library.ualberta.ca/science/article/pii/S0301479719300568](https://www-sciencedirect-com.login.ezproxy.library.ualberta.ca/science/article/pii/S0301479719300568).
- 1701 J. Coen. Some Requirements for Simulating Wildland Fire Behavior Using Insight from Coupled
 1702 Weather—Wildland Fire Models. *Fire*, 1(1):6, feb 2018. ISSN 2571-6255. doi: 10.3390/fire1010006.
 1703 URL <http://www.mdpi.com/2571-6255/1/1/6>.
- 1704 Shane R. Coffield, Casey A. Graff, Yang Chen, Padhraic Smyth, Efi Foufoula-Georgiou, and James T.
 1705 Randerson. Machine learning to predict final fire size at the time of ignition. *International Journal of*
 1706 *Wildland Fire*, 28(11):861, 2019. ISSN 1049-8001. doi: 10.1071/WF19023. URL [http://www.publish.](http://www.publish.csiro.au/?paper=WF19023)
 1707 [csiro.au/?paper=WF19023](http://www.publish.csiro.au/?paper=WF19023).
- 1708 Judah Cohen, Dim Coumou, Jessica Hwang, Lester Mackey, Paulo Orenstein, Sonja Totz, and Eli Tziper-
 1709 man. S2S reboot: An argument for greater inclusion of machine learning in subseasonal to seasonal
 1710 forecasts. *Wiley Interdisciplinary Reviews: Climate Change*, 10(2), mar 2019. ISSN 1757-7780. doi:
 1711 10.1002/wcc.567. URL <https://onlinelibrary.wiley.com/doi/abs/10.1002/wcc.567>.
- 1712 L. Collins, P. Griffioen, G. Newell, and A. Mellor. The utility of Random Forests for wildfire severity
 1713 mapping. *Remote Sensing of Environment*, 216:374–384, oct 2018. ISSN 0034-4257. doi: 10.1016/J.RSE.
 1714 2018.07.005. URL <https://www.sciencedirect.com/science/article/pii/S0034425718303328>.
- 1715 Sean C.P. Coogan, François Nicolas Robinne, Piyush Jain, and Mike D. Flannigan. Scientists’ warning on
 1716 wildfire — a canadian perspective. *Canadian Journal of Forest Research*, 49(9):1015–1023, 2019. ISSN
 1717 12086037. doi: 10.1139/cjfr-2019-0094.
- 1718 Michelle Coppoletta, Kyle E. Merriam, and Brandon M. Collins. Post-fire vegetation and fuel development
 1719 influences fire severity patterns in reburns. *Ecological Applications*, 26(3):686–699, apr 2016. ISSN
 1720 1051-0761. doi: 10.1890/15-0225.
- 1721 A. Cordoba, R. Vilar, A. Lavrov, A. B. Utkin, and A. Fernandes. Multi-objective optimisation of lidar
 1722 parameters for forest-fire detection on the basis of a genetic algorithm. *Optics and Laser Technology*, 36
 1723 (5):393–400, 2004. ISSN 00303992. doi: 10.1016/j.optlastec.2003.10.010.
- 1724 Paulo Cortez and Anibal de Jesus Raimundo Morais. A data mining approach to predict forest fires using
 1725 meteorological data. 2007. URL <https://repositorium.sdum.uminho.pt/handle/1822/8039>.
- 1726 Sergi Costafreda-Aumedes, Adrian Cardil, Domingo M. Molina, Sarah N. Daniel, Robert Mavsar, and
 1727 Cristina Vega-Garcia. Analysis of factors influencing deployment of fire suppression resources in Spain
 1728 using artificial neural networks. *iForest*, 9(Feb 2016):138–145, 2015. ISSN 19717458. doi: 10.3832/
 1729 ifor1329-008.

- 1730 Michael A. Crimmins. Synoptic climatology of extreme fire-weather conditions across the southwest United
 1731 States. *International Journal of Climatology*, 26(8):1001–1016, jun 2006. ISSN 0899-8418. doi: 10.1002/
 1732 joc.1300. URL <http://doi.wiley.com/10.1002/joc.1300>.
- 1733 Morgan A. Crowley, Jeffrey A. Cardille, Joanne C. White, and Michael A. Wulder. Multi-sensor, multi-
 1734 scale, Bayesian data synthesis for mapping within-year wildfire progression. *Remote Sensing Letters*,
 1735 10(3):302–311, 2019. ISSN 2150-704X. doi: 10.1080/2150704X.2018.1536300. URL <https://www.tandfonline.com/doi/full/10.1080/2150704X.2018.1536300>.
- 1737 Thomas Curt, Laurent Borgniet, Thomas Ibanez, Vincent Moron, and Christelle Hély. Understanding fire
 1738 patterns and fire drivers for setting a sustainable management policy of the New-Caledonian biodiversity
 1739 hotspot. *Forest Ecology and Management*, 337:48–60, 2015. ISSN 03781127. doi: 10.1016/j.foreco.2014.
 1740 10.032. URL <http://dx.doi.org/10.1016/j.foreco.2014.10.032>.
- 1741 Thomas Curt, Thibaut Fréjaville, and Sébastien Lahaye. Modelling the spatial patterns of ignition causes
 1742 and fire regime features in southern France: Implications for fire prevention policy. *International Journal*
 1743 *of Wildland Fire*, 25(7):785–796, 2016. ISSN 10498001. doi: 10.1071/WF15205.
- 1744 James Richard Davis, J R L Hoare, and P M Nanninga. Developing a fire man-
 1745 agement expert system for Kakadu National Park, Australia. *Journal of Environmental*
 1746 *Management*, 1986. URL [https://publications.csiro.au/rpr/pub?list=BR0{&}pid=procite:
 1747 f221b911-2e97-4c9f-be9b-f1155bf48c24](https://publications.csiro.au/rpr/pub?list=BR0{&}pid=procite:f221b911-2e97-4c9f-be9b-f1155bf48c24).
- 1748 J.R. Davis, P.M. Nanninga, J.R.L. Hoare, and A.J. Press. Transferring scientific knowledge to natu-
 1749 ral resource managers using artificial intelligence concepts. *Ecological Modelling*, 46(1-2):73–89, jul
 1750 1989. ISSN 0304-3800. doi: 10.1016/0304-3800(89)90070-7. URL [https://www.sciencedirect.com/
 1751 science/article/pii/0304380089900707](https://www.sciencedirect.com/science/article/pii/0304380089900707).
- 1752 Raymond Davis, Zhiqiang Yang, Andrew Yost, Cole Belongie, and Warren Cohen. The normal fire en-
 1753 vironment—Modeling environmental suitability for large forest wildfires using past, present, and fu-
 1754 ture climate normals. *Forest Ecology and Management*, 390:173–186, apr 2017. ISSN 0378-1127.
 1755 doi: 10.1016/J.FORECO.2017.01.027. URL [https://www.sciencedirect.com/science/article/
 1756 pii/S0378112716309318](https://www.sciencedirect.com/science/article/pii/S0378112716309318).
- 1757 Antonella De Angelis, Carlo Ricotta, Marco Conedera, and Gianni Boris Pezzatti. Modelling the Meteor-
 1758 ological Forest Fire Niche in Heterogeneous Pyrologic Conditions. *PLOS ONE*, 10(2):e0116875, feb 2015.
 1759 ISSN 1932-6203. doi: 10.1371/journal.pone.0116875. URL [https://dx.plos.org/10.1371/journal.
 1760 pone.0116875](https://dx.plos.org/10.1371/journal.pone.0116875).
- 1761 P. P. de Bem, O. A. de Carvalho Júnior, E. A. T. Matricardi, R. F. Guimarães, and R. A. T. Gomes.
 1762 Predicting wildfire vulnerability using logistic regression and artificial neural networks: a case study in
 1763 Brazil’s Federal District. *International Journal of Wildland Fire*, 28(1):35, feb 2018. ISSN 1049-8001.
 1764 doi: 10.1071/wf18018. URL <http://www.publish.csiro.au/?paper=WF18018>.
- 1765 Marla Jose Perestrello De Vasconcelos, Sara Sllva, Margarlda Tome, Margarlda Alvim, Jose Mlguel, and
 1766 Cardoso Perelra. Spatial Prediction of Fire Ignition Probabilities: Comparing Logistic Regression and
 1767 Neural Networks. *Photogrammetric Engineering & Remote Sensing*, 67(1):73–81, 2001.
- 1768 Haifa Debouk, Ramon Riera-Tatché, and Cristina Vega-García. Assessing Post-Fire Regeneration in a
 1769 Mediterranean Mixed Forest Using Lidar Data and Artificial Neural Networks. *Photogrammetric Engi-
 1770 neering & Remote Sensing*, 2013. ISSN 00991112. doi: 10.14358/PERS.79.12.1121.
- 1771 Rosario Delgado, José-Luis González, Andrés Sotoca, and Xavier-Andoni Tibau. Archetypes of
 1772 Wildfire Arsonists: An Approach by Using Bayesian Networks. In *Forest Fire*. InTech, may

- 1773 2018. doi: 10.5772/intechopen.72615. URL [http://www.intechopen.com/books/forest-fire/
1774 archetypes-of-wildfire-arsonists-an-approach-by-using-bayesian-networks](http://www.intechopen.com/books/forest-fire/archetypes-of-wildfire-arsonists-an-approach-by-using-bayesian-networks).
- 1775 Mónica Denham and Karina Laneri. Using efficient parallelization in Graphic Processing Units to param-
1776 eterize stochastic fire propagation models. *Journal of Computational Science*, 25:76–88, 2018. ISSN
1777 18777503. doi: 10.1016/j.jocs.2018.02.007. URL <https://doi.org/10.1016/j.jocs.2018.02.007>.
- 1778 Mónica Denham, Kerstin Wendt, Germán Bianchini, Ana Cortés, and Tomàs Margalef. Dynamic Data-
1779 Driven Genetic Algorithm for forest fire spread prediction. *Journal of Computational Science*, 3(5):
1780 398–404, 2012. ISSN 18777503. doi: 10.1016/j.jocs.2012.06.002. URL [http://dx.doi.org/10.1016/j.
1781 jocs.2012.06.002](http://dx.doi.org/10.1016/j.jocs.2012.06.002).
- 1782 Peter J. Diggle, Raquel Menezes, and Ting-li Su. Geostatistical inference under preferential sam-
1783 pling. *Journal of the Royal Statistical Society: Series C (Applied Statistics)*, 59(2):191–232, mar 2010.
1784 ISSN 00359254. doi: 10.1111/j.1467-9876.2009.00701.x. URL [http://doi.wiley.com/10.1111/j.
1785 1467-9876.2009.00701.x](http://doi.wiley.com/10.1111/j.1467-9876.2009.00701.x).
- 1786 Luca Antonio Dimuccio, Rui Ferreira, Lucio Cunha, and Antonio Campar de Almeida. Regional forest-fire
1787 susceptibility analysis in central Portugal using a probabilistic ratings procedure and artificial neural
1788 network weights assignment. *International Journal of Wildland Fire*, 20(6):776, oct 2011. ISSN 1049-
1789 8001. doi: 10.1071/WF09083. URL <http://www.publish.csiro.au/?paper=WF09083>.
- 1790 T. L. Divya and M. N. Vijayalakshmi. Inference of Replanting in Forest Fire Affected Land Using Data
1791 Mining Technique. pages 121–129. Springer, New Delhi, 2016. doi: 10.1007/978-81-322-2734-2_13. URL
1792 http://link.springer.com/10.1007/978-81-322-2734-2_{_}13.
- 1793 Wisdom M. Dlamini. A Bayesian belief network analysis of factors influencing wildfire occurrence in
1794 Swaziland. *Environmental Modelling and Software*, 25(2):199–208, 2010. ISSN 13648152. doi: 10.1016/
1795 j.envsoft.2009.08.002. URL <http://dx.doi.org/10.1016/j.envsoft.2009.08.002>.
- 1796 Wisdom Mdumiseni Dlamini. Application of Bayesian networks for fire risk mapping using GIS and remote
1797 sensing data. *GeoJournal*, 76(3):283–296, jun 2011. ISSN 0343-2521. doi: 10.1007/s10708-010-9362-x.
1798 URL <http://link.springer.com/10.1007/s10708-010-9362-x>.
- 1799 E. Dragozi, I. Gitas, D. Stavrakoudis, and J. Theocharis. Burned Area Mapping Using Support Vector
1800 Machines and the FuzCoC Feature Selection Method on VHR IKONOS Imagery. *Remote Sensing*, 6
1801 (12):12005–12036, dec 2014. ISSN 2072-4292. doi: 10.3390/rs61212005. URL [http://www.mdpi.com/
1802 2072-4292/6/12/12005](http://www.mdpi.com/2072-4292/6/12/12005).
- 1803 Andrea Duane, Míriam Piqué, Marc Castellnou, and Lluís Brotons. Predictive modelling of fire occurrences
1804 from different fire spread patterns in Mediterranean landscapes. *International Journal of Wildland Fire*,
1805 24(3):407, jun 2015. ISSN 1049-8001. doi: 10.1071/WF14040. URL [http://www.publish.csiro.au/
1806 ?paper=WF14040](http://www.publish.csiro.au/?paper=WF14040).
- 1807 Ritaban Dutta, Jagannath Aryal, Aruneema Das, and Jamie B. Kirkpatrick. Deep cognitive imaging
1808 systems enable estimation of continental-scale fire incidence from climate data. *Scientific Reports*, 3,
1809 2013. ISSN 20452322. doi: 10.1038/srep03188.
- 1810 Ritaban Dutta, Aruneema Das, and Jagannath Aryal. Big data integration shows Australian bush-fire
1811 frequency is increasing significantly. *Royal Society Open Science*, 3(2):150241, feb 2016. ISSN 2054-5703.
1812 doi: 10.1098/rsos.150241. URL <https://royalsocietypublishing.org/doi/10.1098/rsos.150241>.
- 1813 Francis K. Dwomoh and Michael C. Wimberly. Fire regimes and their drivers in the Upper Guinean Region
1814 of West Africa. *Remote Sensing*, 9(11), 2017. ISSN 20724292. doi: 10.3390/rs9111117.

- 1815 J.B. Theocharis E. Dragozi, I. Z. Gitas, D.G. Stavrakoudis. A Performance Evaluation Of Sup-
 1816 port Vector Machines And The Nearest Neighbor Classifier In Classifying Image Objects For
 1817 Burned Area Mapping. In *Proceedings of the 8th International EARSeL FF-SIG Workshop*, Stresa,
 1818 Italy, 2011. URL [https://publications.europa.eu/en/publication-detail/-/publication/
 1819 5e90be79-efe8-43dd-8670-2518a43155f4/language-en](https://publications.europa.eu/en/publication-detail/-/publication/5e90be79-efe8-43dd-8670-2518a43155f4/language-en).
- 1820 Hamid Ebrahimi, Aliakbar Rasuly, *, and Davoud Mokhtari. Development of a Web GIS Sys-
 1821 tem Based on the MaxEnt Approach for Wildfire Management: A Case Study of East Azer-
 1822 baijan. *ECOPERSIA*, 5(3):1859–1873, 2017. URL [https://pdfs.semanticscholar.org/46b2/
 1823 fd74419232dbe2dedccaaca40bab6dbf50b8.pdf](https://pdfs.semanticscholar.org/46b2/fd74419232dbe2dedccaaca40bab6dbf50b8.pdf).
- 1824 J. Elith, J. R. Leathwick, and T. Hastie. A working guide to boosted regression trees. *Journal of Animal
 1825 Ecology*, 77(4):802–813, jul 2008. ISSN 0021-8790. doi: 10.1111/j.1365-2656.2008.01390.x. URL [http:
 1826 //doi.wiley.com/10.1111/j.1365-2656.2008.01390.x](http://doi.wiley.com/10.1111/j.1365-2656.2008.01390.x).
- 1827 Jane Elith, Steven J. Phillips, Trevor Hastie, Miroslav Dudík, Yung En Chee, and Colin J. Yates. A
 1828 statistical explanation of MaxEnt for ecologists. *Diversity and Distributions*, 17(1):43–57, jan 2011.
 1829 ISSN 13669516. doi: 10.1111/j.1472-4642.2010.00725.x. URL [http://doi.wiley.com/10.1111/j.
 1830 1472-4642.2010.00725.x](http://doi.wiley.com/10.1111/j.1472-4642.2010.00725.x).
- 1831 Thomas A. Fairman, Lauren T. Bennett, Shauna Tupper, and Craig R. Nitschke. Frequent wildfires erode
 1832 tree persistence and alter stand structure and initial composition of a fire-tolerant sub-alpine forest.
 1833 *Journal of Vegetation Science*, 28(6):1151–1165, 2017. ISSN 16541103. doi: 10.1111/jvs.12575.
- 1834 Lei Fang, Jian Yang, Megan White, and Zhihua Liu. Predicting potential fire severity using vegetation,
 1835 topography and surface moisture availability in a Eurasian boreal forest landscape. *Forests*, 9(3):1–26,
 1836 2018. ISSN 19994907. doi: 10.3390/f9030130.
- 1837 Armando M. Fernandes, Andrei B. Utkin, Alexander V. Lavrov, and Rui M. Vilar. Neural Network Based
 1838 Recognition of Smoke Signatures from Lidar Signals. *Neural Processing Letters*, 19(3):175–189, jun
 1839 2004a. ISSN 1370-4621. doi: 10.1023/B:NEPL.0000035598.19042.42. URL [http://link.springer.
 1840 com/10.1023/B:NEPL.0000035598.19042.42](http://link.springer.com/10.1023/B:NEPL.0000035598.19042.42).
- 1841 Armando M. Fernandes, Andrei B. Utkin, Alexander V. Lavrov, and Rui M. Vilar. Development of neural
 1842 network committee machines for automatic forest fire detection using lidar. *Pattern Recognition*, 37(10):
 1843 2039–2047, oct 2004b. ISSN 00313203. doi: 10.1016/j.patcog.2004.04.002. URL [http://linkinghub.
 1844 elsevier.com/retrieve/pii/S0031320304001360](http://linkinghub.elsevier.com/retrieve/pii/S0031320304001360).
- 1845 Paulo M. Fernandes, Tiago Monteiro-Henriques, Nuno Guiomar, Carlos Loureiro, and Ana M.G. Barros.
 1846 Bottom-Up Variables Govern Large-Fire Size in Portugal. *Ecosystems*, 19(8):1362–1375, 2016. ISSN
 1847 14350629. doi: 10.1007/s10021-016-0010-2.
- 1848 Alfonso Fernandez-Manso, Carmen Quintano, and Dar A. Roberts. Burn severity analysis in Mediterranean
 1849 forests using maximum entropy model trained with EO-1 Hyperion and LiDAR data. *ISPRS Journal of
 1850 Photogrammetry and Remote Sensing*, 155(July):102–118, 2019. ISSN 09242716. doi: 10.1016/j.isprsjprs.
 1851 2019.07.003. URL <https://doi.org/10.1016/j.isprsjprs.2019.07.003>.
- 1852 Mark A Finney. FARSITE: Fire Area Simulator-Model Development and Evaluation. Technical report,
 1853 Rocky Mountain Research Station, 2004. URL <http://www.farsite.org>.
- 1854 Mark A. Finney. The challenge of quantitative risk analysis for wildland fire. In *Forest Ecology and
 1855 Management*, volume 211, pages 97–108, jun 2005. doi: 10.1016/j.foreco.2005.02.010.

- 1856 Marisa G. Fonseca, Luiz Eduardo O. C. Aragão, André Lima, Yosio E. Shimabukuro, Egidio Arai, and
 1857 Liana O. Anderson. Modelling fire probability in the Brazilian Amazon using the maximum entropy
 1858 method. *International Journal of Wildland Fire*, 25(9):955, sep 2016. doi: 10.1071/WF15216. URL
 1859 <http://www.publish.csiro.au/?paper=WF15216>.
- 1860 Matthias Forkel, Wouter Dorigo, Gitta Lasslop, Irene Teubner, Emilio Chuvieco, and Kirsten Thonicke. A
 1861 data-driven approach to identify controls on global fire activity from satellite and climate observations
 1862 (SOFIA V1). *Geoscientific Model Development*, 10(12):4443–4476, dec 2017. ISSN 1991-9603. doi:
 1863 10.5194/gmd-10-4443-2017. URL <https://www.geosci-model-dev.net/10/4443/2017/>.
- 1864 Matthias Forkel, Niels Andela, Sandy P Harrison, Gitta Lasslop, Margreet Van Marle, Emilio Chuvieco,
 1865 Wouter Dorigo, Matthew Forrest, Stijn Hantson, Angelika Heil, Fang Li, Joe Melton, Stephen Sitch,
 1866 Chao Yue, and Almut Arneth. Emergent relationships with respect to burned area in global satellite
 1867 observations and fire-enabled vegetation models. *Biogeosciences*, 16(1):57–76, 2019. ISSN 17264189. doi:
 1868 10.5194/bg-16-57-2019.
- 1869 Yoav Freund and Robert E Shapire. A decision-theoretic generalization of on-line learning and an ap-
 1870 plication to boosting. In *Computational Learning Theory: Eurocolt '95*, pages 23–37. Springer-Verlag,
 1871 1995.
- 1872 Jerome H. Friedman. Greedy function approximation: A gradient boosting machine. *Ann. Statist.*, 29(5):
 1873 1189–1232, 10 2001. doi: 10.1214/aos/1013203451. URL <https://doi.org/10.1214/aos/1013203451>.
- 1874 Sigfredo Fuentes, Eden Jane Tongson, Roberta De Bei, Claudia Gonzalez Viejo, Renata Ristic, Stephen
 1875 Tyerman, and Kerry Wilkinson. Non-Invasive Tools to Detect Smoke Contamination in Grapevine
 1876 Canopies, Berries and Wine: A Remote Sensing and Machine Learning Modeling Approach. *Sensors*, 19
 1877 (15):3335, jul 2019. ISSN 1424-8220. doi: 10.3390/s19153335. URL [https://www.mdpi.com/1424-8220/
 1878 19/15/3335](https://www.mdpi.com/1424-8220/19/15/3335).
- 1879 Mariano García, David Riaño, Emilio Chuvieco, Javier Salas, and F. Mark Danson. Multispectral and
 1880 LiDAR data fusion for fuel type mapping using Support Vector Machine and decision rules. *Remote
 1881 Sensing of Environment*, 115(6):1369–1379, 2011. ISSN 00344257. doi: 10.1016/j.rse.2011.01.017. URL
 1882 <http://dx.doi.org/10.1016/j.rse.2011.01.017>.
- 1883 Paula García-Llamas, Susana Suárez-Seoane, Angela Taboada, Alfonso Fernández-Manso, Carmen Quin-
 1884 tano, Víctor Fernández-García, José Manuel Fernández-Guisuraga, Elena Marcos, and Leonor Calvo.
 1885 Environmental drivers of fire severity in extreme fire events that affect Mediterranean pine forest
 1886 ecosystems. *Forest Ecology and Management*, 433(October 2018):24–32, 2019. ISSN 03781127. doi:
 1887 10.1016/j.foreco.2018.10.051. URL <https://doi.org/10.1016/j.foreco.2018.10.051>.
- 1888 Stuart Geman, Elie Bienenstock, and René Doursat. Neural Networks and the Bias/Variance Dilemma.
 1889 *Neural Computation*, 4(1):1–58, jan 1992. ISSN 0899-7667. doi: 10.1162/neco.1992.4.1.1.
- 1890 Andre Gensler, Janosch Henze, Bernhard Sick, and Nils Raabe. Deep Learning for solar power forecasting
 1891 - An approach using AutoEncoder and LSTM Neural Networks. In *2016 IEEE International Conference
 1892 on Systems, Man, and Cybernetics, SMC 2016 - Conference Proceedings*, pages 2858–2865. Institute
 1893 of Electrical and Electronics Engineers Inc., feb 2017. ISBN 9781509018970. doi: 10.1109/SMC.2016.
 1894 7844673.
- 1895 Omid Ghorbanzadeh, Thomas Blaschke, Khalil Gholamnia, and Jagannath Aryal. Forest Fire Susceptibility
 1896 and Risk Mapping Using Social/Infrastructural Vulnerability and Environmental Variables. *Fire*, 2(3):
 1897 50, sep 2019a. ISSN 2571-6255. doi: 10.3390/fire2030050. URL [https://www.mdpi.com/2571-6255/2/
 1898 3/50](https://www.mdpi.com/2571-6255/2/3/50).

- 1899 Omid Ghorbanzadeh, Khalil Valizadeh Kamran, Thomas Blaschke, Jagannath Aryal, Amin Naboureh,
1900 Jamshid Einali, and Jinhu Bian. Spatial Prediction of Wildfire Susceptibility Using Field Survey GPS
1901 Data and Machine Learning Approaches. *Fire*, 2(3):43, jul 2019b. ISSN 2571-6255. doi: 10.3390/
1902 fire2030043. URL <https://www.mdpi.com/2571-6255/2/3/43>.
- 1903 Louis Giglio, Luigi Boschetti, David P. Roy, Michael L. Humber, and Christopher O. Justice. The Collection
1904 6 MODIS burned area mapping algorithm and product. *Remote Sensing of Environment*, 217:72–85,
1905 nov 2018. ISSN 00344257. doi: 10.1016/j.rse.2018.08.005.
- 1906 Ljubomir Gigović, Hamid Reza Pourghasemi, Siniša Drobnjak, and Shibiao Bai. Testing a New Ensemble
1907 Model Based on SVM and Random Forest in Forest Fire Susceptibility Assessment and Its Mapping in
1908 Serbia’s Tara National Park. *Forests*, 10(5):408, may 2019. ISSN 1999-4907. doi: 10.3390/f10050408.
1909 URL <https://www.mdpi.com/1999-4907/10/5/408>.
- 1910 Y.J. Goldarag, Ali Mohammadzadeh, and A. S. Ardakani. Fire Risk Assessment Using Neural Network
1911 and Logistic Regression. *Journal of the Indian Society of Remote Sensing*, 44(6):885–894, 2016. ISSN
1912 09743006. doi: 10.1007/s12524-016-0557-6. URL <http://dx.doi.org/10.1007/s12524-016-0557-6>.
- 1913 Carla Gomes. Computational Sustainability: Computational Methods for a Sustainable Environment,
1914 Economy, and Society. *The Bridge, National Academy of Engineering*, 39(4), 2009.
- 1915 Israel Gómez and M. Pilar Martín. Prototyping an artificial neural network for burned area mapping on
1916 a regional scale in Mediterranean areas using MODIS images. *International Journal of Applied Earth
1917 Observation and Geoinformation*, 13(5):741–752, 2011. ISSN 15698432. doi: 10.1016/j.jag.2011.05.002.
- 1918 Noel Gorelick, Matt Hancher, Mike Dixon, Simon Ilyushchenko, David Thau, and Rebecca Moore. Google
1919 Earth Engine: Planetary-scale geospatial analysis for everyone. *Remote Sensing of Environment*, 2017.
1920 doi: 10.1016/j.rse.2017.06.031. URL <https://doi.org/10.1016/j.rse.2017.06.031>.
- 1921 Futao Guo, Guangyu Wang, Zhangwen Su, Huiling Liang, Wenhui Wang, Fangfang Lin, and Aiqin Liu.
1922 What drives forest fire in Fujian, China? Evidence from logistic regression and Random Forests. *In-
1923 ternational Journal of Wildland Fire*, 25(5):505, may 2016a. ISSN 1049-8001. doi: 10.1071/WF15121.
1924 URL <http://www.publish.csiro.au/?paper=WF15121>.
- 1925 Futao Guo, Lianjun Zhang, Sen Jin, Muluaem Tigabu, Zhangwen Su, Wenhui Wang, Futao Guo, Lianjun
1926 Zhang, Sen Jin, Muluaem Tigabu, Zhangwen Su, and Wenhui Wang. Modeling Anthropogenic Fire
1927 Occurrence in the Boreal Forest of China Using Logistic Regression and Random Forests. *Forests*, 7(12):
1928 250, oct 2016b. ISSN 1999-4907. doi: 10.3390/f7110250. URL [http://www.mdpi.com/1999-4907/7/
1929 11/250](http://www.mdpi.com/1999-4907/7/11/250).
- 1930 Dale Hamilton, Barry Myers, and Jonathan Branham. Evaluation Of Texture As An Input Of Spatial
1931 Context For Machine Learning Mapping Of Wildland Fire Effects. *An International Journal (SIPIJ)*, 8
1932 (5), 2017. doi: 10.5121/sipij.2017.8501.
- 1933 Jie Han, Zehao Shen, Lingxiao Ying, Guixiang Li, and Anping Chen. Early post-fire regeneration of
1934 a fire-prone subtropical mixed Yunnan pine forest in Southwest China: Effects of pre-fire vegetation,
1935 fire severity and topographic factors. *Forest Ecology and Management*, 356(2015):31–40, 2015. ISSN
1936 03781127. doi: 10.1016/j.foreco.2015.06.016. URL [http://dx.doi.org/10.1016/j.foreco.2015.06.
1937 016](http://dx.doi.org/10.1016/j.foreco.2015.06.016).
- 1938 Lucas Harris and Alan H. Taylor. Previous burns and topography limit and reinforce fire severity in a
1939 large wildfire. *Ecosphere*, 8(11), 2017. ISSN 21508925. doi: 10.1002/ecs2.2019.
- 1940 Trevor Hastie, Jerome Friedman, and Robert Tibshirani. *The Elements of Statistical Learning: Data
1941 Mining, Inference, and Prediction*. Springer, New York, NY, 2009. URL [https://doi.org/10.1007/
1942 978-0-387-21606-5](https://doi.org/10.1007/978-0-387-21606-5).

- 1943 Todd J. Hawbaker, Melanie K. Vanderhoof, Yen Ju Beal, Joshua D. Takacs, Gail L. Schmidt, Jeff T.
1944 Falgout, Brad Williams, Nicole M. Fairaux, Megan K. Caldwell, Joshua J. Picotte, Stephen M. Howard,
1945 Susan Stitt, and John L. Dwyer. Mapping burned areas using dense time-series of Landsat data. *Remote*
1946 *Sensing of Environment*, 198:504–522, sep 2017. ISSN 00344257. doi: 10.1016/j.rse.2017.06.027.
- 1947 Marti A. Hearst, Susan T Dumais, Edgar Osuna, John Platt, and Bernhard Scholkopf. Support vector
1948 machines. *IEEE Intelligent Systems and their applications*, 13(4):18–28, 1998.
- 1949 Robert Hecht-Nielsen. Theory of the backpropagation neural network. In *Neural networks for perception*,
1950 pages 65–93. Elsevier, 1992.
- 1951 Risto K. Heikkinen, Mathieu Marmion, and Miska Luoto. Does the interpolation accuracy of species
1952 distribution models come at the expense of transferability? *Ecography*, 35(3):276–288, mar 2012.
1953 ISSN 09067590. doi: 10.1111/j.1600-0587.2011.06999.x. URL [http://doi.wiley.com/10.1111/j.](http://doi.wiley.com/10.1111/j.1600-0587.2011.06999.x)
1954 [1600-0587.2011.06999.x](http://doi.wiley.com/10.1111/j.1600-0587.2011.06999.x).
- 1955 Txomin Hermosilla, Michael A. Wulder, Joanne C. White, Nicholas C. Coops, and Geordie W. Hobart.
1956 Regional detection, characterization, and attribution of annual forest change from 1984 to 2012 using
1957 Landsat-derived time-series metrics. *Remote Sensing of Environment*, 170:121–132, 2015. ISSN 00344257.
1958 doi: 10.1016/j.rse.2015.09.004. URL <http://dx.doi.org/10.1016/j.rse.2015.09.004>.
- 1959 Sepp Hochreiter and Jürgen Schmidhuber. Long short-term memory. *Neural computation*, 9(8):1735–1780,
1960 1997.
- 1961 Jonathan L. Hodges and Brian Y. Lattimer. Wildland Fire Spread Modeling Using Convolutional Neural
1962 Networks. *Fire Technology*, nov 2019. ISSN 15728099. doi: 10.1007/s10694-019-00846-4.
- 1963 C. M. Hoffman, J. Canfield, R. R. Linn, W. Mell, C. H. Sieg, F. Pimont, and J. Ziegler. Evaluating Crown
1964 Fire Rate of Spread Predictions from Physics-Based Models. *Fire Technology*, 52(1):221–237, jan 2016.
1965 ISSN 15728099. doi: 10.1007/s10694-015-0500-3.
- 1966 Zachary A. Holden, Penelope Morgan, and Jeffrey S. Evans. A predictive model of burn severity based on
1967 20-year satellite-inferred burn severity data in a large southwestern US wilderness area. *Forest Ecology*
1968 *and Management*, 258(11):2399–2406, 2009. ISSN 03781127. doi: 10.1016/j.foreco.2009.08.017.
- 1969 Baisravan Homchaudhuri, Sheng Zhao, Kelly Cohen, and Manish Kumar. Generation of optimal fire-line
1970 for fighting wildland fires using genetic algorithms. *Proceedings of the ASME Dynamic Systems and*
1971 *Control Conference 2009, DSCC2009*, (PART A):111–118, 2010. doi: 10.1115/DSCC2009-2707.
- 1972 Haoyuan Hong, Paraskevas Tsangaratos, Ioanna Iliia, Junzhi Liu, A-Xing Zhu, and Chong Xu. Applying
1973 genetic algorithms to set the optimal combination of forest fire related variables and model forest fire
1974 susceptibility based on data mining models. The case of Dayu County, China. *Science of The Total*
1975 *Environment*, 630:1044–1056, jul 2018. ISSN 0048-9697. doi: 10.1016/J.SCITOTENV.2018.02.278.
1976 URL <https://www.sciencedirect.com/science/article/pii/S004896971830679X>.
- 1977 F M Anim Hossain, Youmin Zhang, Chi Yuan, and Chun-Yi Su. Wildfire Flame and Smoke Detection Using
1978 Static Image Features and Artificial Neural Network. In *2019 1st International Conference on Industrial*
1979 *Artificial Intelligence (IAI)*, pages 1–6. Institute of Electrical and Electronics Engineers (IEEE), oct
1980 2019. doi: 10.1109/iciai.2019.8850811.
- 1981 Bronwyn A. Hradsky, Trent D. Penman, Dan Ababei, Anca Hanea, Euan G. Ritchie, Alan York, and
1982 Julian Di Stefano. Bayesian networks elucidate interactions between fire and other drivers of terrestrial
1983 fauna distributions. *Ecosphere*, 8(8):e01926, aug 2017. ISSN 21508925. doi: 10.1002/ecs2.1926. URL
1984 <http://doi.wiley.com/10.1002/ecs2.1926>.

- 1985 Carolynne Hultquist, Gang Chen, and Kaiguang Zhao. A comparison of Gaussian process regression,
 1986 random forests and support vector regression for burn severity assessment in diseased forests. *Re-*
 1987 *remote Sensing Letters*, 5(8):723–732, aug 2014. ISSN 2150-704X. doi: 10.1080/2150704X.2014.963733.
 1988 URL <http://dx.doi.org/10.1080/2150704X.2014.963733>[http://www.tandfonline.com/doi/abs/](http://www.tandfonline.com/doi/abs/10.1080/2150704X.2014.963733)
 1989 [10.1080/2150704X.2014.963733](http://www.tandfonline.com/doi/abs/10.1080/2150704X.2014.963733).
- 1990 L.S. Iliadis. A decision support system applying an integrated fuzzy model for long-term forest fire
 1991 risk estimation. *Environmental Modelling & Software*, 20(5):613–621, may 2005. ISSN 1364-8152.
 1992 doi: 10.1016/J.ENVSOFT.2004.03.006. URL [https://www.sciencedirect.com/science/article/](https://www.sciencedirect.com/science/article/pii/S1364815204000933)
 1993 [pii/S1364815204000933](https://www.sciencedirect.com/science/article/pii/S1364815204000933).
- 1994 Abolfazl Jaafari. Factors Influencing Regional-Scale Wildfire Probability in Iran: An Applica-
 1995 tion of Random Forest and Support Vector Machine. *Spatial Modeling in GIS and R for*
 1996 *Earth and Environmental Sciences*, pages 607–619, jan 2019. doi: 10.1016/B978-0-12-815226-3.
 1997 00028-4. URL [https://www-sciencedirect-com.login.ezproxy.library.ualberta.ca/science/](https://www-sciencedirect-com.login.ezproxy.library.ualberta.ca/science/article/pii/B9780128152263000284)
 1998 [article/pii/B9780128152263000284](https://www-sciencedirect-com.login.ezproxy.library.ualberta.ca/science/article/pii/B9780128152263000284).
- 1999 Abolfazl Jaafari, Eric K. Zenner, and Binh Thai Pham. Wildfire spatial pattern analysis in the Za-
 2000 gros Mountains, Iran: A comparative study of decision tree based classifiers. *Ecological Informat-*
 2001 *ics*, 43:200–211, jan 2018. ISSN 1574-9541. doi: 10.1016/J.ECOINF.2017.12.006. URL [https://](https://www.sciencedirect.com/science/article/pii/S157495411730167X)
 2002 www.sciencedirect.com/science/article/pii/S157495411730167X.
- 2003 Abolfazl Jaafari, Eric K. Zenner, Mahdi Panahi, and Himan Shahabi. Hybrid artificial intelligence mod-
 2004 els based on a neuro-fuzzy system and metaheuristic optimization algorithms for spatial prediction of
 2005 wildfire probability. *Agricultural and Forest Meteorology*, 266-267:198–207, mar 2019. ISSN 01681923.
 2006 doi: 10.1016/j.agrformet.2018.12.015. URL [https://www.sciencedirect.com/science/article/pii/](https://www.sciencedirect.com/science/article/pii/S0168192318304088)
 2007 [S0168192318304088](https://www.sciencedirect.com/science/article/pii/S0168192318304088).
- 2008 Jacek Jakubowski, Maciej Solarczyk, and Michał Wiśnios. Smoke detection in a digital image with the use
 2009 of convolutional network. page 14. *SPIE-Intl Soc Optical Eng*, mar 2019. ISBN 9781510627857. doi:
 2010 10.1117/12.2524560.
- 2011 J. . R. Jang. Anfis: adaptive-network-based fuzzy inference system. *IEEE Transactions on Systems, Man,*
 2012 *and Cybernetics*, 23(3):665–685, May 1993. doi: 10.1109/21.256541.
- 2013 Maria João Sousa, Alexandra Moutinho, and Miguel Almeida. Wildfire detection using transfer learning
 2014 on augmented datasets. *Expert Systems with Applications*, page 112975, sep 2019. ISSN 09574174. doi:
 2015 10.1016/j.eswa.2019.112975.
- 2016 Torres João, Gonçalves João, Marcos Bruno, and Honrado João. Indicator-based assessment of post-fire re-
 2017 covery dynamics using satellite NDVI time-series. *Ecological Indicators*, 89(January):199–212, 2018. ISSN
 2018 1470160X. doi: 10.1016/j.ecolind.2018.02.008. URL [https://doi.org/10.1016/j.ecolind.2018.02.](https://doi.org/10.1016/j.ecolind.2018.02.008)
 2019 [008](https://doi.org/10.1016/j.ecolind.2018.02.008).
- 2020 Jill F. Johnstone, Teresa N. Hollingsworth, F. Stuart Chapin, and Michelle C. Mack. Changes in fire regime
 2021 break the legacy lock on successional trajectories in Alaskan boreal forest. *Global Change Biology*, 16
 2022 (4):1281–1295, 2010. ISSN 13541013. doi: 10.1111/j.1365-2486.2009.02051.x.
- 2023 Kyle D. Julian and Mykel J. Kochenderfer. Autonomous distributed wildfire surveillance using deep
 2024 reinforcement learning. In *AIAA Guidance, Navigation, and Control Conference, 2018*, number 210039.
 2025 American Institute of Aeronautics and Astronautics Inc, AIAA, jan 2018a. ISBN 9781624105265. doi:
 2026 10.2514/6.2018-1589.

- 2027 Kyle D. Julian and Mykel J. Kochenderfer. Distributed Wildfire Surveillance with Autonomous Aircraft
 2028 using Deep Reinforcement Learning. *Journal of Guidance, Control, and Dynamics*, 42(8):1768–1778, oct
 2029 2018b. URL <http://arxiv.org/abs/1810.04244>.
- 2030 Martin Jung, Susanne Tautenhahn, Christian Wirth, and Jens Kattge. Estimating Basal Area of Spruce
 2031 and Fir in Post-fire Residual Stands in Central Siberia Using Quickbird, Feature Selection, and Random
 2032 Forests. *Procedia Computer Science*, 18:2386–2395, jan 2013. ISSN 1877-0509. doi: 10.1016/J.PROCS.
 2033 2013.05.410. URL <https://www.sciencedirect.com/science/article/pii/S187705091300553X>.
- 2034 M. Njoki Kahi and N. P. Hanan. Fire in sub-Saharan Africa: The fuel, cure and connectivity hypothesis.
 2035 *Global Ecology and Biogeography*, 27(8):946–957, aug 2018. ISSN 1466822X. doi: 10.1111/geb.12753.
 2036 URL <http://doi.wiley.com/10.1111/geb.12753>.
- 2037 Van R. Kane, C. Alina Cansler, Nicholas A. Povak, Jonathan T. Kane, Robert J. McGaughey, James A.
 2038 Lutz, Derek J. Churchill, and Malcolm P. North. Mixed severity fire effects within the Rim fire:
 2039 Relative importance of local climate, fire weather, topography, and forest structure. *Forest Ecology
 2040 and Management*, 358:62–79, dec 2015. ISSN 0378-1127. doi: 10.1016/J.FORECO.2015.09.001. URL
 2041 <https://www.sciencedirect.com/science/article/pii/S0378112715004697>.
- 2042 Anuj Karpatne, Imme Ebert-Uphoff, Sai Ravela, Hassan Ali Babaie, and Vipin Kumar. Machine Learning
 2043 for the Geosciences: Challenges and Opportunities. nov 2017. URL [http://arxiv.org/abs/1711.
 2044 04708](http://arxiv.org/abs/1711.04708).
- 2045 Robert E. Keane, Geoffrey J. Cary, Ian D. Davies, Michael D. Flannigan, Robert H. Gardner, Sandra
 2046 Lavorel, James M. Lenihan, Chao Li, and T. Scott Rupp. A classification of landscape fire succession
 2047 models: Spatial simulations of fire and vegetation dynamics. *Ecological Modelling*, 179(1-2):3–27, nov
 2048 2004. ISSN 03043800. doi: 10.1016/j.ecolmodel.2004.03.015.
- 2049 Nima Khakzad. Modeling wildfire spread in wildland-industrial interfaces using dynamic Bayesian network.
 2050 *Reliability Engineering & System Safety*, 189:165–176, sep 2019. ISSN 0951-8320. doi: 10.1016/J.RESS.
 2051 2019.04.006. URL <https://www.sciencedirect.com/science/article/pii/S0951832018313887>.
- 2052 Sea Jin Kim, Chul-Hee Lim, Gang Sun Kim, Jongyeol Lee, Tobias Geiger, Omid Rahmati, Yowhan Son,
 2053 and Woo-Kyun Lee. Multi-Temporal Analysis of Forest Fire Probability Using Socio-Economic and
 2054 Environmental Variables. *Remote Sensing*, 11(1):86, jan 2019. ISSN 2072-4292. doi: 10.3390/rs11010086.
 2055 URL <https://www.mdpi.com/2072-4292/11/1/86>.
- 2056 Seongchan Kim, Seungkyun Hong, Minsu Joh, and Sa-kwang Song. DeepRain: ConvLSTM Network for
 2057 Precipitation Prediction using Multichannel Radar Data. nov 2017. URL [http://arxiv.org/abs/
 2058 1711.02316](http://arxiv.org/abs/1711.02316).
- 2059 ByoungChul Ko, Kwang-Ho Cheong, and Jae-Yeal Nam. Early fire detection algorithm based on irreg-
 2060 ular patterns of flames and hierarchical Bayesian Networks. *Fire Safety Journal*, 45(4):262–270, jun
 2061 2010. ISSN 0379-7112. doi: 10.1016/J.FIRESAF.2010.04.001. URL [https://www.sciencedirect.com/
 2062 science/article/pii/S0379711210000378](https://www.sciencedirect.com/science/article/pii/S0379711210000378).
- 2063 Alexandru Korotcov, Valery Tkachenko, Daniel P. Russo, and Sean Ekins. Comparison of Deep Learning
 2064 with Multiple Machine Learning Methods and Metrics Using Diverse Drug Discovery Data Sets. *Molec-
 2065 ular Pharmaceutics*, 14(12):4462–4475, dec 2017. ISSN 15438392. doi: 10.1021/acs.molpharmaceut.
 2066 7b00578.
- 2067 Peter Kourtz. Artificial intelligence: a new tool for forest management. *Canadian Journal of For-
 2068 est Research*, 20(4):428–437, apr 1990. ISSN 0045-5067. doi: 10.1139/x90-060. URL [http://www.
 2069 nrcresearchpress.com/doi/10.1139/x90-060](http://www.nrcresearchpress.com/doi/10.1139/x90-060).

- 2070 P.H. Kourtz. Artificial intelligence applications in the next generation Canadian forest fire control system,
2071 1993. URL <https://cfs.nrcan.gc.ca/publications?id=10775>.
- 2072 V. I. Kozik, E. S. Nezhevenko, and A. S. Feoktistov. Adaptive prediction of forest fire behavior on the basis
2073 of recurrent neural networks. *Optoelectronics, Instrumentation and Data Processing*, 49(3):250–259, may
2074 2013. ISSN 8756-6990. doi: 10.3103/S8756699013030060. URL [http://link.springer.com/10.3103/
2075 S8756699013030060](http://link.springer.com/10.3103/S8756699013030060).
- 2076 V. I. Kozik, E. S. Nezhevenko, and A. S. Feoktistov. Studying the method of adaptive prediction of
2077 forest fire evolution on the basis of recurrent neural networks. *Optoelectronics, Instrumentation and
2078 Data Processing*, 50(4):395–401, jul 2014. ISSN 8756-6990. doi: 10.3103/S8756699014040116. URL
2079 <http://link.springer.com/10.3103/S8756699014040116>.
- 2080 Max Kuhn and Kjell Johnson. *Applied predictive modeling*. Springer New York, jan 2013. ISBN
2081 9781461468493. doi: 10.1007/978-1-4614-6849-3.
- 2082 Jan Kukačka, Vladimir Golkov, and Daniel Cremers. Regularization for Deep Learning: A Taxonomy. oct
2083 2017. URL <http://arxiv.org/abs/1710.10686>.
- 2084 Nataliia Kussul, Mykola Lavreniuk, Sergii Skakun, and Andrii Shelestov. Deep Learning Classification
2085 of Land Cover and Crop Types Using Remote Sensing Data. *IEEE Geoscience and Remote Sensing
2086 Letters*, 14(5):778–782, may 2017. ISSN 1545-598X. doi: 10.1109/LGRS.2017.2681128. URL [http:
2087 //ieeexplore.ieee.org/document/7891032/](http://ieeexplore.ieee.org/document/7891032/).
- 2088 Ryan Lagerquist, Mike D. Flannigan, Xianli Wang, and Ginny A. Marshall. Automated prediction of
2089 extreme fire weather from synoptic patterns in northern Alberta, Canada. *Canadian Journal of Forest
2090 Research*, 47(9):1175–1183, sep 2017. ISSN 0045-5067. doi: 10.1139/cjfr-2017-0063. URL [http://www.
2091 nrcresearchpress.com/doi/10.1139/cjfr-2017-0063](http://www.nrcresearchpress.com/doi/10.1139/cjfr-2017-0063).
- 2092 Zachary Langford, Jitendra Kumar, and Forrest Hoffman. Wildfire mapping in interior alaska using deep
2093 neural networks on imbalanced datasets. In *IEEE International Conference on Data Mining Workshops,
2094 ICDMW*, volume 2018-Novem, pages 770–778. IEEE Computer Society, feb 2019. ISBN 9781538692882.
2095 doi: 10.1109/ICDMW.2018.00116.
- 2096 David J. Lary, Amir H. Alavi, Amir H. Gandomi, and Annette L. Walker. Machine learning in geosciences
2097 and remote sensing. *Geoscience Frontiers*, 7(1):3–10, jan 2016. ISSN 16749871. doi: 10.1016/j.gsf.2015.
2098 07.003. URL <https://www.sciencedirect.com/science/article/pii/S1674987115000821>.
- 2099 Don J Latham. Artificial Intelligence Applications To Fire Management. In *Proceedings Of The Symposium
2100 On Wildland Fire*, South Lake Tahoe, 1987.
- 2101 Christopher J. Lauer, Claire A. Montgomery, and Thomas G. Dieterich. Spatial interactions and opti-
2102 mal forest management on a fire-threatened landscape. *Forest Policy and Economics*, 83:107–120, oct
2103 2017. ISSN 1389-9341. doi: 10.1016/J.FORPOL.2017.07.006. URL [https://www.sciencedirect.com/
2104 science/article/pii/S1389934116304749](https://www.sciencedirect.com/science/article/pii/S1389934116304749).
- 2105 Yann LeCun, Yoshua Bengio, and Geoffrey Hinton. Deep learning. *nature*, 521(7553):436, 2015.
- 2106 Michael Leuenberger, Joana Parente, Marj Tonini, Mário Gonzalez Pereira, and Mikhail Kanevski. Wildfire
2107 susceptibility mapping: Deterministic vs. stochastic approaches. *Environmental Modelling & Software*,
2108 101:194–203, mar 2018. ISSN 1364-8152. doi: 10.1016/J.ENVSOFT.2017.12.019. URL [https://www.
2109 sciencedirect.com/science/article/pii/S1364815217303316](https://www.sciencedirect.com/science/article/pii/S1364815217303316).

- 2110 Danielle Levac, Heather Colquhoun, and Kelly K O'Brien. Scoping studies: advancing
 2111 the methodology. *Implementation science : IS*, 5:69, sep 2010. ISSN 1748-5908.
 2112 doi: 10.1186/1748-5908-5-69. URL <http://www.ncbi.nlm.nih.gov/pubmed/20854677><http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=PMC2954944>.
 2113
- 2114 Berangere A. Leys, Julie L. Commerford, and Kendra K. McLauchlan. Reconstructing grassland fire history
 2115 using sedimentary charcoal: Considering count, size and shape. *PLOS ONE*, 12(4):e0176445, apr 2017.
 2116 ISSN 1932-6203. doi: 10.1371/journal.pone.0176445. URL <https://dx.plos.org/10.1371/journal.pone.0176445>.
 2117
- 2118 Hanchao Li, Xiang Fei, and Chaobo He. Study on Most Important Factor and Most Vulnerable Location
 2119 for a Forest Fire Case Using Various Machine Learning Techniques. *2018 Sixth International Conference
 2120 on Advanced Cloud and Big Data (CBD)*, pages 298–303, 2018a. doi: 10.1109/CBD.2018.00060. URL
 2121 <https://ieeexplore.ieee.org/document/8530856/>.
- 2122 Jin Li, Andrew D. Heap, Anna Potter, and James J. Daniell. Application of machine learning methods to
 2123 spatial interpolation of environmental variables. *Environmental Modelling and Software*, 26(12):1647–
 2124 1659, dec 2011. ISSN 13648152. doi: 10.1016/j.envsoft.2011.07.004.
- 2125 Li-Ming Li, Wei-Guo Song, Jian Ma, and Kohyu Satoh. Artificial neural network approach for modeling
 2126 the impact of population density and weather parameters on forest fire risk. *International Journal
 2127 of Wildland Fire*, 18(6):640, oct 2009. ISSN 1049-8001. doi: 10.1071/WF07136. URL <http://www.publish.csiro.au/?paper=WF07136>.
 2128
- 2129 Shufeng Li, Alice C. Hughes, Tao Su, Julie Lebreton Anberrée, Alexei A. Oskolski, Mei Sun, David K.
 2130 Ferguson, and Zhekun Zhou. Fire dynamics under monsoonal climate in Yunnan, SW China: past,
 2131 present and future. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 465:168–176, jan 2017. ISSN
 2132 0031-0182. doi: 10.1016/J.PALAEO.2016.10.028. URL [https://www.sciencedirect.com/science/
 2133 article/pii/S0031018216306411](https://www.sciencedirect.com/science/article/pii/S0031018216306411).
- 2134 T. Li, E. Zhao, J. Zhang, and C. Hu. Detection of Wildfire Smoke Images Based on a Densely Dilated Con-
 2135 volutional Network. *Electronics*, 8(10):1131, oct 2019. ISSN 2079-9292. doi: 10.3390/electronics8101131.
 2136 URL <https://www.mdpi.com/2079-9292/8/10/1131>.
- 2137 Xiaolian Li, Weiguo Song, Liping Lian, and Xiaoge Wei. Forest Fire Smoke Detection Using Back-
 2138 Propagation Neural Network Based on MODIS Data. *Remote Sensing*, 7(4):4473–4498, apr 2015. ISSN
 2139 2072-4292. doi: 10.3390/rs70404473. URL <http://www.mdpi.com/2072-4292/7/4/4473>.
- 2140 Xiuqing Li, Zhenxue Chen, Q. M.J. Wu, and Chengyun Liu. 3D Parallel Fully Convolutional Networks for
 2141 Real-time Video Wildfire Smoke Detection, 2018b. ISSN 10518215.
- 2142 Hao Liang, Meng Zhang, and Hailan Wang. A Neural Network Model for Wildfire Scale Prediction Using
 2143 Meteorological Factors. *IEEE Access*, 7:176746–176755, 2019. ISSN 21693536. doi: 10.1109/ACCESS.
 2144 2019.2957837.
- 2145 Chul-Hee Lim, You Seung Kim, Myungsoo Won, Sea Jin Kim, and Woo-Kyun Lee. Can satellite-based data
 2146 substitute for surveyed data to predict the spatial probability of forest fire? A geostatistical approach to
 2147 forest fire in the Republic of Korea. *Geomatics, Natural Hazards and Risk*, 10(1):719–739, jan 2019. ISSN
 2148 1947-5705. doi: 10.1080/19475705.2018.1543210. URL [https://www.tandfonline.com/doi/full/10.
 2149 1080/19475705.2018.1543210](https://www.tandfonline.com/doi/full/10.1080/19475705.2018.1543210).
- 2150 Y Liu, Y Yang, C Liu, and Yu Gu. Forest Fire Detection Using Artificial Neural Network Algorithm Imple-
 2151 mented in Wireless Sensor Networks. *ZTE Communications*, jun 2015. URL [http://wwen.zte.com.
 2152 cn/endata/magazine/ztecommunications/2015/2/articles/201507/t20150724{}_443252.html](http://wwen.zte.com.cn/endata/magazine/ztecommunications/2015/2/articles/201507/t20150724{}_443252.html).

- 2153 Zelin Liu, Changhui Peng, Timothy Work, Jean-Noel Candau, Annie DesRochers, and Daniel Kneeshaw.
 2154 Application of machine-learning methods in forest ecology: recent progress and future challenges. *En-*
 2155 *vironmental Reviews*, 26(4):339–350, dec 2018. ISSN 1181-8700. doi: 10.1139/er-2018-0034. URL
 2156 <http://www.nrcresearchpress.com/doi/10.1139/er-2018-0034>.
- 2157 Zhihua Liu and Michael C. Wimberly. Climatic and Landscape Influences on Fire Regimes from 1984
 2158 to 2010 in the Western United States. *PLOS ONE*, 10(10):e0140839, oct 2015. ISSN 1932-6203. doi:
 2159 10.1371/journal.pone.0140839. URL <http://dx.plos.org/10.1371/journal.pone.0140839>.
- 2160 Zhihua Liu and Michael C. Wimberly. Direct and indirect effects of climate change on projected future
 2161 fire regimes in the western United States. *Science of The Total Environment*, 542:65–75, jan 2016.
 2162 ISSN 0048-9697. doi: 10.1016/J.SCITOTENV.2015.10.093. URL [https://www.sciencedirect.com/
 2163 science/article/pii/S0048969715309098](https://www.sciencedirect.com/science/article/pii/S0048969715309098).
- 2164 Zhihua Liu and Jian Yang. Quantifying ecological drivers of ecosystem productivity of the early-successional
 2165 boreal *Larix gmelinii* forest. *Ecosphere*, 5(7):art84, jul 2014. ISSN 2150-8925. doi: 10.1890/ES13-00372.1.
 2166 URL <http://doi.wiley.com/10.1890/ES13-00372.1>.
- 2167 Zhihua Liu, Jian Yang, and Hong S. He. Identifying the Threshold of Dominant Controls on Fire Spread
 2168 in a Boreal Forest Landscape of Northeast China. *PLoS ONE*, 8(1):e55618, jan 2013. ISSN 1932-6203.
 2169 doi: 10.1371/journal.pone.0055618. URL <https://dx.plos.org/10.1371/journal.pone.0055618>.
- 2170 Pablito M. López-Serrano, Carlos A. López-Sánchez, Juan G. Álvarez-González, and Jorge García-
 2171 Gutiérrez. A Comparison of Machine Learning Techniques Applied to Landsat-5 TM Spectral Data
 2172 for Biomass Estimation. *Canadian Journal of Remote Sensing*, 42(6):690–705, nov 2016. ISSN 0703-
 2173 8992. doi: 10.1080/07038992.2016.1217485. URL [https://www.tandfonline.com/doi/full/10.1080/
 2174 07038992.2016.1217485](https://www.tandfonline.com/doi/full/10.1080/07038992.2016.1217485).
- 2175 F. Javier Lozano, S. Suárez-Seoane, M. Kelly, and E. Luis. A multi-scale approach for modeling fire
 2176 occurrence probability using satellite data and classification trees: A case study in a mountain-
 2177 ous Mediterranean region. *Remote Sensing of Environment*, 112(3):708–719, mar 2008. ISSN 0034-
 2178 4257. doi: 10.1016/J.RSE.2007.06.006. URL [https://www.sciencedirect.com/science/article/
 2179 pii/S003442570700243X](https://www.sciencedirect.com/science/article/pii/S003442570700243X).
- 2180 V Lozhkin, D Tarkhov, V Timofeev, O Lozhkina, and A Vasilyev. Differential neural network ap-
 2181 proach in information process for prediction of roadside air pollution by peat fire. *IOP Conference*
 2182 *Series: Materials Science and Engineering*, 158(1):012063, nov 2016. ISSN 1757-8981. doi: 10.
 2183 1088/1757-899X/158/1/012063. URL [http://stacks.iop.org/1757-899X/158/i=1/a=012063?key=
 2184 crossref.7abf8c3fd66f7ce48986b4554f7aecd5](http://stacks.iop.org/1757-899X/158/i=1/a=012063?key=crossref.7abf8c3fd66f7ce48986b4554f7aecd5).
- 2185 Guilan Luo, Mei Zhang, Zizhong Yang, and Mingmei Song. Data mining of correlation between fire
 2186 disturbance habitat factors and spider communities. In *2017 4th International Conference on Systems*
 2187 *and Informatics (ICSAI)*, pages 1471–1476. IEEE, nov 2017. ISBN 978-1-5386-1107-4. doi: 10.1109/
 2188 ICSAI.2017.8248518. URL <http://ieeexplore.ieee.org/document/8248518/>.
- 2189 Ruisen Luo, Yingying Dong, Muye Gan, Dejun Li, Shuli Niu, Amy Oliver, Ke Wang, and Yiqi Luo. Global
 2190 Analysis of Influencing Forces of Fire Activity: the Threshold Relationships between Vegetation and
 2191 Fire. *Life Science Journal*, 10(2):15–24, 2013. ISSN 0300-9165.
- 2192 Duncan C Lutes, Robert E Keane, John F Caratti, Carl H Key, Nathan C Benson, Steve Sutherland,
 2193 and Larry J Gangi. FIREMON: Fire effects monitoring and inventory system. *Gen. Tech. Rep. RMRS-*
 2194 *GTR-164*. Fort Collins, CO: US Department of Agriculture, Forest Service, Rocky Mountain Research
 2195 Station. 1 CD., 164, 2006.

- 2196 Jamie M. Lydersen, Malcolm P. North, and Brandon M. Collins. Severity of an uncharacteristically large
 2197 wildfire, the Rim Fire, in forests with relatively restored frequent fire regimes. *Forest Ecology and*
 2198 *Management*, 328:326–334, sep 2014. ISSN 0378-1127. doi: 10.1016/J.FORECO.2014.06.005. URL
 2199 <https://www.sciencedirect.com/science/article/pii/S0378112714003661>.
- 2200 Jamie M. Lydersen, Brandon M. Collins, Matthew L. Brooks, John R. Matchett, Kristen L. Shive,
 2201 Nicholas A. Povak, Van R. Kane, and Douglas F. Smith. Evidence of fuels management and fire weather
 2202 influencing fire severity in an extreme fire event. *Ecological Applications*, 27(7):2013–2030, oct 2017.
 2203 ISSN 10510761. doi: 10.1002/eap.1586. URL <http://doi.wiley.com/10.1002/eap.1586>.
- 2204 James MacQueen et al. Some methods for classification and analysis of multivariate observations. In
 2205 *Proceedings of the fifth Berkeley symposium on mathematical statistics and probability*, volume 1, pages
 2206 281–297. Oakland, CA, USA, 1967.
- 2207 Eduardo Eiji Maeda, Antonio Roberto Formaggio, Yosio Edemir Shimabukuro, Gustavo Felipe Balu e
 2208 Arcoverde, and Matthew C. Hansen. Predicting forest fire in the Brazilian Amazon using MODIS
 2209 imagery and artificial neural networks. *International Journal of Applied Earth Observation and*
 2210 *Geoinformation*, 11(4):265–272, aug 2009. ISSN 0303-2434. doi: 10.1016/J.JAG.2009.03.003. URL
 2211 <https://www.sciencedirect.com/science/article/pii/S0303243409000233>.
- 2212 Nyasha Magadzire, Helen M. Klerk, Karen J. Esler, and Jasper A. Slingsby. Fire and life history affect the
 2213 distribution of plant species in a biodiversity hotspot. *Diversity and Distributions*, 25(7):1012–1023, jul
 2214 2019. ISSN 1366-9516. doi: 10.1111/ddi.12921. URL [https://onlinelibrary.wiley.com/doi/abs/](https://onlinelibrary.wiley.com/doi/abs/10.1111/ddi.12921)
 2215 [10.1111/ddi.12921](https://onlinelibrary.wiley.com/doi/abs/10.1111/ddi.12921).
- 2216 G. Mallinis, F. Maris, I. Kalinderis, and N. Koutsias. Assessment of Post-fire Soil Erosion Risk in Fire-
 2217 Affected Watersheds Using Remote Sensing and GIS. *GIScience & Remote Sensing*, 46(4):388–410, oct
 2218 2009. ISSN 1548-1603. doi: 10.2747/1548-1603.46.4.388. URL [https://www.tandfonline.com/doi/](https://www.tandfonline.com/doi/full/10.2747/1548-1603.46.4.388)
 2219 [full/10.2747/1548-1603.46.4.388](https://www.tandfonline.com/doi/full/10.2747/1548-1603.46.4.388).
- 2220 Nicolas Mansuy, Carol Miller, Marc-Andr e Parisien, Sean A Parks, Enric Batllori, and Max A Moritz. Con-
 2221 trasting human influences and macro-environmental factors on fire activity inside and outside protected
 2222 areas of North America. *Environmental Research Letters*, 14(6):064007, may 2019. ISSN 1748-9326.
 2223 doi: 10.1088/1748-9326/ab1bc5. URL [https://iopscience.iop.org/article/10.1088/1748-9326/](https://iopscience.iop.org/article/10.1088/1748-9326/ab1bc5)
 2224 [ab1bc5](https://iopscience.iop.org/article/10.1088/1748-9326/ab1bc5).
- 2225 Natasha Markuzon and Stephan Kolitz. Data driven approach to estimating fire danger from satellite
 2226 images and weather information. In *2009 IEEE Applied Imagery Pattern Recognition Workshop (AIPR*
 2227 *2009)*, pages 1–7. IEEE, oct 2009. ISBN 978-1-4244-5146-3. doi: 10.1109/AIPR.2009.5466309. URL
 2228 <http://ieeexplore.ieee.org/document/5466309/>.
- 2229 David L. Martell. A review of recent forest and wildland fire management decision support systems research,
 2230 jun 2015. ISSN 21986436.
- 2231 Yago Mart n, Mar a Z niga-Ant n, and Marcos Rodrigues Mimbbrero. Modelling temporal variation of
 2232 fire-occurrence towards the dynamic prediction of human wildfire ignition danger in northeast Spain.
 2233 *Geomatics, Natural Hazards and Risk*, 10(1):385–411, jan 2019. ISSN 1947-5705. doi: 10.1080/19475705.
 2234 2018.1526219. URL <https://www.tandfonline.com/doi/full/10.1080/19475705.2018.1526219>.
- 2235 S. Mart n-Alc n and L Coll. Unraveling the relative importance of factors driving post-fire regenera-
 2236 tion trajectories in non-serotinous Pinus nigra forests. *Forest Ecology and Management*, 361:13–22, feb
 2237 2016. ISSN 0378-1127. doi: 10.1016/J.FORECO.2015.11.006. URL [https://www-sciencedirect-com.](https://www-sciencedirect-com.login.ezproxy.library.ualberta.ca/science/article/pii/S037811271500612X)
 2238 [login.ezproxy.library.ualberta.ca/science/article/pii/S037811271500612X](https://www-sciencedirect-com.login.ezproxy.library.ualberta.ca/science/article/pii/S037811271500612X).

- 2239 Arif Masrur, Andrey N. Petrov, and John DeGroot. Circumpolar spatio-temporal patterns and contribut-
 2240 ing climatic factors of wildfire activity in the Arctic tundra from 2001-2015. *Environmental Research*
 2241 *Letters*, 13(1), jan 2018. ISSN 17489326. doi: 10.1088/1748-9326/aa9a76.
- 2242 Robert Mavsar, Armando González Cabán, and Elsa Varela. The state of development of fire management
 2243 decision support systems in America and Europe. *Forest Policy and Economics*, 29:45–55, apr 2013.
 2244 ISSN 13899341. doi: 10.1016/j.forpol.2012.11.009.
- 2245 R. Stockton Maxwell, Alan H. Taylor, Carl N. Skinner, Hugh D. Safford, Rachel E. Isaacs, Catherine Airey,
 2246 and Amanda B. Young. Landscape-scale modeling of reference period forest conditions and fire behavior
 2247 on heavily logged lands. *Ecosphere*, 5(3):art32, mar 2014. ISSN 2150-8925. doi: 10.1890/ES13-00294.1.
 2248 URL <http://doi.wiley.com/10.1890/ES13-00294.1>.
- 2249 M.J. Mayr, K.A. Vanselow, and C. Samimi. Fire regimes at the arid fringe: A 16-year remote sensing
 2250 perspective (2000–2016) on the controls of fire activity in Namibia from spatial predictive models. *Eco-*
 2251 *logical Indicators*, 91:324–337, aug 2018. ISSN 1470-160X. doi: 10.1016/J.ECOLIND.2018.04.022. URL
 2252 <https://www.sciencedirect.com/science/article/pii/S1470160X18302759>.
- 2253 Ronald J McCormick, Thomas A Brandner, and Timothy F H Allen. TOWARD A THEORY OF MESO-
 2254 SCALE WILDFIRE MODELING-A COMPLEX SYSTEMS APPROACH USING ARTIFICIAL NEU-
 2255 RAL NETWORKS. In *The Joint Fire Science Conference And Workshop*, Boise, Idaho, 1999.
- 2256 Amy McGovern, Kimberly L. Elmore, David John Gagne, Sue Ellen Haupt, Christopher D. Karstens, Ryan
 2257 Lagerquist, Travis Smith, John K. Williams, Amy McGovern, Kimberly L. Elmore, David John Gagne
 2258 II, Sue Ellen Haupt, Christopher D. Karstens, Ryan Lagerquist, Travis Smith, and John K. Williams.
 2259 Using Artificial Intelligence to Improve Real-Time Decision-Making for High-Impact Weather. *Bulletin*
 2260 *of the American Meteorological Society*, 98(10):2073–2090, oct 2017. ISSN 0003-0007. doi: 10.1175/
 2261 BAMS-D-16-0123.1. URL <http://journals.ametsoc.org/doi/10.1175/BAMS-D-16-0123.1>.
- 2262 Amy McGovern, Ryan Lagerquist, David John Gagne, G. Eli Jergensen, Kimberly L. Elmore, Cameron R.
 2263 Homeyer, and Travis Smith. Making the Black Box More Transparent: Understanding the Physical
 2264 Implications of Machine Learning. *Bulletin of the American Meteorological Society*, 100(11):2175–2199,
 2265 nov 2019. ISSN 0003-0007. doi: 10.1175/BAMS-D-18-0195.1. URL <http://journals.ametsoc.org/doi/10.1175/BAMS-D-18-0195.1>.
- 2267 Sean McGregor, Rachel Houtman, Hailey Buckingham, Claire Montgomery, Ronald Metoyer, and
 2268 Thomas G Dietterich. Fast simulation for computational sustainability sequential decision making prob-
 2269 lems. In *Proceedings of the 4th International Conference on Computational Sustainability*, pages 5–7,
 2270 2016.
- 2271 Sean McGregor, Rachel Houtman, Claire Montgomery, Ronald Metoyer, and Thomas G. Dietterich. Fast
 2272 Optimization of Wildfire Suppression Policies with SMAC. *arXiv preprint*, mar 2017. URL <http://arxiv.org/abs/1703.09391>.
- 2274 James P. Minas, John W. Hearne, and John W. Handmer. A review of operations research methods
 2275 applicable to wildfire management. *International Journal of Wildland Fire*, 21(3):189, may 2012. ISSN
 2276 1049-8001. doi: 10.1071/WF10129. URL <http://www.publish.csiro.au/?paper=WF10129>.
- 2277 Yosune Miquelajauregui, Steven G. Cumming, and Sylvie Gauthier. Modelling Variable Fire Severity in
 2278 Boreal Forests: Effects of Fire Intensity and Stand Structure. *PLOS ONE*, 11(2):e0150073, feb 2016.
 2279 ISSN 1932-6203. doi: 10.1371/journal.pone.0150073. URL <https://dx.plos.org/10.1371/journal.pone.0150073>.
- 2281 Melanie Mitchell. *An introduction to genetic algorithms*. MIT Press, 1996. ISBN 9780262133166.

- 2282 T.M. Mitchell. *Machine Learning*. McGraw-Hill, 1997. ISBN 0071154671.
- 2283 Varun Mithal, Guruprasad Nayak, Ankush Khandelwal, Vipin Kumar, Ramakrishna Nemani, and Nikunj
2284 Oza. Mapping burned areas in tropical forests using a novel machine learning framework. *Remote*
2285 *Sensing*, 10(1):69, 2018.
- 2286 Nikolaos E. Mitrakis, Giorgos Mallinis, Nikos Koutsias, and John B. Theocharis. Burned area mapping
2287 in Mediterranean environment using medium-resolution multi-spectral data and a neuro-fuzzy classifier.
2288 *International Journal of Image and Data Fusion*, 3(4):299–318, dec 2012. ISSN 1947-9832. doi: 10.
2289 1080/19479832.2011.635604. URL [http://www.tandfonline.com/doi/abs/10.1080/19479832.2011.](http://www.tandfonline.com/doi/abs/10.1080/19479832.2011.635604)
2290 [635604](http://www.tandfonline.com/doi/abs/10.1080/19479832.2011.635604).
- 2291 Ioannis Mitsopoulos and Giorgos Mallinis. A data-driven approach to assess large fire size generation in
2292 Greece. *Natural Hazards*, 88(3):1591–1607, sep 2017. ISSN 0921-030X. doi: 10.1007/s11069-017-2934-z.
2293 URL <http://link.springer.com/10.1007/s11069-017-2934-z>.
- 2294 J. R. Molina, A. Lora, C. Prades, and F. Rodríguez y Silva. Roadside vegetation planning and con-
2295 servation: New approach to prevent and mitigate wildfires based on fire ignition potential. *Forest*
2296 *Ecology and Management*, 444:163–173, jul 2019. ISSN 0378-1127. doi: 10.1016/J.FORECO.2019.
2297 04.034. URL [https://www-sciencedirect-com.login.ezproxy.library.ualberta.ca/science/](https://www-sciencedirect-com.login.ezproxy.library.ualberta.ca/science/article/pii/S0378112719301501)
2298 [article/pii/S0378112719301501](https://www-sciencedirect-com.login.ezproxy.library.ualberta.ca/science/article/pii/S0378112719301501).
- 2299 Max A. Moritz, Marc-André Parisien, Enric Batllori, Meg A. Krawchuk, Jeff Van Dorn, David J. Ganz, and
2300 Katharine Hayhoe. Climate change and disruptions to global fire activity. *Ecosphere*, 3(6):art49, jun 2012.
2301 ISSN 2150-8925. doi: 10.1890/ES11-00345.1. URL <http://doi.wiley.com/10.1890/ES11-00345.1>.
- 2302 Amir Mosavi, Pinar Ozturk, Kwok-wing Chau, Amir Mosavi, Pinar Ozturk, and Kwok-wing Chau. Flood
2303 Prediction Using Machine Learning Models: Literature Review. *Water*, 10(11):1536, oct 2018. ISSN
2304 2073-4441. doi: 10.3390/w10111536. URL <http://www.mdpi.com/2073-4441/10/11/1536>.
- 2305 Mohsen MOSTAFA, Shaban SHATAEE JOUIBARY, Majid LOTFALIAN, and Amir SADODDIN. WA-
2306 TERSHED ROAD NETWORK ANALYSIS WITH AN EMPHASIS ON FIRE FIGHTING MANAGE-
2307 MENT. *Journal of Environmental Engineering and Landscape Management*, 25(4):342–353, dec 2017.
2308 ISSN 1648-6897. doi: 10.3846/16486897.2017.1281816. URL [http://journals.vgtu.lt/index.php/](http://journals.vgtu.lt/index.php/JEELM/article/view/1712)
2309 [JEELM/article/view/1712](http://journals.vgtu.lt/index.php/JEELM/article/view/1712).
- 2310 Kudzai S. Mpakairi, Paradzayi Tagwireyi, Henry Ndaimani, and Hilary T. Madiri. Distribution of wild-
2311 land fires and possible hotspots for the Zimbabwean component of Kavango-Zambezi Transfrontier Con-
2312 servation Area. *South African Geographical Journal*, 101(1):110–120, jan 2019. ISSN 0373-6245. doi:
2313 10.1080/03736245.2018.1541023. URL [https://www.tandfonline.com/doi/full/10.1080/03736245.](https://www.tandfonline.com/doi/full/10.1080/03736245.2018.1541023)
2314 [2018.1541023](https://www.tandfonline.com/doi/full/10.1080/03736245.2018.1541023).
- 2315 Khan Muhammad, Jamil Ahmad, and Sung Wook Baik. Early fire detection using convolutional neural
2316 networks during surveillance for effective disaster management. *Neurocomputing*, 288:30–42, may 2018.
2317 ISSN 18728286. doi: 10.1016/j.neucom.2017.04.083.
- 2318 Kevin Murphy. *Machine Learning: A Probabilistic Perspective*. MIT Press, 2012. URL [http://www.](http://www.amazon.com/Machine-Learning-Probabilistic-Perspective-Computation/dp/0262018020)
2319 [amazon.com/Machine-Learning-Probabilistic-Perspective-Computation/dp/0262018020](http://www.amazon.com/Machine-Learning-Probabilistic-Perspective-Computation/dp/0262018020).
- 2320 Khurram Nadeem, S. W. Taylor, Douglas G. Woolford, and C. B. Dean. Mesoscale spatiotemporal pre-
2321 dictive models of daily human- and lightning-caused wildland fire occurrence in British Columbia. *In-*
2322 *ternational Journal of Wildland Fire*, 29(1):11, 2020. ISSN 1049-8001. doi: 10.1071/WF19058. URL
2323 <http://www.publish.csiro.au/?paper=WF19058>.

- 2324 Hariharan Naganathan, Sudarshan P Seshasayee, Jonghoon Kim, Wai K Chong, and Jui-Sheng Chou.
 2325 Wildfire Predictions: Determining Reliable Models using Fused Dataset. *Global Journal of Computer*
 2326 *Science and Technology*, 16(4C), 2016. URL [https://computerresearch.org/index.php/computer/](https://computerresearch.org/index.php/computer/article/view/1437)
 2327 [article/view/1437](https://computerresearch.org/index.php/computer/article/view/1437).
- 2328 Nicholas J. Nauslar, Benjamin J. Hatchett, Timothy J. Brown, Michael L. Kaplan, and John F. Mejia.
 2329 Impact of the North American monsoon on wildfire activity in the southwest United States. *International*
 2330 *Journal of Climatology*, 39(3):1539–1554, mar 2019. ISSN 08998418. doi: 10.1002/joc.5899. URL
 2331 <http://doi.wiley.com/10.1002/joc.5899>.
- 2332 Trisalyn A. Nelson, Wiebe Nijland, Mathieu L. Bourbonnais, and Michael A. Wulder. Regression Tree
 2333 Modeling of Spatial Pattern and Process Interactions. In *Mapping Forest Landscape Patterns*, pages
 2334 187–212. Springer New York, New York, NY, 2017. doi: 10.1007/978-1-4939-7331-6_5. URL [http://](http://link.springer.com/10.1007/978-1-4939-7331-6_5)
 2335 link.springer.com/10.1007/978-1-4939-7331-6_5.
- 2336 Nguyen Ngoc Thach, Dang Bao-Toan Ngo, Pham Xuan-Canh, Nguyen Hong-Thi, Bui Hang Thi, Hoang
 2337 Nhat-Duc, and Tien Bui Dieu. Spatial pattern assessment of tropical forest fire danger at Thuan Chau
 2338 area (Vietnam) using GIS-based advanced machine learning algorithms: A comparative study. *Ecological*
 2339 *Informatics*, 46:74–85, jul 2018. ISSN 1574-9541. doi: 10.1016/J.ECOINF.2018.05.009. URL [https://](https://www.sciencedirect.com/science/article/pii/S1574954118300852)
 2340 www.sciencedirect.com/science/article/pii/S1574954118300852.
- 2341 I. Nitze, G. Grosse, B. M. Jones, V. E. Romanovsky, and J. Boike. Remote sensing quantifies widespread
 2342 abundance of permafrost region disturbances across the Arctic and Subarctic. *Nature Communications*,
 2343 9(1), dec 2018. ISSN 20411723. doi: 10.1038/s41467-018-07663-3.
- 2344 Christopher D. O’ O’Connor, David E. Calkin, Matthew P. Thompson, Christopher D.O. O’Connor,
 2345 David E. Calkin, and Matthew P. Thompson. An empirical machine learning method for predict-
 2346 ing potential fire control locations for pre-fire planning and operational fire management. *Interna-*
 2347 *tional Journal of Wildland Fire*, 26(7):587, jul 2017. ISSN 1049-8001. doi: 10.1071/WF16135. URL
 2348 <http://www.publish.csiro.au/?paper=WF16135>.
- 2349 Julian D. Olden, Joshua J. Lawler, and N. LeRoy Poff. Machine Learning Methods Without Tears: A
 2350 Primer for Ecologists. *The Quarterly Review of Biology*, 83(2):171–193, jun 2008. ISSN 0033-5770. doi:
 2351 10.1086/587826. URL <https://www.journals.uchicago.edu/doi/10.1086/587826>.
- 2352 Sandra Oliveira, Friderike Oehler, Jesús San-Miguel-Ayanz, Andrea Camia, and José M.C. Pereira.
 2353 Modeling spatial patterns of fire occurrence in Mediterranean Europe using Multiple Regression
 2354 and Random Forest. *Forest Ecology and Management*, 275:117–129, jul 2012. ISSN 0378-1127.
 2355 doi: 10.1016/J.FORECO.2012.03.003. URL [https://www.sciencedirect.com/science/article/](https://www.sciencedirect.com/science/article/pii/S0378112712001272)
 2356 [pii/S0378112712001272](https://www.sciencedirect.com/science/article/pii/S0378112712001272).
- 2357 A. Murat Özbayoğlu and Recep Bozer. Estimation of the Burned Area in Forest Fires Using Compu-
 2358 tational Intelligence Techniques. *Procedia Computer Science*, 12:282–287, jan 2012. ISSN 1877-0509.
 2359 doi: 10.1016/J.PROCS.2012.09.070. URL [https://www.sciencedirect.com/science/article/pii/](https://www.sciencedirect.com/science/article/pii/S1877050912006618)
 2360 [S1877050912006618](https://www.sciencedirect.com/science/article/pii/S1877050912006618).
- 2361 P. Papakosta, G. Xanthopoulos, and D. Straub. Probabilistic prediction of wildfire economic losses to
 2362 housing in Cyprus using Bayesian network analysis. *International Journal of Wildland Fire*, 26(1):10, feb
 2363 2017. ISSN 1049-8001. doi: 10.1071/WF15113. URL <http://www.publish.csiro.au/?paper=WF15113>.
- 2364 Marc-André Parisien and Max A. Moritz. Environmental controls on the distribution of wildfire at multiple
 2365 spatial scales. *Ecological Monographs*, 79(1):127–154, feb 2009. ISSN 0012-9615. doi: 10.1890/07-1289.1.
 2366 URL <http://doi.wiley.com/10.1890/07-1289.1>.

- 2367 Marc-André Parisien, Sean A. Parks, Meg A. Krawchuk, John M. Little, Mike D. Flannigan, Lynn M.
 2368 Gowman, and Max A. Moritz. An analysis of controls on fire activity in boreal Canada: comparing
 2369 models built with different temporal resolutions. *Ecological Applications*, 24(6):1341–1356, sep 2014.
 2370 ISSN 1051-0761. doi: 10.1890/13-1477.1. URL <http://doi.wiley.com/10.1890/13-1477.1>.
- 2371 Marc And  Parisien, Susan Snetsinger, Jonathan A. Greenberg, Cara R. Nelson, Tania Schoennagel,
 2372 Solomon Z. Dobrowski, and Max A. Moritz. Spatial variability in wildfire probability across the west-
 2373 ern United States. *International Journal of Wildland Fire*, 21(4):313–327, 2012. ISSN 10498001. doi:
 2374 10.1071/WF11044.
- 2375 Sean A. Parks, Carol Miller, Marc-Andr  Parisien, Lisa M. Holsinger, Solomon Z. Dobrowski, and John
 2376 Abatzoglou. Wildland fire deficit and surplus in the western United States, 1984–2012. *Ecosphere*, 6
 2377 (12):art275, dec 2015. ISSN 2150-8925. doi: 10.1890/ES15-00294.1. URL <http://doi.wiley.com/10.1890/ES15-00294.1>.
- 2379 Sean A Parks, Carol Miller, John T Abatzoglou, Lisa M Holsinger, Marc-Andr  Parisien, and
 2380 Solomon Z Dobrowski. How will climate change affect wildland fire severity in the west-
 2381 ern US? *Environmental Research Letters*, 11(3):035002, mar 2016. ISSN 1748-9326. doi:
 2382 10.1088/1748-9326/11/3/035002. URL <http://stacks.iop.org/1748-9326/11/i=3/a=035002?key=crossref.4d33abcb068f5458baf3b94828ca073e>.
- 2384 Sean A Parks, Lisa M Holsinger, Matthew H Panunto, W Matt Jolly, Solomon Z Dobrowski, and
 2385 Gregory K Dillon. High-severity fire: evaluating its key drivers and mapping its probability across
 2386 western US forests. *Environmental Research Letters*, 13(4):044037, apr 2018. ISSN 1748-9326.
 2387 doi: 10.1088/1748-9326/aab791. URL <http://stacks.iop.org/1748-9326/13/i=4/a=044037?key=crossref.5c2b6b1d5870d4a9269af3badf873e81>.
- 2389 Judea Pearl. *Probabilistic Reasoning in Intelligent Systems: Networks of Plausible Inference*. Morgan Kauf-
 2390 mann, San Mateo, California, 1988.
- 2391 T. D. Penman, O. Price, and R. A. Bradstock. Bayes Nets as a method for analysing the influence of
 2392 management actions in fire planning. *International Journal of Wildland Fire*, 20(8):909–920, 2011.
 2393 ISSN 10498001. doi: 10.1071/WF10076.
- 2394 T. D. Penman, R. A. Bradstock, and O. F. Price. Reducing wildfire risk to urban developments:
 2395 Simulation of cost-effective fuel treatment solutions in south eastern Australia. *Environmental Mod-
 2396 elling and Software*, 52:166–175, 2014. ISSN 13648152. doi: 10.1016/j.envsoft.2013.09.030. URL
 2397 <http://dx.doi.org/10.1016/j.envsoft.2013.09.030>.
- 2398 T. D. Penman, A. E. Nicholson, R. A. Bradstock, L. Collins, S. H. Penman, and O. F. Price. Reducing
 2399 the risk of house loss due to wildfires. *Environmental Modelling and Software*, 67:12–25, 2015. ISSN
 2400 13648152. doi: 10.1016/j.envsoft.2014.12.020.
- 2401 Allan Pereira, Jos  Pereira, Renata Libonati, Duarte Oom, Alberto Setzer, Fabiano Morelli, Fausto
 2402 Machado-Silva, Luis de Carvalho, Allan A. Pereira, Jos  M. C. Pereira, Renata Libonati, Duarte Oom,
 2403 Alberto W. Setzer, Fabiano Morelli, Fausto Machado-Silva, and Luis Marcelo Tavares de Carvalho.
 2404 Burned Area Mapping in the Brazilian Savanna Using a One-Class Support Vector Machine Trained by
 2405 Active Fires. *Remote Sensing*, 9(11):1161, nov 2017. ISSN 2072-4292. doi: 10.3390/rs9111161. URL
 2406 <http://www.mdpi.com/2072-4292/9/11/1161>.
- 2407 George L. W. Perry, Janet M. Wilmshurst, Matt S. McGlone, and Aaron Napier. Reconstructing spatial
 2408 vulnerability to forest loss by fire in pre-historic New Zealand. *Global Ecology and Biogeography*, 21
 2409 (10):1029–1041, oct 2012. ISSN 1466822X. doi: 10.1111/j.1466-8238.2011.00745.x. URL <http://doi.wiley.com/10.1111/j.1466-8238.2011.00745.x>.

- 2411 Matthew P. Peters and Louis R. Iverson. Incorporating fine-scale drought information into an eastern US
 2412 wildfire hazard model. *International Journal of Wildland Fire*, 26(5):393, may 2017. ISSN 1049-8001.
 2413 doi: 10.1071/WF16130. URL <http://www.publish.csiro.au/?paper=WF16130>.
- 2414 Matthew P. Peters, Louis R. Iverson, Stephen N. Matthews, and Anantha M. Prasad. Wildfire hazard
 2415 mapping: exploring site conditions in eastern US wildland–urban interfaces. *International Journal of*
 2416 *Wildland Fire*, 22(5):567, aug 2013. ISSN 1049-8001. doi: 10.1071/WF12177. URL [http://www.](http://www.publish.csiro.au/?paper=WF12177)
 2417 [publish.csiro.au/?paper=WF12177](http://www.publish.csiro.au/?paper=WF12177).
- 2418 G. P. Petropoulos, W. Knorr, M. Scholze, L. Boschetti, and G. Karantounias. Combining ASTER
 2419 multispectral imagery analysis and support vector machines for rapid and cost-effective post-fire as-
 2420 sessment: a case study from the Greek wildland fires of 2007. *Natural Hazards and Earth Sys-*
 2421 *tem Science*, 10(2):305–317, feb 2010. ISSN 1684-9981. doi: 10.5194/nhess-10-305-2010. URL
 2422 <http://www.nat-hazards-earth-syst-sci.net/10/305/2010/>.
- 2423 George P. Petropoulos, Charalambos Kontoes, and Iphigenia Keramitsoglou. Burnt area delineation from
 2424 a uni-temporal perspective based on Landsat TM imagery classification using Support Vector Ma-
 2425 chines. *International Journal of Applied Earth Observation and Geoinformation*, 13(1):70–80, feb 2011.
 2426 ISSN 0303-2434. doi: 10.1016/J.JAG.2010.06.008. URL [https://www.sciencedirect.com/science/](https://www.sciencedirect.com/science/article/pii/S0303243410000784)
 2427 [article/pii/S0303243410000784](https://www.sciencedirect.com/science/article/pii/S0303243410000784).
- 2428 M.T. Pham, A. Rajić, J.D. Greig, J.M. Sargeant, A. Papadopoulos, and S.A. Mcewen. A scoping review of
 2429 scoping reviews: Advancing the approach and enhancing the consistency. *Research Synthesis Methods*,
 2430 5(4):371–385, dec 2014. ISSN 17592887. doi: 10.1002/jrsm.1123.
- 2431 Thanh Cong Phan and Thanh Tam Nguyen. Remote Sensing meets Deep Learning: Exploiting Spatio-
 2432 Temporal-Spectral Satellite Images for Early Wildfire Detection. Technical report, 2019. URL [https:](https://infoscience.epfl.ch/record/270339)
 2433 [://infoscience.epfl.ch/record/270339](https://infoscience.epfl.ch/record/270339).
- 2434 Sharon B. Phillips, Viney P. Aneja, Daiwen Kang, and S. Pal Arya. Modelling and analysis of the atmo-
 2435 spheric nitrogen deposition in North Carolina. In *International Journal of Global Environmental Issues*,
 2436 volume 6, pages 231–252, 2006. doi: 10.1016/j.ecolmodel.2005.03.026.
- 2437 Andrew D. Pierce, Calvin A. Farris, and Alan H. Taylor. Use of random forests for modeling and mapping
 2438 forest canopy fuels for fire behavior analysis in Lassen Volcanic National Park, California, USA. *Forest*
 2439 *Ecology and Management*, 279:77–89, sep 2012. ISSN 0378-1127. doi: 10.1016/J.FORECO.2012.05.010.
 2440 URL <https://www.sciencedirect.com/science/article/pii/S0378112712002654>.
- 2441 David L Poole and Alan K Mackworth. *Artificial Intelligence: foundations of computational agents*. Cam-
 2442 bridge University Press, 2010.
- 2443 Patrick Poon, Alicia Kinoshita, Patrick K. Poon, and Alicia M. Kinoshita. Estimating Evapotranspiration
 2444 in a Post-Fire Environment Using Remote Sensing and Machine Learning. *Remote Sensing*, 10(11):1728,
 2445 nov 2018. ISSN 2072-4292. doi: 10.3390/rs10111728. URL [http://www.mdpi.com/2072-4292/10/11/](http://www.mdpi.com/2072-4292/10/11/1728)
 2446 [1728](http://www.mdpi.com/2072-4292/10/11/1728).
- 2447 Zohre Sadat Pourtaghi, Hamid Reza Pourghasemi, Roberta Aretano, and Teodoro Semeraro. Investigation
 2448 of general indicators influencing on forest fire and its susceptibility modeling using different data mining
 2449 techniques. *Ecological Indicators*, 64:72–84, may 2016. ISSN 1470-160X. doi: 10.1016/J.ECOLIND.
 2450 2015.12.030. URL <https://www.sciencedirect.com/science/article/pii/S1470160X15007359>.
- 2451 Ruiliang Pu and Peng Gong. Determination of Burnt Scars Using Logistic Regression and
 2452 Neural Network Techniques from a Single Post-Fire Landsat 7 ETM + Image. *Photogram-*
 2453 *metric Engineering & Remote Sensing*, 70(7):841–850, jul 2004. ISSN 00991112. doi: 10.

- 14358/PERS.70.7.841. URL <http://openurl.ingenta.com/content/xref?genre=article{&}issn=0099-1112{&}volume=70{&}issue=7{&}spage=841>.
- J Ross Quinlan. C 4.5: Programs for machine learning. *The Morgan Kaufmann Series in Machine Learning, San Mateo, CA: Morgan Kaufmann,— c1993*, 1993.
- Carmen Quintano, Alfonso Fernández-Manso, Leonor Calvo, and Dar A. Roberts. Vegetation and Soil Fire Damage Analysis Based on Species Distribution Modeling Trained with Multispectral Satellite Data. *Remote Sensing*, 11(15):1832, aug 2019. ISSN 2072-4292. doi: 10.3390/rs11151832. URL <https://www.mdpi.com/2072-4292/11/15/1832>.
- Natalia Quintero, Olga Viedma, Itziar R. Urbieto, and José M. Moreno. Assessing Landscape Fire Hazard by Multitemporal Automatic Classification of Landsat Time Series Using the Google Earth Engine in West-Central Spain. *Forests*, 10(6):518, jun 2019. ISSN 1999-4907. doi: 10.3390/f10060518. URL <https://www.mdpi.com/1999-4907/10/6/518>.
- David Radke, Anna Hessler, and Dan Ellsworth. FireCast: Leveraging Deep Learning to Predict Wildfire Spread. In *Proceedings of the Twenty-Eighth International Joint Conference on Artificial Intelligence (IJCAI-19)*, pages 4575–4581. International Joint Conferences on Artificial Intelligence, jul 2019. doi: 10.24963/ijcai.2019/636.
- M. Raissi, P. Perdikaris, and G. E. Karniadakis. Physics-informed neural networks: A deep learning framework for solving forward and inverse problems involving nonlinear partial differential equations. *Journal of Computational Physics*, 378:686–707, feb 2019. ISSN 10902716. doi: 10.1016/j.jcp.2018.10.045.
- Maziar Raissi and George Em Karniadakis. Hidden physics models: Machine learning of nonlinear partial differential equations. *Journal of Computational Physics*, 357:125–141, mar 2018. ISSN 10902716. doi: 10.1016/j.jcp.2017.11.039.
- Carl Edward Rasmussen and Christopher KI Williams. *Gaussian processes for machine learning*, volume 1. MIT press Cambridge, 2006.
- Stephan Rasp and Sebastian Lerch. Neural Networks for Postprocessing Ensemble Weather Forecasts. *Monthly Weather Review*, 146(11):3885–3900, nov 2018. ISSN 0027-0644. doi: 10.1175/MWR-D-18-0187.1. URL <http://journals.ametsoc.org/doi/10.1175/MWR-D-18-0187.1>.
- Markus Reichstein, Gustau Camps-Valls, Bjorn Stevens, Martin Jung, Joachim Denzler, Nuno Carvalhais, and Prabhat. Deep learning and process understanding for data-driven Earth system science. *Nature*, 566(7743):195–204, feb 2019. ISSN 0028-0836. doi: 10.1038/s41586-019-0912-1. URL <http://www.nature.com/articles/s41586-019-0912-1>.
- Colleen E. Reid, Michael Jerrett, Maya L. Petersen, Gabriele G. Pfister, Philip E. Morefield, Ira B. Tager, Sean M. Raffuse, and John R. Balmes. Spatiotemporal Prediction of Fine Particulate Matter During the 2008 Northern California Wildfires Using Machine Learning. *Environmental Science & Technology*, 49(6):3887–3896, mar 2015. ISSN 0013-936X. doi: 10.1021/es505846r. URL <http://pubs.acs.org/doi/10.1021/es505846r>.
- Quentin Renard, Raphaël Péliissier, B. R. Ramesh, and Narendran Kodandapani. Environmental susceptibility model for predicting forest fire occurrence in the Western Ghats of India. *International Journal of Wildland Fire*, 21(4):368, jul 2012. ISSN 1049-8001. doi: 10.1071/WF10109. URL <http://www.publish.csiro.au/?paper=WF10109>.

- 2495 April E. Reside, Jeremy VanDerWal, Alex Kutt, Ian Watson, and Stephen Williams. Fire regime shifts af-
 2496 fect bird species distributions. *Diversity and Distributions*, 18(3):213–225, mar 2012. ISSN 13669516. doi:
 2497 10.1111/j.1472-4642.2011.00818.x. URL <http://doi.wiley.com/10.1111/j.1472-4642.2011.00818.x>.
 2498 x.
- 2499 D. Riaño, S. L. Ustin, L. Usero, and M. A. Patricio. Estimation of Fuel Moisture Content Using Neural
 2500 Networks. pages 489–498. Springer, Berlin, Heidelberg, 2005. doi: 10.1007/11499305_50. URL http://link.springer.com/10.1007/11499305_50.
 2501 //link.springer.com/10.1007/11499305_50.
- 2502 Marco Tulio Ribeiro, Sameer Singh, and Carlos Guestrin. Model-Agnostic Interpretability of Machine
 2503 Learning. jun 2016. URL <http://arxiv.org/abs/1606.05386>.
- 2504 Rihan, Zhao, Zhang, Guo, Ying, Deng, and Li. Wildfires on the Mongolian Plateau: Identifying Drivers
 2505 and Spatial Distributions to Predict Wildfire Probability. *Remote Sensing*, 11(20):2361, oct 2019. ISSN
 2506 2072-4292. doi: 10.3390/rs11202361. URL <https://www.mdpi.com/2072-4292/11/20/2361>.
- 2507 David R. Roberts, Volker Bahn, Simone Ciuti, Mark S. Boyce, Jane Elith, Gurutzeta Guillera-Arroita, Sev-
 2508 erin Hauenstein, José J. Lahoz-Monfort, Boris Schröder, Wilfried Thuiller, David I. Warton, Brendan A.
 2509 Wintle, Florian Hartig, and Carsten F. Dormann. Cross-validation strategies for data with temporal,
 2510 spatial, hierarchical, or phylogenetic structure. *Ecography*, 40(8):913–929, aug 2017. ISSN 09067590.
 2511 doi: 10.1111/ecog.02881. URL <http://doi.wiley.com/10.1111/ecog.02881>.
- 2512 Marcos Rodrigues and Juan De la Riva. An insight into machine-learning algorithms to model human-
 2513 caused wildfire occurrence. *Environmental Modelling and Software*, 57:192–201, 2014. ISSN 13648152.
 2514 doi: 10.1016/j.envsoft.2014.03.003. URL <http://dx.doi.org/10.1016/j.envsoft.2014.03.003>.
- 2515 Marcos Rodrigues, Fermín Alcasena, and Cristina Vega-García. Modeling initial attack success of wildfire
 2516 suppression in Catalonia, Spain. *Science of The Total Environment*, 666:915–927, may 2019. ISSN
 2517 00489697. doi: 10.1016/j.scitotenv.2019.02.323. URL <https://linkinghub.elsevier.com/retrieve/pii/S0048969719308319>.
 2518 pii/S0048969719308319.
- 2519 Roque Rodriguez, Ana Cortés, Tomás Margalef, and Emilio Luque. An Adaptive System for Forest
 2520 Fire Behavior Prediction. In *2008 11th IEEE International Conference on Computational Science and
 2521 Engineering*, pages 275–282. IEEE, jul 2008. ISBN 978-0-7695-3193-9. doi: 10.1109/CSE.2008.15. URL
 2522 <http://ieeexplore.ieee.org/document/4578243/>.
- 2523 Roque Rodríguez, Ana Cortés, and Tomás Margalef. Injecting Dynamic Real-Time Data into a DDDAS
 2524 for Forest Fire Behavior Prediction. pages 489–499. Springer, Berlin, Heidelberg, 2009. doi: 10.1007/
 2525 978-3-642-01973-9_55. URL http://link.springer.com/10.1007/978-3-642-01973-9_55.
- 2526 Yuji Roh, Geon Heo, and Steven Euijong Whang. A Survey on Data Collection for Machine Learning: a
 2527 Big Data – AI Integration Perspective. *IEEE Transactions on Knowledge and Data Engineering*, pages
 2528 1–1, nov 2018. URL <http://arxiv.org/abs/1811.03402>.
- 2529 David Rolnick, Priya L. Donti, Lynn H. Kaack, Kelly Kochanski, Alexandre Lacoste, Kris Sankaran, An-
 2530 drew Slavin Ross, Nikola Milojevic-Dupont, Natasha Jaques, Anna Waldman-Brown, Alexandra Luc-
 2531 cioni, Tegan Maharaj, Evan D. Sherwin, S. Karthik Mukkavilli, Konrad P. Kording, Carla Gomes,
 2532 Andrew Y. Ng, Demis Hassabis, John C. Platt, Felix Creutzig, Jennifer Chayes, and Yoshua Bengio.
 2533 Tackling Climate Change with Machine Learning. jun 2019. URL <http://arxiv.org/abs/1906.05433>.
- 2534 J. Ruffault and F. Mouillot. How a new fire-suppression policy can abruptly reshape the fire-weather
 2535 relationship. *Ecosphere*, 6(10):art199, oct 2015. ISSN 2150-8925. doi: 10.1890/ES15-00182.1. URL
 2536 <http://doi.wiley.com/10.1890/ES15-00182.1>.

- 2537 Jakob Runge, Sebastian Bathiany, Erik Bollt, Gustau Camps-Valls, Dim Coumou, Ethan Deyle, Clark
 2538 Glymour, Marlene Kretschmer, Miguel D. Mahecha, Jordi Muñoz-Marí, Egbert H. van Nes, Jonas Peters,
 2539 Rick Quax, Markus Reichstein, Marten Scheffer, Bernhard Schölkopf, Peter Spirtes, George Sugihara, Jie
 2540 Sun, Kun Zhang, and Jakob Zscheischler. Inferring causation from time series in Earth system sciences.
 2541 *Nature Communications*, 10(1):1–13, dec 2019. ISSN 20411723. doi: 10.1038/s41467-019-10105-3.
- 2542 A. C. L. Sá, J. M. C. Pereira, M. J. P. Vasconcelos, J. M. N. Silva, N. Ribeiro, and A. Awasse. Assessing
 2543 the feasibility of sub-pixel burned area mapping in miombo woodlands of northern Mozambique using
 2544 MODIS imagery. *International Journal of Remote Sensing*, 24(8):1783–1796, jan 2003. ISSN 0143-
 2545 1161. doi: 10.1080/01431160210144750. URL [https://www.tandfonline.com/doi/full/10.1080/
 2546 01431160210144750](https://www.tandfonline.com/doi/full/10.1080/01431160210144750).
- 2547 Shruti Sachdeva, Tarunpreet Bhatia, and A. K. Verma. GIS-based evolutionary optimized Gradient
 2548 Boosted Decision Trees for forest fire susceptibility mapping. *Natural Hazards*, 92(3):1399–1418, jul
 2549 2018. ISSN 0921-030X. doi: 10.1007/s11069-018-3256-5. URL [http://link.springer.com/10.1007/
 2550 s11069-018-3256-5](http://link.springer.com/10.1007/s11069-018-3256-5).
- 2551 Y. Safi and A. Bouroumi. Prediction of forest fires using Artificial neural networks. *Applied Mathematical
 2552 Sciences*, 7(6):271–286, 2013. doi: 10.12988/ams.2013.13025.
- 2553 George E. Sakr, Imad H. Elhajj, George Mitri, and Uchechukwu C. Wejinya. Artificial intelligence for
 2554 forest fire prediction. In *IEEE/ASME International Conference on Advanced Intelligent Mechatronics,
 2555 AIM*, 2010. ISBN 9781424480319. doi: 10.1109/AIM.2010.5695809.
- 2556 George E. Sakr, Imad H. Elhajj, and George Mitri. Efficient forest fire occurrence prediction for
 2557 developing countries using two weather parameters. *Engineering Applications of Artificial Intelli-
 2558 gence*, 24(5):888–894, aug 2011. ISSN 0952-1976. doi: 10.1016/J.ENGAPPAI.2011.02.017. URL
 2559 <https://www.sciencedirect.com/science/article/abs/pii/S0952197611000418>.
- 2560 Jess San-Miguel-Ayanz, Ernst Schulte, Guido Schmuck, Andrea Camia, Peter Strobl, Giorgio Liberta, Cris-
 2561 tiano Giovando, Roberto Boca, Fernando Sedano, Pieter Kempeneers, Daniel McInerney, Ceri Withmore,
 2562 Sandra Santos de Oliveira, Marcos Rodrigues, Tracy Durrant, Paolo Corti, Friderike Oehler, Lara Vilar,
 2563 and Giuseppe Amatulli. Comprehensive Monitoring of Wildfires in Europe: The European Forest Fire
 2564 Information System (EFFIS). In *Approaches to Managing Disaster - Assessing Hazards, Emergencies
 2565 and Disaster Impacts*. InTech, mar 2012. doi: 10.5772/28441.
- 2566 L.A. Sanabria, X. Qin, J. Li, R.P. Cechet, and C. Lucas. Spatial interpolation of McArthur’s Forest Fire
 2567 Danger Index across Australia: Observational study. *Environmental Modelling & Software*, 50:37–50,
 2568 dec 2013. ISSN 1364-8152. doi: 10.1016/J.ENVSOF.2013.08.012. URL [https://www.sciencedirect.
 2569 com/science/article/pii/S1364815213001916?via%3Dihub](https://www.sciencedirect.com/science/article/pii/S1364815213001916?via%3Dihub).
- 2570 Onur Satir, Suha Berberoglu, and Cenk Donmez. Mapping regional forest fire probability using artificial
 2571 neural network model in a Mediterranean forest ecosystem. *Geomatics, Natural Hazards and Risk*, 7
 2572 (5):1645–1658, sep 2016. ISSN 1947-5705. doi: 10.1080/19475705.2015.1084541. URL [http://www.
 2573 tandfonline.com/doi/full/10.1080/19475705.2015.1084541](http://www.tandfonline.com/doi/full/10.1080/19475705.2015.1084541).
- 2574 Younes Oulad Sayad, Hajar Mousannif, and Hassan Al Moatassime. Predictive modeling of wildfires:
 2575 A new dataset and machine learning approach. *Fire Safety Journal*, 104:130–146, mar 2019. ISSN
 2576 03797112. doi: 10.1016/j.firesaf.2019.01.006.
- 2577 Daniel L. Schmoldt. Application of Artificial Intelligence to Risk Analysis for Forested Ecosystems. pages
 2578 49–74. Springer, Dordrecht, 2001. doi: 10.1007/978-94-017-2905-5_3. URL [http://link.springer.
 2579 com/10.1007/978-94-017-2905-5_3](http://link.springer.com/10.1007/978-94-017-2905-5_3).

- 2580 Frederic Paik Schoenberg. A NOTE ON THE CONSISTENT ESTIMATION OF SPATIAL-TEMPORAL
2581 POINT PROCESS PARAMETERS, 2016. URL <https://www.jstor.org/stable/24721302>.
- 2582 Gavin Shaddick and James V. Zidek. A case study in preferential sampling: Long term monitoring of air
2583 pollution in the UK. *Spatial Statistics*, 9:51–65, aug 2014. ISSN 2211-6753. doi: 10.1016/J.SPASTA.2014.
2584 03.008. URL <https://www.sciencedirect.com/science/article/abs/pii/S2211675314000219>.
- 2585 Chaopeng Shen. A Transdisciplinary Review of Deep Learning Research and Its Relevance for Water
2586 Resources Scientists. *Water Resources Research*, 54(11):8558–8593, nov 2018. ISSN 0043-1397. doi:
2587 10.1029/2018WR022643. URL <https://onlinelibrary.wiley.com/doi/abs/10.1029/2018WR022643>.
- 2588 Kirk R. Sherrill and William H. Romme. Spatial Variation in Postfire Cheatgrass: Dinosaur National
2589 Monument, USA. *Fire Ecology*, 8(2):38–56, aug 2012. ISSN 19339747. doi: 10.4996/fireecology.0802038.
2590 URL <http://fireecologyjournal.org/journal/abstract/?abstract=162>.
- 2591 Mengyun Shi, Fengying Xie, Yue Zi, and Jihao Yin. Cloud detection of remote sensing images by deep
2592 learning. In *International Geoscience and Remote Sensing Symposium (IGARSS)*, volume 2016-Novem,
2593 pages 701–704. Institute of Electrical and Electronics Engineers Inc., nov 2016. ISBN 9781509033324.
2594 doi: 10.1109/IGARSS.2016.7729176.
- 2595 Guruh Fajar Shidik and Khabib Mustofa. Predicting Size of Forest Fire Using Hybrid Model. pages
2596 316–327. Springer, Berlin, Heidelberg, 2014. doi: 10.1007/978-3-642-55032-4_31. URL [http://link.
2597 springer.com/10.1007/978-3-642-55032-4_{_}31](http://link.springer.com/10.1007/978-3-642-55032-4_{_}31).
- 2598 Albert J. Simard. Fire severity, changing scales, and how things hang together. *International Journal of*
2599 *Wildland Fire*, 1(1):23–34, 1991. ISSN 10498001. doi: 10.1071/WF9910023.
- 2600 Imas Sukaesih Sitanggang and Mohd Hasmadi Ismail. Classification model for hotspot occurrences using
2601 a decision tree method. *Geomatics, Natural Hazards and Risk*, 2(2):111–121, jun 2011. ISSN 19475705.
2602 doi: 10.1080/19475705.2011.565807.
- 2603 I.S. Sitanggang, R. Yaakob, N. Mustapha, and A.N. Ainuddin. Predictive Models for Hotspots Occurrence
2604 using Decision Tree Algorithms and Logistic Regression. *Journal of Applied Sciences*, 13(2):252–261,
2605 feb 2013. ISSN 18125654. doi: 10.3923/jas.2013.252.261. URL [http://www.scialert.net/abstract/
2606 ?doi=jas.2013.252.261](http://www.scialert.net/abstract/?doi=jas.2013.252.261).
- 2607 Natasa Skific and Jennifer Francis. Self-organizing maps: a powerful tool for the atmospheric sciences.
2608 *Applications of Self-Organizing Maps*, pages 251–268, 2012.
- 2609 W. R. Skinner, M. D. Flannigan, B. J. Stocks, D. L. Martell, B. M. Wotton, J. B. Todd, J. A. Mason,
2610 K. A. Logan, and E. M. Bosch. A 500 hPa synoptic wildland fire climatology for large Canadian forest
2611 fires, 1959-1996. *Theoretical and Applied Climatology*, 71(3-4):157–169, 2002. ISSN 0177798X. doi:
2612 10.1007/s007040200002.
- 2613 Hamdy Soliman, Komal Sudan, and Ashish Mishra. A smart forest-fire early detection sensory system:
2614 Another approach of utilizing wireless sensor and neural networks. In *2010 IEEE Sensors*, pages 1900–
2615 1904. IEEE, nov 2010. ISBN 978-1-4244-8170-5. doi: 10.1109/ICSENS.2010.5690033. URL [http:
2616 //ieeexplore.ieee.org/document/5690033/](http://ieeexplore.ieee.org/document/5690033/).
- 2617 Chao Song, Mei-Po Kwan, Weiguo Song, and Jiping Zhu. A Comparison between Spatial Econometric
2618 Models and Random Forest for Modeling Fire Occurrence. *Sustainability*, 9(5):819, may 2017. ISSN
2619 2071-1050. doi: 10.3390/su9050819. URL <http://www.mdpi.com/2071-1050/9/5/819>.
- 2620 Kalli Srinivasa, Nageswara Prasad, and S Ramakrishna. An Autonomous Forest Fire Detection System
2621 Based On Spatial Data Mining and Fuzzy Logic. Technical Report 12, 2008.

- 2622 B. J. Stocks and David L. Martell. Forest fire management expenditures in Canada: 1970-2013. *Forestry*
2623 *Chronicle*, 92(3):298–306, jun 2016. ISSN 00157546. doi: 10.5558/tfc2016-056.
- 2624 Daniela Stojanova, Andrej Kobler, Sašo Džeroski, and Katerina Taškova. LEARNING TO PREDICT
2625 FOREST FIRES WITH DIFFERENT DATA MINING TECHNIQUES. In *Conference on data mining*
2626 *and data warehouses (SiKDD 2006)*, pages 255–258, 2006. URL [http://www.academia.edu/download/](http://www.academia.edu/download/30570649/10.1.1.116.2555.pdf)
2627 [30570649/10.1.1.116.2555.pdf](http://www.academia.edu/download/30570649/10.1.1.116.2555.pdf).
- 2628 Daniela Stojanova, Andrej Kobler, Peter Ogrinc, Bernard Ženko, and Sašo Džeroski. Estimating the risk of
2629 fire outbreaks in the natural environment. *Data Mining and Knowledge Discovery*, 24(2):411–442, mar
2630 2012. ISSN 1384-5810. doi: 10.1007/s10618-011-0213-2. URL [http://link.springer.com/10.1007/](http://link.springer.com/10.1007/s10618-011-0213-2)
2631 [s10618-011-0213-2](http://link.springer.com/10.1007/s10618-011-0213-2).
- 2632 Jeremy Storer and Robert Green. PSO trained Neural Networks for predicting forest fire size: A comparison
2633 of implementation and performance. In *Proceedings of the International Joint Conference on Neural*
2634 *Networks*, volume 2016-Octob, pages 676–683. Institute of Electrical and Electronics Engineers Inc., oct
2635 2016. ISBN 9781509006199. doi: 10.1109/IJCNN.2016.7727265.
- 2636 Diana Stralberg, Xianli Wang, Marc-André Parisien, François-Nicolas Robinne, Péter Sólymos, C. Lisa
2637 Mahon, Scott E. Nielsen, and Erin M. Bayne. Wildfire-mediated vegetation change in boreal forests of
2638 Alberta, Canada. *Ecosphere*, 9(3):e02156, 2018. ISSN 2150-8925. doi: 10.1002/ecs2.2156.
- 2639 Carolin Strobl, Anne-Laure Boulesteix, Achim Zeileis, and Torsten Hothorn. Bias in random forest variable
2640 importance measures: Illustrations, sources and a solution. *BMC Bioinformatics*, 8(1):25, dec 2007.
2641 ISSN 1471-2105. doi: 10.1186/1471-2105-8-25. URL [https://bmcbioinformatics.biomedcentral.](https://bmcbioinformatics.biomedcentral.com/articles/10.1186/1471-2105-8-25)
2642 [com/articles/10.1186/1471-2105-8-25](https://bmcbioinformatics.biomedcentral.com/articles/10.1186/1471-2105-8-25).
- 2643 Esther D. Stroh, Matthew A. Struckhoff, Michael C. Stambaugh, and Richard P. Guyette. Fire and
2644 Climate Suitability for Woody Vegetation Communities in the South Central United States. *Fire Ecology*
2645 *2018 14:1*, 14(1):106–124, feb 2018. ISSN 1933-9747. doi: 10.4996/FIREECOLOGY.140110612. URL
2646 <https://fireecology.springeropen.com/articles/10.4996/fireecology.140110612>.
- 2647 Sriram Ganapathi Subramanian and Mark Crowley. Learning Forest Wildfire Dynamics from Satellite
2648 Images Using Reinforcement Learning. In *Conference on Reinforcement Learning and Decision Making*,
2649 page 5, 2017. URL <http://www.ospo.noaa.gov/Products/atmosphere/wind.html>.
- 2650 Sriram Ganapathi Subramanian and Mark Crowley. Using Spatial Reinforcement Learning to Build Forest
2651 Wildfire Dynamics Models From Satellite Images. *Frontiers in ICT*, 5:6, apr 2018. ISSN 2297-198X.
2652 doi: 10.3389/fict.2018.00006. URL [http://journal.frontiersin.org/article/10.3389/fict.2018.](http://journal.frontiersin.org/article/10.3389/fict.2018.00006/full)
2653 [00006/full](http://journal.frontiersin.org/article/10.3389/fict.2018.00006/full).
- 2654 AL Sullivan. A review of wildland fire spread modelling, 1990-present 3: Mathematical analogues and
2655 simulation models. *arXiv preprint arXiv:0706.4130*, 2007.
- 2656 Andrew L. Sullivan. Wildland surface fire spread modelling, 1990 - 2007. 3: Simulation and mathematical
2657 analogue models. *International Journal of Wildland Fire*, 18(4):387, 2009a. ISSN 1049-8001. doi:
2658 10.1071/wf06144.
- 2659 Andrew L. Sullivan. Wildland surface fire spread modelling, 1990 - 2007. 2: Empirical and quasi-empirical
2660 models. *International Journal of Wildland Fire*, 18(4):369, 2009b. ISSN 1049-8001. doi: 10.1071/
2661 wf06142.
- 2662 Andrew L. Sullivan. Wildland surface fire spread modelling, 1990 - 2007. 1: Physical and quasi-physical
2663 models. *International Journal of Wildland Fire*, 18(4):349, 2009c. ISSN 1049-8001. doi: 10.1071/wf06143.

- 2664 Brian L. Sullivan, Jocelyn L. Aycrigg, Jessie H. Barry, Rick E. Bonney, Nicholas Bruns, Caren B. Cooper,
 2665 Theo Damoulas, André A. Dhondt, Tom Dietterich, Andrew Farnsworth, Daniel Fink, John W. Fitz-
 2666 patrick, Thomas Fredericks, Jeff Gerbracht, Carla Gomes, Wesley M. Hochachka, Marshall J. Iliff,
 2667 Carl Lagoze, Frank A. La Sorte, Matthew Merrifield, Will Morris, Tina B. Phillips, Mark Reynolds,
 2668 Amanda D. Rodewald, Kenneth V. Rosenberg, Nancy M. Trautmann, Andrea Wiggins, David W. Win-
 2669 kler, Weng Keen Wong, Christopher L. Wood, Jun Yu, and Steve Kelling. The eBird enterprise: An
 2670 integrated approach to development and application of citizen science, jan 2014. ISSN 00063207.
- 2671 Alexander Y. Sun and Bridget R Scanlon. How can big data and machine learning benefit environment and
 2672 water management: A survey of methods, applications, and future directions. *Environmental Research*
 2673 *Letters*, apr 2019. ISSN 1748-9326. doi: 10.1088/1748-9326/ab1b7d. URL [http://iopscience.iop.
 2674 org/article/10.1088/1748-9326/ab1b7d](http://iopscience.iop.org/article/10.1088/1748-9326/ab1b7d).
- 2675 F. Sunar and C. Özkan. Forest fire analysis with remote sensing data. *International Journal of Remote*
 2676 *Sensing*, 22(12):2265–2277, jan 2001. ISSN 0143-1161. doi: 10.1080/01431160118510. URL [https:
 2677 //www.tandfonline.com/doi/full/10.1080/01431160118510](https://www.tandfonline.com/doi/full/10.1080/01431160118510).
- 2678 Richard S Sutton and Andrew G Barto. *Introduction to reinforcement learning*, volume 135. MIT press
 2679 Cambridge, 1998.
- 2680 Richard S Sutton and Andrew G Barto. *Reinforcement learning: An introduction*. MIT press, 2018.
- 2681 Alexandra D. Syphard, Jon E. Keeley, Avi Bar Massada, Teresa J. Brennan, and Volker C. Radeloff.
 2682 Housing Arrangement and Location Determine the Likelihood of Housing Loss Due to Wildfire. *PLoS*
 2683 *ONE*, 7(3):e33954, mar 2012. ISSN 1932-6203. doi: 10.1371/journal.pone.0033954. URL [http://dx.
 2684 plos.org/10.1371/journal.pone.0033954](http://dx.plos.org/10.1371/journal.pone.0033954).
- 2685 Alexandra D. Syphard, Avi Bar Massada, Van Butsic, and Jon E. Keeley. Land Use Planning and Wildfire:
 2686 Development Policies Influence Future Probability of Housing Loss. *PLoS ONE*, 8(8):e71708, aug 2013.
 2687 ISSN 1932-6203. doi: 10.1371/journal.pone.0071708. URL [https://dx.plos.org/10.1371/journal.
 2688 pone.0071708](https://dx.plos.org/10.1371/journal.pone.0071708).
- 2689 Alexandra D. Syphard, Van Butsic, Avi Bar-Massada, Jon E. Keeley, Jeff A. Tracey, and Robert N.
 2690 Fisher. Setting priorities for private land conservation in fire-prone landscapes: Are fire risk reduction
 2691 and biodiversity conservation competing or compatible objectives? *Ecology and Society*, 21(3):art2, jul
 2692 2016. ISSN 1708-3087. doi: 10.5751/ES-08410-210302. URL [http://www.ecologyandsociety.org/
 2693 vol21/iss3/art2/](http://www.ecologyandsociety.org/vol21/iss3/art2/).
- 2694 Alexandra D. Syphard, Heather Rustigian-Romsos, Michael Mann, Erin Conlisk, Max A. Moritz, and
 2695 David Ackerly. The relative influence of climate and housing development on current and projected
 2696 future fire patterns and structure loss across three California landscapes. *Global Environmental Change*,
 2697 56:41–55, may 2019. ISSN 0959-3780. doi: 10.1016/J.GLOENVCHA.2019.03.007. URL [https://www.
 2698 sciencedirect.com/science/article/pii/S0959378018313293](https://www.sciencedirect.com/science/article/pii/S0959378018313293).
- 2699 S. W. Taylor, Douglas G. Woolford, C. B. Dean, and David L. Martell. Wildfire Prediction to Inform Fire
 2700 Management: Statistical Science Challenges. *Statistical Science*, 28(4):586–615, 2013. ISSN 0883-4237.
 2701 doi: 10.1214/13-STS451.
- 2702 S.W. Taylor. Atmospheric cascades shape wildfire fire management decision spaces - a theory unifying fire
 2703 weather and fire management. *Submitted*, 2020.
- 2704 Mahyat Shafapour Tehrani, Simon Jones, Farzin Shabani, Francisco Martínez-Álvarez, and Dieu Tien
 2705 Bui. A novel ensemble modeling approach for the spatial prediction of tropical forest fire susceptibility
 2706 using LogitBoost machine learning classifier and multi-source geospatial data. *Theoretical and Applied*

- 2707 *Climatology*, pages 1–17, sep 2018. ISSN 0177-798X. doi: 10.1007/s00704-018-2628-9. URL <http://link.springer.com/10.1007/s00704-018-2628-9>.
2708
- 2709 Jonathan R. Thompson and Thomas A. Spies. Factors associated with crown damage following recurring
2710 mixed-severity wildfires and post-fire management in southwestern Oregon. *Landscape Ecology*, 25(5):
2711 775–789, may 2010. ISSN 0921-2973. doi: 10.1007/s10980-010-9456-3. URL <http://link.springer.com/10.1007/s10980-010-9456-3>.
2712
- 2713 Matthew P. Thompson and Dave E. Calkin. Uncertainty and risk in wildland fire management: A re-
2714 view. *Journal of Environmental Management*, 92(8):1895–1909, aug 2011. ISSN 0301-4797. doi:
2715 10.1016/J.JENVMAN.2011.03.015. URL <https://www.sciencedirect.com/science/article/pii/S0301479711000818>.
2716
- 2717 Dieu Tien Bui, Quang-Thanh Bui, Quoc-Phi Nguyen, Biswajeet Pradhan, Haleh Nampak, and Phan Trong
2718 Trinh. A hybrid artificial intelligence approach using GIS-based neural-fuzzy inference system and
2719 particle swarm optimization for forest fire susceptibility modeling at a tropical area. *Agricultural and*
2720 *Forest Meteorology*, 233:32–44, feb 2017. ISSN 0168-1923. doi: 10.1016/J.AGRFORMET.2016.11.002.
2721 URL <https://www.sciencedirect.com/science/article/pii/S0168192316304269>.
- 2722 Ahmed Toujani, Hammadi Achour, and Sami Faïz. Estimating Forest Fire Losses Using Stochastic
2723 Approach: Case Study of the Kroumiria Mountains (Northwestern Tunisia). *Applied Artificial*
2724 *Intelligence*, pages 1–25, sep 2018. ISSN 0883-9514. doi: 10.1080/08839514.2018.1514808. URL
2725 <https://www.tandfonline.com/doi/full/10.1080/08839514.2018.1514808>.
- 2726 James L. Tracy, Antonio Trabucco, A. Michelle Lawing, J. Tomasz Giermakowski, Maria Tchakerian,
2727 Gail M. Drus, and Robert N. Coulson. Random subset feature selection for ecological niche models
2728 of wildfire activity in Western North America. *Ecological Modelling*, 383:52–68, sep 2018. ISSN 0304-
2729 3800. doi: 10.1016/J.ECOLMODEL.2018.05.019. URL <https://www.sciencedirect.com/science/article/pii/S0304380018301868>.
2730
- 2731 Cordy Tymstra, Brian J. Stocks, Xinli Cai, and Mike D. Flannigan. Wildfire management in Canada:
2732 Review, challenges and opportunities. *Progress in Disaster Science*, page 100045, 2019. ISSN
2733 25900617. doi: 10.1016/j.pdisas.2019.100045. URL <https://linkinghub.elsevier.com/retrieve/pii/S2590061719300456>.
2734
- 2735 A.B. Utkin, Armando M Fernandes, Fernando Simões, and R. Vilar. Forest-fire detection by means of
2736 lidar. *Proceedings of IV International Conference on Forest Fire Research*, (1993):1–14, 2002.
- 2737 Giorgio Vacchiano, Cristiano Foderi, Roberta Berretti, Enrico Marchi, and Renzo Motta. Modeling
2738 anthropogenic and natural fire ignitions in an inner-alpine valley. *Natural Hazards and Earth Sys-*
2739 *tem Sciences*, 18(3):935–948, mar 2018. ISSN 1684-9981. doi: 10.5194/nhess-18-935-2018. URL
2740 <https://www.nat-hazards-earth-syst-sci.net/18/935/2018/>.
- 2741 D. Vakalis, H. Sarimveis, C. Kiranoudis, A. Alexandridis, and G. Bafas. A GIS based operational system
2742 for wildland fire crisis management I. Mathematical modelling and simulation. *Applied Mathematical*
2743 *Modelling*, 28(4):389–410, 2004. ISSN 0307904X. doi: 10.1016/j.apm.2003.10.005.
- 2744 Miguel Conrado Valdez, Kang-Tsung Chang, Chi-Farn Chen, Shou-Hao Chiang, and Jorge Luis Santos.
2745 Modelling the spatial variability of wildfire susceptibility in Honduras using remote sensing and geo-
2746 graphical information systems. *Geomatics, Natural Hazards and Risk*, 8(2):876–892, dec 2017. ISSN
2747 1947-5705. doi: 10.1080/19475705.2016.1278404. URL <https://www.tandfonline.com/doi/full/10.1080/19475705.2016.1278404>.
2748

- 2749 Ashley E. Van Beusekom, William A. Gould, A. Carolina Monmany, Azad Henareh Khalyani, Maya
 2750 Quiñones, Stephen J. Fain, Maria José Andrade-Núñez, and Grizelle González. Fire weather and
 2751 likelihood: characterizing climate space for fire occurrence and extent in Puerto Rico. *Climatic*
 2752 *Change*, 146(1-2):117–131, jan 2018. ISSN 0165-0009. doi: 10.1007/s10584-017-2045-6. URL <http://link.springer.com/10.1007/s10584-017-2045-6>.
 2753
- 2754 P. van Breugel, I. Friis, Sebsebe Demissew, Jens-Peter Barnekow Lillesø, and Roeland Kindt. Current and
 2755 Future Fire Regimes and Their Influence on Natural Vegetation in Ethiopia. *Ecosystems*, 19(2):369–386,
 2756 mar 2016. ISSN 1432-9840. doi: 10.1007/s10021-015-9938-x. URL <http://link.springer.com/10.1007/s10021-015-9938-x>.
 2757
- 2758 Ce Van Wagner. *Development and Structure of the Canadian Forest Fire Weather Index System*. 1987.
 2759 ISBN 0662151984. doi: 19927.
- 2760 Thomas Vandal, Evan Kodra, and Auroop R. Ganguly. Intercomparison of machine learning methods for
 2761 statistical downscaling: the case of daily and extreme precipitation. *Theoretical and Applied Climatology*,
 2762 pages 1–14, sep 2018. ISSN 0177-798X. doi: 10.1007/s00704-018-2613-3. URL <http://link.springer.com/10.1007/s00704-018-2613-3>.
 2763
- 2764 Christos Vasilakos, Kostas Kalabokidis, John Hatzopoulos, George Kallos, and Yiannis Matsinos. Integrat-
 2765 ing new methods and tools in fire danger rating. *International Journal of Wildland Fire*, 16(3):306–316,
 2766 2007. ISSN 10498001. doi: 10.1071/WF05091.
- 2767 Christos Vasilakos, Kostas Kalabokidis, John Hatzopoulos, and Ioannis Matsinos. Identifying wildland fire
 2768 ignition factors through sensitivity analysis of a neural network. *Natural Hazards*, 50(1):125–143, jul
 2769 2009. ISSN 0921-030X. doi: 10.1007/s11069-008-9326-3. URL <http://link.springer.com/10.1007/s11069-008-9326-3>.
 2770
- 2771 Daniel Vecín-Arias, Fernando Castedo-Dorado, Celestino Ordóñez, and José Ramón Rodríguez-Pérez. Bio-
 2772 physical and lightning characteristics drive lightning-induced fire occurrence in the central plateau
 2773 of the Iberian Peninsula. *Agricultural and Forest Meteorology*, 225:36–47, sep 2016. ISSN 0168-
 2774 1923. doi: 10.1016/J.AGRFORMET.2016.05.003. URL <https://www.sciencedirect.com/science/article/pii/S0168192316302593>.
 2775
- 2776 C. Vega-Garcia, B.S. Lee, P.M. Woodard, and S.J. Titus. Applying neural network technology to human-
 2777 caused wildfire occurrence prediction. *AI Applications*, pages 9–18, 1996. URL <https://cfs.nrcan.gc.ca/publications?id=18949>.
 2778
- 2779 O Viedma, L A Arroyo, R Mateo, A De Santis, and J M Moreno. EFFECTS OF ENVIRONMEN-
 2780 TAL PROPERTIES, BURNING CONDITIONS AND HUMAN-RELATED VARIABLES ON FIRE
 2781 SEVERITY DERIVED FROM LANDSAT TM IMAGES FOR A LARGE FIRE IN CENTRAL SPAIN.
 2782 In *Advances in Remote Sensing and GIS applications in Forest Fire Management From local to global*
 2783 *assessments*, page 157. 2011.
- 2784 Olga Viedma, Juan Quesada, Iván Torres, Angela De Santis, and José M. Moreno. Fire Severity in a Large
 2785 Fire in a Pinus pinaster Forest is Highly Predictable from Burning Conditions, Stand Structure, and
 2786 Topography. *Ecosystems*, 18(2):237–250, mar 2015. ISSN 1432-9840. doi: 10.1007/s10021-014-9824-y.
 2787 URL <http://link.springer.com/10.1007/s10021-014-9824-y>.
- 2788 Domingos Xavier Viegas, Karin L. Riley, Isaac C. Grenfell, Mark A. Finney, and Nicholas L.
 2789 Crookston. *Utilizing random forests imputation of forest plot data for landscape-level wild-*
 2790 *fire analyses*. Imprensa da Universidade de Coimbra, Coimbra, 2014. ISBN 978-989-26-0884-
 2791 6 (PDF). doi: 10.14195/978-989-26-0884-6.67. URL [https://digitalis.uc.pt/en/livro/
 2792 utilizing_{random}_{forests}_{imputation}_{forest}_{plot}_{data}_{landscape}_{level}_{wildfire-](https://digitalis.uc.pt/en/livro/utilizing_{random}_{forests}_{imputation}_{forest}_{plot}_{data}_{landscape}_{level}_{wildfire)

- 2793 Dinesh Babu Irulappa Pillai Vijayakumar, Frédéric Raulier, Pierre Y. Bernier, Sylvie Gauthier, Yves
 2794 Bergeron, and David Pothier. Lengthening the historical records of fire history over large areas of boreal
 2795 forest in eastern Canada using empirical relationships. *Forest Ecology and Management*, 347:30–39, jul
 2796 2015. ISSN 0378-1127. doi: 10.1016/J.FORECO.2015.03.011. URL [#}b0140](https://www.sciencedirect.com/science/article/pii/S0378112715001310).
 2797
- 2798 Dinesh Babu Irulappa Pillai Vijayakumar, Frédéric Raulier, Pierre Bernier, David Paré, Sylvie Gauthier,
 2799 Yves Bergeron, and David Pothier. Cover density recovery after fire disturbance controls landscape
 2800 aboveground biomass carbon in the boreal forest of eastern Canada. *Forest Ecology and Management*,
 2801 360:170–180, jan 2016. ISSN 0378-1127. doi: 10.1016/J.FORECO.2015.10.035. URL [https://www.
 2802 sciencedirect.com/science/article/pii/S0378112715005927?via{=}3Dihub](https://www.sciencedirect.com/science/article/pii/S0378112715005927?via{=}3Dihub).
- 2803 Lara Vilar, Israel Gómez, Javier Martínez-Vega, Pilar Echavarría, David Riaño, and M. Pilar Martín.
 2804 Multitemporal Modelling of Socio-Economic Wildfire Drivers in Central Spain between the 1980s and
 2805 the 2000s: Comparing Generalized Linear Models to Machine Learning Algorithms. *PLOS ONE*, 11(8):
 2806 e0161344, aug 2016. ISSN 1932-6203. doi: 10.1371/journal.pone.0161344. URL [http://dx.plos.org/
 2807 10.1371/journal.pone.0161344](http://dx.plos.org/10.1371/journal.pone.0161344).
- 2808 Yan Wang, Chunyu Yu, Ran Tu, and Yongming Zhang. Fire detection model in Tibet based on grey-fuzzy
 2809 neural network algorithm. *Expert Systems with Applications*, 38(8):9580–9586, aug 2011. ISSN 0957-
 2810 4174. doi: 10.1016/J.ESWA.2011.01.163. URL [https://www.sciencedirect.com/science/article/
 2811 pii/S0957417411001965](https://www.sciencedirect.com/science/article/pii/S0957417411001965).
- 2812 Yuanbin Wang, Langfei Dang, and Jieying Ren. Forest fire image recognition based on convolutional
 2813 neural network. *Journal of Algorithms & Computational Technology*, 13:174830261988768, jan 2019.
 2814 ISSN 1748-3026. doi: 10.1177/1748302619887689.
- 2815 Gregory L. Watson, Donatello Telesca, Colleen E. Reid, Gabriele G. Pfister, and Michael Jerrett. Machine
 2816 learning models accurately predict ozone exposure during wildfire events. *Environmental Pollution*, 254:
 2817 112792, nov 2019. ISSN 18736424. doi: 10.1016/j.envpol.2019.06.088.
- 2818 David H Wolpert. The lack of a priori distinctions between learning algorithms. *Neural computation*, 8(7):
 2819 1341–1390, 1996.
- 2820 Zhiwei Wu, Hong S. He, Jian Yang, Zhihua Liu, and Yu Liang. Relative effects of climatic and local
 2821 factors on fire occurrence in boreal forest landscapes of northeastern China. *Science of The Total
 2822 Environment*, 493:472–480, sep 2014. ISSN 0048-9697. doi: 10.1016/J.SCITOTENV.2014.06.011. URL
 2823 <https://www.sciencedirect.com/science/article/pii/S0048969714008547>.
- 2824 Zhiwei Wu, Hong S. He, Jian Yang, and Yu Liang. Defining fire environment zones in the boreal forests
 2825 of northeastern China. *Science of The Total Environment*, 518-519:106–116, jun 2015. ISSN 0048-9697.
 2826 doi: 10.1016/J.SCITOTENV.2015.02.063. URL [https://www.sciencedirect.com/science/article/
 2827 pii/S0048969715002065](https://www.sciencedirect.com/science/article/pii/S0048969715002065).
- 2828 Dexten D Z Xi, Stephen W Taylor, Douglas G Woolford, and C B Dean. Statistical Models of
 2829 Key Components of Wildfire Risk. 2019. doi: 10.1146/annurev-statistics-031017-100450. URL
 2830 www.annualreviews.org.
- 2831 Dao Wen Xie and Shi Liang Shi. Prediction for Burned Area of Forest Fires Based on SVM Model. *Applied
 2832 Mechanics and Materials*, 513-517:4084–4089, feb 2014. ISSN 1662-7482. doi: 10.4028/www.scientific.
 2833 net/AMM.513-517.4084. URL <https://www.scientific.net/AMM.513-517.4084>.
- 2834 Ying Xie and Minggang Peng. Forest fire forecasting using ensemble learning approaches. *Neural Computing
 2835 and Applications*, 31(9):4541–4550, sep 2019. ISSN 0941-0643. doi: 10.1007/s00521-018-3515-0. URL
 2836 <http://link.springer.com/10.1007/s00521-018-3515-0>.

- 2837 Jiayun Yao, Michael Brauer, Sean Raffuse, and Sarah B. Henderson. Machine Learning Approach To
2838 Estimate Hourly Exposure to Fine Particulate Matter for Urban, Rural, and Remote Populations during
2839 Wildfire Seasons. *Environmental Science & Technology*, 52(22):13239–13249, nov 2018a. ISSN 0013-
2840 936X. doi: 10.1021/acs.est.8b01921. URL <http://pubs.acs.org/doi/10.1021/acs.est.8b01921>.
- 2841 Jiayun Yao, Sean M. Raffuse, Michael Brauer, Grant J. Williamson, David M.J.S. Bowman, Fay H.
2842 Johnston, and Sarah B. Henderson. Predicting the minimum height of forest fire smoke within
2843 the atmosphere using machine learning and data from the CALIPSO satellite. *Remote Sensing of*
2844 *Environment*, 206:98–106, mar 2018b. ISSN 0034-4257. doi: 10.1016/J.RSE.2017.12.027. URL
2845 <https://www.sciencedirect.com/science/article/abs/pii/S003442571730603X>.
- 2846 Lingxiao Ying, Jie Han, Yongsheng Du, and Zehao Shen. Forest fire characteristics in China: Spatial pat-
2847 terns and determinants with thresholds. *Forest Ecology and Management*, 424:345–354, sep 2018. ISSN
2848 0378-1127. doi: 10.1016/J.FORECO.2018.05.020. URL [https://www.sciencedirect.com/science/](https://www.sciencedirect.com/science/article/pii/S0378112717317668)
2849 [article/pii/S0378112717317668](https://www.sciencedirect.com/science/article/pii/S0378112717317668).
- 2850 Adam M. Young, Philip E. Higuera, Paul A. Duffy, and Feng Sheng Hu. Climatic thresholds shape northern
2851 high-latitude fire regimes and imply vulnerability to future climate change. *Ecography*, 40(5):606–617,
2852 may 2017. ISSN 16000587. doi: 10.1111/ecog.02205.
- 2853 Adam M. Young, Philip E. Higuera, John T. Abatzoglou, Paul A. Duffy, and Feng Sheng Hu. Consequences
2854 of climatic thresholds for projecting fire activity and ecological change. *Global Ecology and Biogeography*,
2855 28(4):521–532, apr 2019. ISSN 14668238. doi: 10.1111/geb.12872.
- 2856 Bo Yu, Fang Chen, Bin Li, Li Wang, and Mingquan Wu. Fire Risk Prediction Using Remote Sensed
2857 Products: A Case of Cambodia. *Photogrammetric Engineering & Remote Sensing*, 83(1):19–25, jan 2017.
2858 ISSN 0099-1112. doi: 10.14358/PERS.83.1.19. URL [http://www.ingentaconnect.com/content/10.](http://www.ingentaconnect.com/content/10.14358/PERS.83.1.19)
2859 [14358/PERS.83.1.19](http://www.ingentaconnect.com/content/10.14358/PERS.83.1.19).
- 2860 Yong Poh Yu, Rosli Omar, Rhett D Harrison, Mohan Kumar Sammathuria, and Abdul Rahim Nik. Pattern
2861 clustering of forest fires based on meteorological variables and its classification using hybrid data mining
2862 methods. *Journal of Computational Biology and Bioinformatics Research*, 3(4):47–52, 2011. URL [http:](http://www.academicjournals.org/jcbbbr)
2863 [://www.academicjournals.org/jcbbbr](http://www.academicjournals.org/jcbbbr).
- 2864 Jie Yuan, Lidong Wang, Peng Wu, Chao Gao, and Lingqing Sun. Detection of Wildfires along Transmission
2865 Lines Using Deep Time and Space Features. *Pattern Recognition and Image Analysis*, 28(4):805–812,
2866 oct 2018. ISSN 15556212. doi: 10.1134/S1054661818040168.
- 2867 Bianca Zadrozny. Learning and evaluating classifiers under sample selection bias. In *Twenty-first interna-*
2868 *tional conference on Machine learning - ICML '04*, page 114, New York, New York, USA, 2004. ACM
2869 Press. ISBN 1581138285. doi: 10.1145/1015330.1015425. URL [http://portal.acm.org/citation.](http://portal.acm.org/citation.cfm?doid=1015330.1015425)
2870 [cfm?doid=1015330.1015425](http://portal.acm.org/citation.cfm?doid=1015330.1015425).
- 2871 Harold S.J. Zald and Christopher J. Dunn. Severe fire weather and intensive forest management increase fire
2872 severity in a multi-ownership landscape. *Ecological Applications*, 28(4):1068–1080, 2018. ISSN 19395582.
2873 doi: 10.1002/eap.1710.
- 2874 Olivier Zammit, Xavier Descombes, and Josiane Zerubia. Burnt area mapping using Support Vector
2875 Machines. *Forest Ecology and Management*, 234:S240, 2006. ISSN 03781127. doi: 10.1016/j.foreco.2006.
2876 08.269. URL <http://linkinghub.elsevier.com/retrieve/pii/S0378112706008097>.
- 2877 Bin Zhang, Wei Wei, Bingqian He, and Chuanlei Guo. Early wildfire smoke detection based on
2878 improved codebook model and convolutional neural networks. In Xudong Jiang and Jenq-
2879 Neng Hwang, editors, *Tenth International Conference on Digital Image Processing (ICDIP*

- 2880 2018), page 120. SPIE, aug 2018a. ISBN 9781510621992. doi: 10.1117/12.2502974. URL
 2881 [https://www.spiedigitallibrary.org/conference-proceedings-of-spie/10806/2502974/
 2882 Early-wildfire-smoke-detection-based-on-improved-codebook-model-and/10.1117/12.
 2883 2502974.full](https://www.spiedigitallibrary.org/conference-proceedings-of-spie/10806/2502974/Early-wildfire-smoke-detection-based-on-improved-codebook-model-and/10.1117/12.2502974.full).
- 2884 Guoli Zhang, Ming Wang, and Kai Liu. Forest Fire Susceptibility Modeling Using a Convolutional Neural
 2885 Network for Yunnan Province of China. *International Journal of Disaster Risk Science*, 10(3):1–18, sep
 2886 2019. ISSN 2095-0055. doi: 10.1007/s13753-019-00233-1. URL [http://link.springer.com/10.1007/
 2887 s13753-019-00233-1](http://link.springer.com/10.1007/s13753-019-00233-1).
- 2888 Qi Xing Zhang, Gao Hua Lin, Yong Ming Zhang, Gao Xu, and Jin Jun Wang. Wildland Forest Fire Smoke
 2889 Detection Based on Faster R-CNN using Synthetic Smoke Images. In *Procedia Engineering*, volume 211,
 2890 pages 441–446. Elsevier Ltd, 2018b. doi: 10.1016/j.proeng.2017.12.034.
- 2891 Qingjie Zhang, Jiaolong Xu, Liang Xu, and Haifeng Guo. Deep Convolutional Neural Networks for Forest
 2892 Fire Detection. Atlantis Press, 2016. doi: 10.2991/ifmeita-16.2016.105.
- 2893 Zhanqing Li, A. Khananian, R.H. Fraser, and J. Cihlar. Automatic detection of fire smoke using artificial
 2894 neural networks and threshold approaches applied to AVHRR imagery. *IEEE Transactions on Geoscience
 2895 and Remote Sensing*, 39(9):1859–1870, 2001. ISSN 01962892. doi: 10.1109/36.951076. URL [http:
 2896 //ieeexplore.ieee.org/document/951076/](http://ieeexplore.ieee.org/document/951076/).
- 2897 Feng Zhao, Chengquan Huang, and Zhiliang Zhu. Use of Vegetation Change Tracker and Support Vector
 2898 Machine to Map Disturbance Types in Greater Yellowstone Ecosystems in a 1984–2010 Landsat Time
 2899 Series. *IEEE Geoscience and Remote Sensing Letters*, 12(8):1650–1654, aug 2015. ISSN 1545-598X. doi:
 2900 10.1109/LGRS.2015.2418159. URL <http://ieeexplore.ieee.org/document/7088596/>.
- 2901 Jianhui Zhao, Zhong Zhang, Shizhong Han, Chengzhang Qu, Zhiyong Yuan, and Dengyi Zhang. SVM based
 2902 forest fire detection using static and dynamic features. *Computer Science and Information Systems*, 8
 2903 (3):821–841, 2011. ISSN 1820-0214. doi: 10.2298/csis101012030z.
- 2904 Yi Zhao, Jiale Ma, Xiaohui Li, and Jie Zhang. Saliency Detection and Deep Learning-Based Wildfire
 2905 Identification in UAV Imagery. *Sensors*, 18(3):712, feb 2018. ISSN 1424-8220. doi: 10.3390/s18030712.
 2906 URL <http://www.mdpi.com/1424-8220/18/3/712>.
- 2907 Zhong Zheng, Wei Huang, Songnian Li, and Yongnian Zeng. Forest fire spread simulating model us-
 2908 ing cellular automaton with extreme learning machine. *Ecological Modelling*, 348(May 2018):33–43,
 2909 2017. ISSN 03043800. doi: 10.1016/j.ecolmodel.2016.12.022. URL [http://dx.doi.org/10.1016/j.
 2910 ecolmodel.2016.12.022](http://dx.doi.org/10.1016/j.ecolmodel.2016.12.022).
- 2911 Yufei Zou, Susan M. O’Neill, Narasimhan K. Larkin, Ernesto C. Alvarado, Robert Solomon, Clifford Mass,
 2912 Yang Liu, M. Talat Odman, and Huizhong Shen. Machine Learning-Based Integration of High-Resolution
 2913 Wildfire Smoke Simulations and Observations for Regional Health Impact Assessment. *International
 2914 Journal of Environmental Research and Public Health*, 16(12):2137, jun 2019. ISSN 1660-4601. doi:
 2915 10.3390/ijerph16122137. URL <https://www.mdpi.com/1660-4601/16/12/2137>.
- 2916 K. Zwirgmaier, P. Papakosta, and D. Straub. Learning a Bayesian network model for predicting wildfire
 2917 behavior. In *ICOSSAR 2013*, 2013.