

Spring mires fed by hot artesian water in Kruger National Park, South Africa

A.P. Grootjans^{1,2} P-L. Grundling³, A. Grundling^{3,4}, A. Linström⁵, J. Engelbrecht⁵ & J.S. Price³

¹University of Groningen and ²Radboud University Nijmegen, The Netherlands; ³University of Waterloo, Canada; ⁴Agricultural Research Council (Institute for Soil, Climate and Water) and ⁵Golder Associates Africa, South Africa

SUMMARY

This article describes two spring mire complexes in the Kruger National Park (South Africa) that are fed by thermal water with a temperature of 37–42°C. The mires are small (1–20 m in diameter). The peat thickness is 1–2.5 m, of which 1–1.5 m is elevated above the surroundings. Some of the domes have dried out severely and show signs of erosion due to water flow and trampling by large animals. The mires lie in an almost straight line, supporting the hypothesis that the water originates from deep aquifers which discharge at geological faults. The long-term existence of these spring mire complexes may not be threatened because young stages of mire formation are present, but research to elucidate the timescales of peat development is needed to make a valid prognosis.

KEY WORDS: artesian water, erosion, hydrology, peat domes.

INTRODUCTION

Seasonal and temporal wetlands are common in Kruger National Park (KNP). Although 5–8% of the KNP area is covered by wetlands, most are associated with preferential water flows in river valleys, depressions or hill-slope seeps, and much less than 1% of them have more than 30 cm of accumulated peat. Peat-forming systems can exist here only if they have ground or surface water sources. Unlike other rural areas, the wetlands in Kruger are not used by humans.

Although mires that are fed by warm water are globally rare, according to Scott & Vogel (1983) there are 50 thermal springs in South Africa, some with organic deposits. An example is the Wonderkrater spring mound which is 3.5 m high, 8 m deep, 35,000 years old (McCarthy *et al.* 2010) and appears to be of ambient temperature, but there is a thermal spring with a water temperature of 36 ± 4°C in the close vicinity (Mazor & Vehagen 1983).

Thermal springs originate from rainwater that percolates into permeable rocks or joints and becomes heated underground. The water reaches substantial depths and its temperature increases at a rate of 2–3°C per 100 m due to the geothermal gradient (Press & Siever 1986). Convection currents in the groundwater are created when the heated water expands and rises. A thermal spring is sometimes formed when the heated groundwater in an aquifer encounters a fracture zone or an aquitard such as a dyke which forces it to rise to the surface (Olivier *et al.* 2008).

Growth of *Sphagnum* on hot wet soils (40–60°C)

has been reported from the sub-Antarctic (Ile St Paul; J. Whigham, pers. comm.). Thermal springs also occur in Iceland (Guðmundsson 2007), but the springs themselves are too hot to allow plant growth and thus peat accumulation, and bog vegetation types are found only where the water has cooled down sufficiently to allow vegetation to grow in surface water systems (A.J.M. Jansen, pers. comm.). A similar situation exists in Liard River Hot Springs Provincial Park in Canada, where the discharge of hot springs flows into a marsh (see also Weed 1889). We found no reports of discharging hot groundwater directly feeding a mire.

Local managers of KNP drew attention to the thermal mires because some of them had been badly damaged by wild animals and they wished to know if measures to protect the mires were needed. The objective of this paper is to present field data which were collected to support the formulation of hypotheses on the functioning of the mires, and to consider possible measures to protect them.

METHODS

Study area

Two hot spring mire areas in the north-eastern part of KNP were studied. The Malahlapanga spring mire (22° 53' 20.0" S, 31° 02' 25.8" E) is situated near the western boundary of KNP on a small tributary stream, close to its confluence with the Mphongolo River. The Mfayeni spring mire (23° 00' 47.4" S, 31° 14' 15.9" E) has a more easterly location (Figure 1).




Figure 1. Map of the northern part of Kruger National Park showing the locations of the Malahlapanga (No. 1) and Mfayeni (No. 2) thermal spring mires in relation to mean annual rainfall distribution, adapted from Venter & Gertenbach (1986).

The KNP has a sub-tropical climate with hot and humid summers and rather mild, dry winters (Venter & Gertenbach 1986). Temperatures range from $>40^{\circ}\text{C}$ in summer to about 0°C in winter. The mean annual temperature in the northern part of Kruger is 23.2°C ([/www.sanparks.org](http://www.sanparks.org)). Annual precipitation is 550–600 mm at Malahlapanga and 450–500 mm at Mfayeni (Gertenbach 1980), and the area has a strong soil water deficit (Venter & Gertenbach 1986, Mucina & Rutherford 2006).

Malahlapanga is underlain by the Goudplaats Gneiss. There is a major fault zone about 10 km to the north, with the Dzundwini and Nyunyani Faults striking from east to west (Figure 2). This fault zone dips northwards away from the site; but an offshoot from the southern Nyunyani Fault strikes roughly

north-south in line with, but stopping 2 km short of, Malahlapanga (Brandl 1981).

Gillson & Duffin (2010) carried out radiocarbon dating (AMS) on peat cores from both Malahlapanga and Mfayeni. From their research it can be concluded that the Mfayeni (110 cm depth) sequence began accumulating between 1640 and 1680 BP (before present). Malahlapanga is much older and in the Gillson & Duffin profile (72 cm) the peat began accumulating between 4940 and 4810 BP.

Data collection

The Malahlapanga mires were visited on 16 April 2009 (temperature measurements, peat coring) and in July 2010 (vegetation descriptions). Mfayeni was

visited on 27 April 2010 (temperature measurements, peat coring and water sampling) and in July 2010 (vegetation description).

The vegetation descriptions were restricted to dominant or important diagnostic species, including *Acacia grandicornuta*, *Bolboschoenus maritimus*, *Capparis tomentosa*, *Cyperus sexangularis*, *Echinochloa colona*, *Eustachys paspaloides*, *Fuirena pubescens*, *Hemarthria altissima*, *Ischaemum afrum*, *Hibiscus surattensis*, *Leptochloa chinensis*, *L. fusca*, *Miscanthus junceus*, *Paspalum distachyum*, *Pavonia* spp., *Phoenix reclinata*, *Phragmites australis*, *Pycnus polystachyos*, *Sesbania bispinosa*, *S. sesban*, *Sphaeranthus peduncularis*, *Thelypteris confluens* and *Trichoneura grandiglumis* (Burrows 1990, Retief & Herman 1997, Pooley 1998, Van Oudtshoorn 1999, Schmidt *et al.* 2002, Leistner 2005).

Temperature and electrical conductivity were measured with a combined EC/Temperature device capable of measuring directly in peat (WTW, LF96) attached to a two-metre-long probe. Peat depth was determined with a 'Russian type' (closed chamber) corer. Water samples were analysed at the Institute for Soil, Climate and Water (ARC.LNR) in Pretoria.

HCO_3^- was measured by titration; Cl^- and SO_4^{2-} were analysed using a Dionex Ionchromatograph; and Na, Ca, Mg, were measured using an ICP (Inductively Coupled Plasma) mass spectrometer (Wirsam).

RESULTS

Malahlapanga

At Malahlapanga, several thermal spring mires were present within an area of 4–6 hectares (Figures 2 and 5). The most spectacular view was of three small (*ca.* 200–500m²) peat cupolas rising 1–2 m above their surroundings. Smaller wetland patches (1–10 m²) were present between the larger mires. These were sparsely vegetated and damaged by animals such as elephant, rhinoceros and buffalo that had trampled the peat in order to get to the water. The tops of the two largest cupolas (A and C in Figure 5) were apparently dry and lacked wetland vegetation, which was present only at the bases of these domes. Two smaller and wetter cupolas lying between the two dried-out domes had many wetland species across their entire surfaces.




Figure 2. Schematic overview ('working model' *sensu* Grootjans *et al.* 2010) of the Malahlapanga wetland site showing four peat domes (A–D), as well as a mineral spring (E) that discharges warm groundwater (37°C) from a deep aquifer.

Peat coring in the largest cupola (A) showed that the total peat depth was approximately four metres, of which one metre lay below the level of the surrounding plateau. This cupola was covered by grasses and tall herbs with some large trees, and its edges were severely damaged by wild animal trampling. Although it had dried out considerably and a tree (*Acacia grandicornuta*) had established at the dry outer edge (Figures 3 and 5), some typical wetland species such as *Phragmites australis* and *Bolboschoenus maritimus* were still present but with low cover. Wetland vegetation occurred only at the base of the dome and along the eroding rivulets that transported spring water out of the wetland. The temperature of the water escaping from the base of the dome was *ca.* 33°C, but after a few hundred metres dropped to 23°C. Thus, it appeared that this dome was no longer fed by groundwater.

The cupola (D) on the right in Figure 2 was less damaged by animals but again there were no wetland species on its summit. It was covered by shrubs and grasses, with wetland species growing only around its base (Figure 5). A temperature profile to a depth of two metres showed that the warm water escaped at the base of the dome and did not reach the surface of the mire (Figure 6). On the other hand, a temperature profile in one of the two small and wet mires (C) showed that warm water rose to a height of 0.4 m below the top of the dome, although temperatures in the main rooting zone were close to air temperature. Species of healthy wetlands including *Thelypteris confluenta*, *Leptochloa fusca* and *Bolboschoenus maritimus* were present in the cupola vegetation, indicating active wetland processes (Figure 4).

The mires were surrounded by numerous ponds, which had evidently been created by wild animals. These ponds collected surface water that was discharging from the mires and from a small

unvegetated spring. The temperature of most pools was between 22°C and 23°C, which was about the same as the air temperature. The water discharging from a small spring pond was much warmer, with a temperature of 37°C. The electrical conductivity (EC), which is an indication of the total amount of dissolved minerals, showed that this water was rich in minerals (EC = 102 mS m⁻¹) but still in the range of fresh to slightly brackish water. The water emerging from the spring discharged into elongated muddy pools and watercourses, finally leaving the wetland *via* a small rivulet. The EC values of stagnant water in the muddy pools was very high (400–800 mS m⁻¹), most likely due to evaporation and consequent concentration of solutes.

It was evident in the field that the small mineral spring was the point of greatest groundwater discharge, since the surface water outflow was visibly greater than from the peat domes and other springs. We speculate that minor tectonic activity may have caused changes in flow patterns of the groundwater and that the groundwater discharge to the largest mires has decreased, causing compaction in the peat body. This may have forced groundwater to escape at the bases of the cupolas, instead of discharging at their summits. The concentrated groundwater outflow at the bases of the cupolas probably triggered further erosion around them.

The temperature profiles (Figure 6) indeed suggest that the upward flow of groundwater in the two largest peat domes has been blocked, forcing the water to escape sideways.

Elephant and other large animal activity is probably so severe in the larger domes because they have dried out. In order to reach the muddy water, animals dig and trample in the more central parts, probably just because they can. They appear to avoid the smaller wetter mire, probably because they do not want to risk sinking into the peat.

<i>Phragmites australis</i>							
<i>Bolboschoenus maritimus</i>							
<i>Miscanthus junceus</i>							
<i>Capparis tomentosa</i>							
<i>Echinochloa colona</i>							
<i>Leptochloa chinensis</i>							
<i>Hibiscus surattensis</i>							
<i>Capparis tomentosa</i>							
<i>Acacia grandicornuta</i>							
distance from edge of dome (m)	0		40			80	

Figure 3. Occurrence of some species on a transect (80 m long) running from the outer edge to the centre of the dried-out cupola (A) at Malahlapanga.

<i>Thelypteris confluenta</i>							
<i>Leptochloa fusca</i>							
<i>Hemarthria altissima</i>							
<i>Bolboschoenus maritimus</i>							
<i>Pycreus polystachyos</i>							
<i>Fuirena</i> spp.							
<i>Eustachys paspaloides</i>							
<i>Sesbania sesban</i>							
<i>Miscanthus junceus</i>							
<i>Pavonia</i> spp.							
distance from edge of dome (m)	0	15	30	45			

Figure 4. Occurrence of some species on a transect (45 m long) running from the outer edge to the centre of a wet cupola (C) at Malahlapanga.



Figure 5. Severely eroded peat cupolas in the Malahlapanga spring mire complex with trees (top left, A in Figure 2) or shrubs (top right, D in Figure 2) on their summits. A smaller peat dome (bottom left, C in Figure 2) is much wetter and covered with typical wetland species. The small pool (bottom right, E in Figure 2) is a spring that discharges groundwater at 37°C from a deep aquifer.




Figure 6. Temperature profiles in Malahlapanga for the two dried-out peat cupolas A (left) and D (right) and one cupola (C) whose summit is still wet (centre right). In Cupola C, warm water is rising higher in the profile than in the dry cupolas. Warm water is escaping at the base of the left-hand mire (A).



Figure 9. The large undisturbed cupola (top left) and the smaller highly mineralised cupolas (top right) at Mfayeni. The large peat dome is very wet and covered with typical wetland species (bottom left). The pools (bottom right) between cupolas and at the southern edge of the wetland are used by animals for water.

The concentrations of dissolved minerals in the groundwater discharging from the spring at Mfayeni were much higher than at Malahlapanga. The Mfayeni groundwater was somewhat brackish and also rich in calcium and sulphate (Table 1), suggesting dissolution of gypsum in deeper layers. The high sodium and chloride contents suggest dissolution of sea salt from marine deposits. The composition of this water was very similar to that of groundwater discharging from a spring located at Matiyovila, more than 700 m to the north-west. Thus, it appeared that these widely separated spring

systems were fed by the same groundwater source.

The area between the smaller cupolas and the large one, as well as areas south of the larger cupola, were heavily trampled by animals in search of water or in need of a mud bath. However, Mfayeni spring mire appeared generally to be in good condition although, judging from the occurrence of species such as *Sphaeranthus peduncularis* and *Trichoneura grandiglumis*, some parts had dried out. This is not unusual in a living spring mire system because groundwater discharge may shift position within the mire from time to time (Grootjans *et al.* 2005).

Table 1. Chemical composition of groundwater discharging from the spring in the large mire cupola at Mfayeni (0.5 m below the surface) and from another spring *ca.* 700 metres to the north-east (Matiyovila).

	Temp (°C)	EC (mS m ⁻¹)	Cl ⁻ (Mmol L ⁻¹)	SO ₄ ²⁻ (Mmol L ⁻¹)	HCO ₃ ⁻ (Mmol L ⁻¹)	Na ⁺ (Mmol L ⁻¹)	Ca ²⁺ (Mmol L ⁻¹)	Mg ²⁺ (Mmol L ⁻¹)
Mfayeni	42	407	20.2	6.8	0.3	21.2	16.6	0.01
Matiyovila	38.1	416	20.7	7.3	0.6	21.6	17.6	0.1




Figure 10. Temperature profiles ($^{\circ}\text{C}$) in the large cupola in Mfayeni. Warm water was rising very high in the peat dome and cooled down only in the top 0.2 m. We were unable to investigate the dry cupola(s) shown on the left-hand side with the temperature probe, due to the high penetration resistance of the soil.

DISCUSSION

Timescales

Spring mires are well known for their very dynamic histories, observed over timescales of centuries or even decades. Water flow to a spring mire may change through time, and some spring mires may be partly cut off from groundwater discharge as a result of tectonic activity. Also, water may shift to other areas within the spring mire itself, triggering drying out and erosion of some parts and stimulating peat growth in others (Wolejko *et al.* 1994). Such processes promote decomposition of the peat and thus complicate precise dating of the mire. Gillson & Duffin (2010) estimated that peat formation started *ca.* 5000 BP at Malahlapanga and *ca.* 1400 BP at Mfayeni. These are under-estimates for both mires, since the larger cupolas are much deeper than the maximum (110 cm) profile depth reported by these authors. Mfayeni, for instance, has a peat depth of 245 cm. The uppermost 50 cm consists almost entirely of water, but almost two metres of peat is present beneath. Some of the domes at Malahlapanga have similar thickness. Thus, a rather

conservative estimate, based on peat thickness and accumulation rates from Gillson & Duffin (2010), suggests that some domes at Malahlapanga may be 7,000–14,000 years old and the large dome at Mfayeni could be 3,000–4,000 years old.

Restoration measures needed?

The observations and measurements described above indicate that the best-developed spring mire cupolas have warm water high in the profile, whereas in very degraded ones groundwater escapes from the base of the mire or no groundwater is discharged at all. Trampling and erosion of the peat was evident in practically all cupolas, but was most severe in those where groundwater no longer reached the tops of the domes. On one occasion we identified ‘piping’, which is preferential water flow through underground erosion channels in the mire. Such structures can intercept rising groundwater before it reaches the top of the dome, and in most cases cannot be repaired (Koska & Stegmann 2001).

The question that now arises is whether the KNP managers should take measures to protect the mires

against wild animals and/or to restore them. Our preliminary conclusion is that restoration of the damaged spring cupolas is not an urgent priority at present because younger stages of dome formation are still present and animals do not go into the very wet and growing parts of these mires. Damage is most severe in cupolas that have lost their groundwater supplies, most likely due to natural causes, but we cannot state this with certainty. We do not think that the future of thermal spring mires in the KNP will be threatened unless the discharge of groundwater within the whole system decreases or even stops. However, some monitoring of the situation is recommended.

Future research

Research on the geology of this area is important in order to identify future threats to the warm-water-fed spring mires within the National Park. Further knowledge about the timescales of peat formation within the spring area could also inform prognoses for recovery of the existing spring cupolas and/or the development of new ones.

ACKNOWLEDGEMENTS

We wish to thank Freek Venter and Cathy Griever of SANPARKS (South African National Parks) for drawing our attention to the thermal spring mires, and the local manager and scientific services staff (Marius Renke and Jacob Mlangeni) of KNP for showing us the locations of the spring mires. Baps Snijdewind helped with the fieldwork at Mfayeni. We also thank Golder Associates Africa for supporting the second field excursion.

REFERENCES

Brandl, G. (1981) *Sheet 2230 (Messina)*. 1:250 000 Geological Map Series, Geological Survey South Africa.

Burrows, J.E. (1990) *Southern African Ferns and Fern Allies*. Frandsen Publishers, Sandton, South Africa, 359 pp.

Gertenbach, W.P.D. (1980) Rainfall patterns of the Kruger National Park. *Koedoe*, 23, 35–43.

Gillson, L. & Duffin, K.I. (2010) Thresholds of potential concern as benchmarks in the management of African savannahs. *Philosophical Transactions of the Royal Society B*, 362, 309–319.

Grootjans, A.P., Alserda, A., Bekker, R.E.M., Janáková, M., Kemmers, R.F., Madaras, M., Stanova, V., Ripka, J. Van Delft, B. & Wolejko,

L. (2005) Calcareous spring mires in Slovakia; jewels in the crown of the mire kingdom. In: Steiner, G.M. (ed.) *Mires, from Siberia to Tierra del Fuego*. Stapfia 85, Zugleich Kataloge der Oberösterreichischen Landesmuseen, N.S. 35, 97–115.

Grootjans, A.P., Iturraspe, R., Lanting, A., Fritz, C. & Joosten, H. (2010) Ecohydrological features of some contrasting mires in Tierra del Fuego, Argentina. *Mires and Peat*, 6(01), 1–15.

Guömundsson, A.T. (2007) *Living Earth; Outline of the Geology of Iceland*. Mál og Menning, Reykjavik, Iceland.

Koska, I. & Stegmann, H. (2001) Revitalisierung eines Quellmoorkomplexes am Serbitz-Oberlauf (Re-vitalisation of a spring mire complex in Upper Serbia). In: Succow, M. & Joosten, H. (eds.) *Landschaftsökologische Moorkunde*, Schweizerbart'sche Verlagsbuchhandlung, Stuttgart, 509–517 (in German).

Leistner, O.A. (2005) *Seed Plants of Southern Tropical Africa*. Southern African Botanical Diversity Network Report No. 26, 494 pp.

Mazor, E. & Vehagen, B.Th. (1983) Dissolved ions, stable and radioactive isotopes and noble gases in thermal waters of South Africa. *Journal of Hydrology*, 63, 315–329.

McCarthy, T.S., Ellery, W.N., Backwell, L., Marren, P., de Klerk, B., Tooth, S., Brandt, D. & Woodborne, S. (2010) The character, origin and palaeoenvironmental significance of the Wonderkrater spring mound, South Africa. *Journal of African Earth Sciences*, 58, 115–126.

Mucina, L. & Rutherford, M.C. (eds.) (2006) The vegetation of South Africa, Lesotho and Swaziland. *Strelitzia*, 19, South African National Biodiversity Institute, Pretoria.

Olivier, J.H.J., Van Niekerk, H.J. & Van der Walt, I.J. (2008) Physical and chemical characteristics of thermal springs in the Waterberg area in Limpopo Province, South Africa. *Water SA*, 34, 163–174.

Pooley, E. (1998) *A Field Guide to Wild Flowers Kwazulu-Natal and Eastern Region*. Natal Flora Publications Trust, 603 pp.

Press, F. & Siever, R. (1986) *Earth* (4th edition). Freeman and Co., New York, 626 pp.

Retief, E. & Herman, P.P.J. (1997) *Plants of the Northern Provinces of South Africa: Keys and Diagnostic Characters*. National Botanical Institute, Pretoria, South Africa, 681 pp.

Schmidt, E., Lotter, M. & McClelland (2002) *Trees and Shrubs of Mpumalanga and Kruger National Park*. Jacana, South Africa, 702 pp.

Scott, L. & Vogel, J.C. (1983) Late quaternary pollen profile from the Transvaal Highveld,

- South-Africa. *South African Journal of Science*, 79, 266–272.
- Van Oudtshoorn, F. (1999) *Guide to Grasses of Southern Africa*. Briza, Pretoria, South Africa, 301 pp.
- Venter, F.J. & Gertenbach, W.P.D. (1986) A cursory review of the climate and vegetation of the Kruger National Park. *Koedoe*, 29, 139–148.
- Weed, W.H. (1889) The vegetation of hot springs. *American Naturalist*, 23, 394–398.
- Wolejko, L., Aggenbach, C., Van Diggelen, R. & Grootjans, A.P. (1994) Vegetation and hydrology in a spring mire complex in western Pomerania, Poland. *Proceedings of the Royal Academy of Sciences of the Netherlands*, 97, 219–245.
- Submitted 05 Nov 2010, revision 26 Nov 2010*
Editor: Olivia Bragg
-

Author for correspondence:

Dr Ab P. Grootjans, Centre for Energy and Environmental Studies (IVEM), University of Groningen, Nijenborgh 4, 9747 AG, Groningen, The Netherlands. Tel: 003150268160; E-mail: A.P.Grootjans@rug.nl